



$$J = 0$$

In the following  $H^0$  refers to the signal that has been discovered in the Higgs searches. Whereas the observed signal is labeled as a spin 0 particle and is called a Higgs Boson, the detailed properties of  $H^0$  and its role in the context of electroweak symmetry breaking need to be further clarified. These issues are addressed by the measurements listed below.

Concerning mass limits and cross section limits that have been obtained in the searches for neutral and charged Higgs bosons, see the sections “Searches for Neutral Higgs Bosons” and “Searches for Charged Higgs Bosons ( $H^\pm$  and  $H^{\pm\pm}$ )”, respectively.

### $H^0$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
<b>125.10±0.14 OUR AVERAGE</b>			
124.86±0.27	<sup>1</sup> AABOUD	18BMATLS	$pp$ , 13 TeV, 36.1 fb <sup>-1</sup> , $\gamma\gamma, ZZ^* \rightarrow 4\ell$
125.26±0.20±0.08	<sup>2</sup> SIRUNYAN	17AV CMS	$pp$ , 13 TeV, $ZZ^* \rightarrow 4\ell$
125.09±0.21±0.11	<sup>1,3</sup> AAD	15B LHC	$pp$ , 7, 8 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
124.79±0.37	<sup>4</sup> AABOUD	18BMATLS	$pp$ , 13 TeV, 36.1 fb <sup>-1</sup> , $ZZ^* \rightarrow 4\ell$
124.93±0.40	<sup>5</sup> AABOUD	18BMATLS	$pp$ , 13 TeV, 36.1 fb <sup>-1</sup> , $\gamma\gamma$
124.97±0.24	<sup>1,6</sup> AABOUD	18BMATLS	$pp$ , 7, 8, 13 TeV, $\gamma\gamma$ , $ZZ^* \rightarrow 4\ell$
125.07±0.25±0.14	<sup>3</sup> AAD	15B LHC	$pp$ , 7, 8 TeV, $\gamma\gamma$
125.15±0.37±0.15	<sup>3</sup> AAD	15B LHC	$pp$ , 7, 8 TeV, $ZZ^* \rightarrow 4\ell$
126.02±0.43±0.27	AAD	15B ATLS	$pp$ , 7, 8 TeV, $\gamma\gamma$
124.51±0.52±0.04	AAD	15B ATLS	$pp$ , 7, 8 TeV, $ZZ^* \rightarrow 4\ell$
125.59±0.42±0.17	AAD	15B CMS	$pp$ , 7, 8 TeV, $ZZ^* \rightarrow 4\ell$
125.02 <sup>+0.26+0.14</sup> <sub>-0.27-0.15</sub>	<sup>7</sup> KHACHATRY...15AM	CMS	$pp$ , 7, 8 TeV
125.36±0.37±0.18	<sup>1,8</sup> AAD	14W ATLS	$pp$ , 7, 8 TeV
125.98±0.42±0.28	<sup>8</sup> AAD	14W ATLS	$pp$ , 7, 8 TeV, $\gamma\gamma$
124.51±0.52±0.06	<sup>8</sup> AAD	14W ATLS	$pp$ , 7, 8 TeV, $ZZ^* \rightarrow 4\ell$
125.6 ±0.4 ±0.2	<sup>9</sup> CHATRCHYAN 14AA	CMS	$pp$ , 7, 8 TeV, $ZZ^* \rightarrow 4\ell$
122 ±7	<sup>10</sup> CHATRCHYAN 14K	CMS	$pp$ , 7, 8 TeV, $\tau\tau$
124.70±0.31±0.15	<sup>11</sup> KHACHATRY...14P	CMS	$pp$ , 7, 8 TeV, $\gamma\gamma$
125.5 ±0.2 <sup>+0.5</sup> <sub>-0.6</sub>	<sup>1,12</sup> AAD	13AK ATLS	$pp$ , 7, 8 TeV
126.8 ±0.2 ±0.7	<sup>12</sup> AAD	13AK ATLS	$pp$ , 7, 8 TeV, $\gamma\gamma$
124.3 <sup>+0.6+0.5</sup> <sub>-0.5-0.3</sub>	<sup>12</sup> AAD	13AK ATLS	$pp$ , 7, 8 TeV, $ZZ^* \rightarrow 4\ell$
125.8 ±0.4 ±0.4	<sup>1,13</sup> CHATRCHYAN 13J	CMS	$pp$ , 7, 8 TeV
126.2 ±0.6 ±0.2	<sup>13</sup> CHATRCHYAN 13J	CMS	$pp$ , 7, 8 TeV, $ZZ^* \rightarrow 4\ell$
126.0 ±0.4 ±0.4	<sup>1,14</sup> AAD	12AI ATLS	$pp$ , 7, 8 TeV
125.3 ±0.4 ±0.5	<sup>1,15</sup> CHATRCHYAN 12N	CMS	$pp$ , 7, 8 TeV

<sup>1</sup> Combined value from  $\gamma\gamma$  and  $ZZ^* \rightarrow 4\ell$  final states.

- <sup>2</sup> SIRUNYAN 17AV use  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$  with  $H^0 \rightarrow ZZ^* \rightarrow 4\ell$  where  $\ell = e, \mu$ .
- <sup>3</sup> ATLAS and CMS data are fitted simultaneously.
- <sup>4</sup> AABOUD 18BM use  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$  with  $H^0 \rightarrow ZZ^* \rightarrow 4\ell$  where  $\ell = e, \mu$ .
- <sup>5</sup> AABOUD 18BM use  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$  with  $H^0 \rightarrow \gamma\gamma$ .
- <sup>6</sup> AABOUD 18BM combine 13 TeV results with 7 and 8 TeV results. Other combined results are summarized in their Fig. 4.
- <sup>7</sup> KHACHATRYAN 15AM use up to  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and up to  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ .
- <sup>8</sup> AAD 14W use  $4.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $20.3 \text{ fb}^{-1}$  at 8 TeV.
- <sup>9</sup> CHATRCHYAN 14AA use  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ .
- <sup>10</sup> CHATRCHYAN 14K use  $4.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ .
- <sup>11</sup> KHACHATRYAN 14P use  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ .
- <sup>12</sup> AAD 13AK use  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $20.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . Superseded by AAD 14W.
- <sup>13</sup> CHATRCHYAN 13J use  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $12.2 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ .
- <sup>14</sup> AAD 12AI obtain results based on  $4.6\text{--}4.8 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $5.8\text{--}5.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . An excess of events over background with a local significance of  $5.9 \sigma$  is observed at  $m_{H^0} = 126 \text{ GeV}$ . See also AAD 12DA.
- <sup>15</sup> CHATRCHYAN 12N obtain results based on  $4.9\text{--}5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $5.1\text{--}5.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . An excess of events over background with a local significance of  $5.0 \sigma$  is observed at about  $m_{H^0} = 125 \text{ GeV}$ . See also CHATRCHYAN 12BY and CHATRCHYAN 13Y.

## $H^0$ SPIN AND $CP$ PROPERTIES

The observation of the signal in the  $\gamma\gamma$  final state rules out the possibility that the discovered particle has spin 1, as a consequence of the Landau-Yang theorem. This argument relies on the assumptions that the decaying particle is an on-shell resonance and that the decay products are indeed two photons rather than two pairs of boosted photons, which each could in principle be misidentified as a single photon.

Concerning distinguishing the spin 0 hypothesis from a spin 2 hypothesis, some care has to be taken in modelling the latter in order to ensure that the discriminating power is actually based on the spin properties rather than on unphysical behavior that may affect the model of the spin 2 state.

Under the assumption that the observed signal consists of a single state rather than an overlap of more than one resonance, it is sufficient to discriminate between distinct hypotheses in the spin analyses. On the other hand, the determination of the  $CP$  properties is in general much more difficult since in principle the observed state could consist of any admixture of  $CP$ -even and  $CP$ -odd components. As a first step, the compatibility of the data with distinct hypotheses of pure  $CP$ -even and pure  $CP$ -odd states with different spin assignments has been investigated. In order to treat the case of a possible mixing of different  $CP$  states, certain cross section ratios are considered. Those cross section ratios need to be distinguished from the amount of mixing between a  $CP$ -even and a  $CP$ -odd state, as the cross section ratios depend

in addition also on the coupling strengths of the  $CP$ -even and  $CP$ -odd components to the involved particles. A small relative coupling implies a small sensitivity of the corresponding cross section ratio to effects of  $CP$  mixing.

VALUE	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
1	AABOUD 18AJ	ATLS	$H^0 \rightarrow ZZ^* \rightarrow 4l$ ( $l = e, \mu$ ), 13TeV
2	SIRUNYAN 17AM	CMS	$pp \rightarrow H^0 + \geq 2j$ , $H^0 \rightarrow 4l$ ( $l = e, \mu$ )
3	AAD 16	ATLS	$H^0 \rightarrow \gamma\gamma$
4	AAD 16BL	ATLS	$pp \rightarrow H^0 jjX$ (VBF), $H^0 \rightarrow \tau\tau$ , 8 TeV
5	KHACHATRYAN 16AB	CMS	$pp \rightarrow WH^0, ZH^0, H^0 \rightarrow b\bar{b}$ , 8 TeV
6	AAD 15AX	ATLS	$H^0 \rightarrow WW^*$
7	AAD 15CI	ATLS	$H^0 \rightarrow ZZ^*, WW^*, \gamma\gamma$
8	AALTONEN 15	TEVA	$p\bar{p} \rightarrow WH^0, ZH^0, H^0 \rightarrow b\bar{b}$
9	AALTONEN 15B	CDF	$p\bar{p} \rightarrow WH^0, ZH^0, H^0 \rightarrow b\bar{b}$
10	KHACHATRYAN 15Y	CMS	$H^0 \rightarrow 4l, WW^*, \gamma\gamma$
11	ABAZOV 14F	D0	$p\bar{p} \rightarrow WH^0, ZH^0, H^0 \rightarrow b\bar{b}$
12	CHATRCHYAN 14AA	CMS	$H^0 \rightarrow ZZ^*$
13	CHATRCHYAN 14G	CMS	$H^0 \rightarrow WW^*$
14	KHACHATRYAN 14P	CMS	$H^0 \rightarrow \gamma\gamma$
15	AAD 13AJ	ATLS	$H^0 \rightarrow \gamma\gamma, ZZ^* \rightarrow 4l, WW^* \rightarrow l\nu l\nu$
16	CHATRCHYAN 13J	CMS	$H^0 \rightarrow ZZ^* \rightarrow 4l$

<sup>1</sup>AABOUD 18AJ study the tensor structure of the Higgs boson couplings using an effective Lagrangian using  $36.1 \text{ fb}^{-1}$  of  $pp$  collision data at  $E_{\text{cm}} = 13 \text{ TeV}$ . Constraints are set on the non-Standard-Model  $CP$ -even and  $CP$ -odd couplings to  $Z$  bosons and on the  $CP$ -odd coupling to gluons. See their Figs. 9 and 10, and Tables 10 and 11.

<sup>2</sup>SIRUNYAN 17AM constrain anomalous couplings of the Higgs boson with  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$ ,  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ , and  $38.6 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Table 3 and Fig. 3, which show 68% CL and 95% CL intervals. A  $CP$  violation parameter  $f_{a3}$  is set to be  $f_{a3}\cos(\phi_{a3}) = [-0.38, 0.46]$  at 95% CL ( $\phi_{a3} = 0$  or  $\pi$ ).

<sup>3</sup>AAD 16 study  $H^0 \rightarrow \gamma\gamma$  with an effective Lagrangian including  $CP$  even and odd terms in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . The data is consistent with the expectations for the Higgs boson of the Standard Model. Limits on anomalous couplings are also given.

<sup>4</sup>AAD 16BL study VBF  $H^0 \rightarrow \tau\tau$  with an effective Lagrangian including a  $CP$  odd term in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . The measurement is consistent with the expectation of the Standard Model. The  $CP$ -mixing parameter  $\tilde{d}$  (a dimensionless coupling  $\tilde{d} = -(m_W^2/\Lambda^2)f_{\tilde{W}W}$ ) is constrained to the interval of  $(-0.11, 0.05)$  at 68% CL under the assumption of  $\tilde{d} = \tilde{d}_B$ .

<sup>5</sup>KHACHATRYAN 16AB search for anomalous pseudoscalar couplings of the Higgs boson to  $W$  and  $Z$  with  $18.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Table 5 and Figs 5 and 6 for limits on possible anomalous pseudoscalar coupling parameters.

<sup>6</sup>AAD 15AX compare the  $J^{CP} = 0^+$  Standard Model assignment with other  $J^{CP}$  hypotheses in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ , using the process  $H^0 \rightarrow WW^* \rightarrow e\nu\mu\nu$ .  $2^+$  hypotheses are excluded at 84.5–99.4%CL,  $0^-$  at 96.5%CL,  $0^+$  (field strength coupling) at 70.8%CL. See their Fig. 19 for limits on possible  $CP$  mixture parameters.

<sup>7</sup>AAD 15CI compare the  $J^{CP} = 0^+$  Standard Model assignment with other  $J^{CP}$  hypotheses in  $4.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ ,

- using the processes  $H^0 \rightarrow ZZ^* \rightarrow 4\ell$ ,  $H^0 \rightarrow \gamma\gamma$  and combine with AAD 15AX data.  $0^+$  (field strength coupling),  $0^-$  and several  $2^+$  hypotheses are excluded at more than 99.9% CL. See their Tables 7–9 for limits on possible  $CP$  mixture parameters.
- <sup>8</sup> AALTONEN 15 combine AALTONEN 15B and ABAZOV 14F data. An upper limit of 0.36 of the Standard Model production rate at 95% CL is obtained both for a  $0^-$  and a  $2^+$  state. Assuming the SM event rate, the  $J^{CP} = 0^-$  ( $2^+$ ) hypothesis is excluded at the  $5.0\sigma$  ( $4.9\sigma$ ) level.
- <sup>9</sup> AALTONEN 15B compare the  $J^{CP} = 0^+$  Standard Model assignment with other  $J^{CP}$  hypotheses in  $9.45 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ , using the processes  $ZH^0 \rightarrow \ell\ell b\bar{b}$ ,  $WH^0 \rightarrow \ell\nu b\bar{b}$ , and  $ZH^0 \rightarrow \nu\nu b\bar{b}$ . Bounds on the production rates of  $0^-$  and  $2^+$  (graviton-like) states are set, see their tables II and III.
- <sup>10</sup> KHACHATRYAN 15Y compare the  $J^{CP} = 0^+$  Standard Model assignment with other  $J^{CP}$  hypotheses in up to  $5.1 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and up to  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ , using the processes  $H^0 \rightarrow 4\ell$ ,  $H^0 \rightarrow WW^*$ , and  $H^0 \rightarrow \gamma\gamma$ .  $0^-$  is excluded at 99.98% CL, and several  $2^+$  hypotheses are excluded at more than 99% CL. Spin 1 models are excluded at more than 99.999% CL in  $ZZ^*$  and  $WW^*$  modes. Limits on anomalous couplings and several cross section fractions, treating the case of  $CP$ -mixed states, are also given.
- <sup>11</sup> ABAZOV 14F compare the  $J^{CP} = 0^+$  Standard Model assignment with  $J^{CP} = 0^-$  and  $2^+$  (graviton-like coupling) hypotheses in up to  $9.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . They use kinematic correlations between the decay products of the vector boson and the Higgs boson in the final states  $ZH \rightarrow \ell\ell b\bar{b}$ ,  $WH \rightarrow \ell\nu b\bar{b}$ , and  $ZH \rightarrow \nu\nu b\bar{b}$ . The  $0^-$  ( $2^+$ ) hypothesis is excluded at 97.6% CL (99.0% CL). In order to treat the case of a possible mixture of a  $0^+$  state with another  $J^{CP}$  state, the cross section fractions  $f_X = \sigma_X / (\sigma_{0^+} + \sigma_X)$  are considered, where  $X = 0^-, 2^+$ . Values for  $f_{0^-}$  ( $f_{2^+}$ ) above 0.80 (0.67) are excluded at 95% CL under the assumption that the total cross section is that of the SM Higgs boson.
- <sup>12</sup> CHATRCHYAN 14AA compare the  $J^{CP} = 0^+$  Standard Model assignment with various  $J^{CP}$  hypotheses in  $5.1 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ .  $J^{CP} = 0^-$  and  $1^\pm$  hypotheses are excluded at 99% CL, and several  $J = 2$  hypotheses are excluded at 95% CL. In order to treat the case of a possible mixture of a  $0^+$  state with another  $J^{CP}$  state, the cross section fraction  $f_{a3} = |a_3|^2 \sigma_3 / (|a_1|^2 \sigma_1 + |a_2|^2 \sigma_2 + |a_3|^2 \sigma_3)$  is considered, where the case  $a_3 = 1$ ,  $a_1 = a_2 = 0$  corresponds to a pure  $CP$ -odd state. Assuming  $a_2 = 0$ , a value for  $f_{a3}$  above 0.51 is excluded at 95% CL.
- <sup>13</sup> CHATRCHYAN 14G compare the  $J^{CP} = 0^+$  Standard Model assignment with  $J^{CP} = 0^-$  and  $2^+$  (graviton-like coupling) hypotheses in  $4.9 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.4 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . Varying the fraction of the production of the  $2^+$  state via  $g\bar{g}$  and  $q\bar{q}$ ,  $2^+$  hypotheses are disfavored at CL between 83.7 and 99.8%. The  $0^-$  hypothesis is disfavored against  $0^+$  at the 65.3% CL.
- <sup>14</sup> KHACHATRYAN 14P compare the  $J^{CP} = 0^+$  Standard Model assignment with a  $2^+$  (graviton-like coupling) hypothesis in  $5.1 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . Varying the fraction of the production of the  $2^+$  state via  $g\bar{g}$  and  $q\bar{q}$ ,  $2^+$  hypotheses are disfavored at CL between 71 and 94%.
- <sup>15</sup> AAD 13AJ compare the spin 0,  $CP$ -even hypothesis with specific alternative hypotheses of spin 0,  $CP$ -odd, spin 1,  $CP$ -even and  $CP$ -odd, and spin 2,  $CP$ -even models using the Higgs boson decays  $H \rightarrow \gamma\gamma$ ,  $H \rightarrow ZZ^* \rightarrow 4\ell$  and  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  and combinations thereof. The data are compatible with the spin 0,  $CP$ -even hypothesis, while all other tested hypotheses are excluded at confidence levels above 97.8%.
- <sup>16</sup> CHATRCHYAN 13J study angular distributions of the lepton pairs in the  $ZZ^*$  channel where both  $Z$  bosons decay to  $e$  or  $\mu$  pairs. Under the assumption that the observed

particle has spin 0, the data are found to be consistent with the pure  $CP$ -even hypothesis, while the pure  $CP$ -odd hypothesis is disfavored.

## $H^0$ DECAY WIDTH

The total decay width for a light Higgs boson with a mass in the observed range is not expected to be directly observable at the LHC. For the case of the Standard Model the prediction for the total width is about 4 MeV, which is three orders of magnitude smaller than the experimental mass resolution. There is no indication from the results observed so far that the natural width is broadened by new physics effects to such an extent that it could be directly observable. Furthermore, as all LHC Higgs channels rely on the identification of Higgs decay products, the total Higgs width cannot be measured indirectly without additional assumptions. The different dependence of on-peak and off-peak contributions on the total width in Higgs decays to  $ZZ^*$  and interference effects between signal and background in Higgs decays to  $\gamma\gamma$  can provide additional information in this context. Constraints on the total width from the combination of on-peak and off-peak contributions in Higgs decays to  $ZZ^*$  rely on the assumption of equal on- and off-shell effective couplings. Without an experimental determination of the total width or further theoretical assumptions, only ratios of couplings can be determined at the LHC rather than absolute values of couplings.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.0144	95	<sup>1</sup> AABOUD	18BP ATLS	$pp$ , 13 TeV, $ZZ \rightarrow 4\ell, 2\ell 2\nu$
<1.10	95	<sup>2</sup> SIRUNYAN	17AV CMS	$pp$ , 13 TeV, $ZZ^* \rightarrow 4\ell$
<b>&lt;0.013</b>	95	<sup>3</sup> KHACHATRYAN...16BA	CMS	$pp$ , 7, 8 TeV, $ZZ^{(*)}, WW^{(*)}$
<1.7	95	<sup>4</sup> KHACHATRYAN...15AM	CMS	$pp$ , 7, 8 TeV
>3.5 $\times 10^{-12}$	95	<sup>5</sup> KHACHATRYAN...15BA	CMS	$pp$ , 7, 8 TeV, flight distance
<5.0	95	<sup>6</sup> AAD	14W ATLS	$pp$ , 7, 8 TeV, $\gamma\gamma$
<2.6	95	<sup>6</sup> AAD	14W ATLS	$pp$ , 7, 8 TeV, $ZZ^* \rightarrow 4\ell$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.026	95	<sup>7</sup> KHACHATRYAN...16BA	CMS	$pp$ , 7, 8 TeV, $WW^{(*)}$
<0.0227	95	<sup>8</sup> AAD	15BE ATLS	$pp$ , 8 TeV, $ZZ^{(*)}, WW^{(*)}$
<0.046	95	<sup>9</sup> KHACHATRYAN...15BA	CMS	$pp$ , 7, 8 TeV, $ZZ^{(*)} \rightarrow 4\ell$
<3.4	95	<sup>10</sup> CHATRCHYAN	14AA CMS	$pp$ , 7, 8 TeV, $ZZ^* \rightarrow 4\ell$
<0.022	95	<sup>11</sup> KHACHATRYAN...14D	CMS	$pp$ , 7, 8 TeV, $ZZ^{(*)}$
<2.4	95	<sup>12</sup> KHACHATRYAN...14P	CMS	$pp$ , 7, 8 TeV, $\gamma\gamma$

<sup>1</sup> AABOUD 18BP use  $36.1 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . An observed upper limit on the off-shell Higgs signal strength of 3.8 is obtained at 95% CL using off-shell Higgs boson production in the  $ZZ \rightarrow 4\ell$  and  $ZZ \rightarrow 2\ell 2\nu$  decay channels ( $\ell = e, \mu$ ). Combining with the on-shell signal strength measurements, the quoted upper limit on the Higgs boson total width is obtained, assuming the ratios of the relevant Higgs-boson couplings to the SM predictions are constant with energy from on-shell production to the high-mass range.

<sup>2</sup> SIRUNYAN 17AV obtain an upper limit on the width from the  $m_{4\ell}$  distribution in  $ZZ^* \rightarrow 4\ell$  ( $\ell = e, \mu$ ) decays. Data of  $35.9 \text{ fb}^{-1}$   $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$  is used. The expected limit is 1.60 GeV.

<sup>3</sup> KHACHATRYAN 16BA combine the  $WW^{(*)}$  result with  $ZZ^{(*)}$  results of KHACHATRYAN 15BA and KHACHATRYAN 14D.

<sup>4</sup> KHACHATRYAN 15AM combine  $\gamma\gamma$  and  $ZZ^* \rightarrow 4\ell$  results. The expected limit is 2.3 GeV.

- <sup>5</sup> KHACHATRYAN 15BA derive a lower limit on the total width from an upper limit on the decay flight distance  $\tau < 1.9 \times 10^{-13}$  s.  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $19.7 \text{ fb}^{-1}$  at 8 TeV are used.
- <sup>6</sup> AAD 14W use  $4.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $20.3 \text{ fb}^{-1}$  at 8 TeV. The expected limit is 6.2 GeV.
- <sup>7</sup> KHACHATRYAN 16BA derive constraints on the total width from comparing  $WW^{(*)}$  production via on-shell and off-shell  $H^0$  using  $4.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $19.4 \text{ fb}^{-1}$  at 8 TeV.
- <sup>8</sup> AAD 15BE derive constraints on the total width from comparing  $ZZ^{(*)}$  and  $WW^{(*)}$  production via on-shell and off-shell  $H^0$  using  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. The K factor for the background processes is assumed to be equal to that for the signal.
- <sup>9</sup> KHACHATRYAN 15BA derive constraints on the total width from comparing  $ZZ^{(*)}$  production via on-shell and off-shell  $H^0$  with an unconstrained anomalous coupling.  $4\ell$  final states in  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV are used.
- <sup>10</sup> CHATRCHYAN 14AA use  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The expected limit is 2.8 GeV.
- <sup>11</sup> KHACHATRYAN 14D derive constraints on the total width from comparing  $ZZ^{(*)}$  production via on-shell and off-shell  $H^0$ .  $4\ell$  and  $\ell\ell\nu\nu$  final states in  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV are used.
- <sup>12</sup> KHACHATRYAN 14P use  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The expected limit is 3.1 GeV.

## $H^0$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $WW^*$		
$\Gamma_2$ $ZZ^*$		
$\Gamma_3$ $\gamma\gamma$		
$\Gamma_4$ $b\bar{b}$		
$\Gamma_5$ $e^+e^-$	$< 1.9 \times 10^{-3}$	95%
$\Gamma_6$ $\mu^+\mu^-$		
$\Gamma_7$ $\tau^+\tau^-$		
$\Gamma_8$ $Z\gamma$		
$\Gamma_9$ $\gamma^*\gamma$		
$\Gamma_{10}$ $J/\psi\gamma$	$< 3.5 \times 10^{-4}$	95%
$\Gamma_{11}$ $\psi(2S)\gamma$	$< 2.0 \times 10^{-3}$	95%
$\Gamma_{12}$ $\Upsilon(1S)\gamma$	$< 4.9 \times 10^{-4}$	95%
$\Gamma_{13}$ $\Upsilon(2S)\gamma$	$< 5.9 \times 10^{-4}$	95%
$\Gamma_{14}$ $\Upsilon(3S)\gamma$	$< 5.7 \times 10^{-4}$	95%
$\Gamma_{15}$ $\rho(770)\gamma$	$< 8.8 \times 10^{-4}$	95%
$\Gamma_{16}$ $\phi(1020)\gamma$	$< 4.8 \times 10^{-4}$	95%
$\Gamma_{17}$ $e\mu$	$< 3.5 \times 10^{-4}$	95%
$\Gamma_{18}$ $e\tau$	$< 6.1 \times 10^{-3}$	95%
$\Gamma_{19}$ $\mu\tau$	$< 2.5 \times 10^{-3}$	95%
$\Gamma_{20}$ invisible	$< 24 \%$	95%

**$H^0$  BRANCHING RATIOS** **$\Gamma(e^+e^-)/\Gamma_{\text{total}}$**  **$\Gamma_5/\Gamma$** 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<b><math>&lt;1.9 \times 10^{-3}</math></b>	95	<sup>1</sup> KHACHATRY...15H	CMS

<sup>1</sup> KHACHATRYAN 15H use  $5.0 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at  $8 \text{ TeV}$ .

 **$\Gamma(J/\psi\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_{10}/\Gamma$** 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;3.5 \times 10^{-4}</math></b>	95	<sup>1</sup> AABOUD	18BL ATLS	13 TeV, $36.1 \text{ fb}^{-1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.5 \times 10^{-3}$	95	<sup>2</sup> KHACHATRY...16B	CMS	8 TeV
-----------------------	----	------------------------------	-----	-------

$<1.5 \times 10^{-3}$	95	<sup>3</sup> AAD	15i ATLS	8 TeV
-----------------------	----	------------------	----------	-------

<sup>1</sup> AABOUD 18BL search for  $H^0 \rightarrow J/\psi\gamma$ ,  $J/\psi \rightarrow \mu^+\mu^-$  with  $36.1 \text{ fb}^{-1}$  of  $pp$  collision data at  $E_{\text{cm}} = 13 \text{ TeV}$ .

<sup>2</sup> KHACHATRYAN 16B use  $19.7 \text{ fb}^{-1}$  of  $pp$  collision data at  $8 \text{ TeV}$ .

<sup>3</sup> AAD 15i use  $19.7 \text{ fb}^{-1}$  of  $pp$  collision data at  $8 \text{ TeV}$ .

 **$\Gamma(\psi(2S)\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_{11}/\Gamma$** 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;2.0 \times 10^{-3}</math></b>	95	<sup>1</sup> AABOUD	18BL ATLS	13 TeV, $36.1 \text{ fb}^{-1}$

<sup>1</sup> AABOUD 18BL search for  $H^0 \rightarrow \psi(2S)\gamma$ ,  $\psi(2S) \rightarrow \mu^+\mu^-$  with  $36.1 \text{ fb}^{-1}$  of  $pp$  collision data at  $E_{\text{cm}} = 13 \text{ TeV}$ .

 **$\Gamma(\Upsilon(1S)\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_{12}/\Gamma$** 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;4.9 \times 10^{-4}</math></b>	95	<sup>1</sup> AABOUD	18BL ATLS	13 TeV, $36.1 \text{ fb}^{-1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.3 \times 10^{-3}$	95	<sup>2</sup> AAD	15i ATLS	8 TeV
-----------------------	----	------------------	----------	-------

<sup>1</sup> AABOUD 18BL search for  $H^0 \rightarrow \Upsilon(1S)\gamma$ ,  $\Upsilon(1S) \rightarrow \mu^+\mu^-$  with  $36.1 \text{ fb}^{-1}$  of  $pp$  collision data at  $E_{\text{cm}} = 13 \text{ TeV}$ .

<sup>2</sup> AAD 15i use  $19.7 \text{ fb}^{-1}$  of  $pp$  collision data at  $8 \text{ TeV}$ .

 **$\Gamma(\Upsilon(2S)\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_{13}/\Gamma$** 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;5.9 \times 10^{-4}</math></b>	95	<sup>1</sup> AABOUD	18BL ATLS	13 TeV, $36.1 \text{ fb}^{-1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.9 \times 10^{-3}$	95	<sup>2</sup> AAD	15i ATLS	8 TeV
-----------------------	----	------------------	----------	-------

<sup>1</sup> AABOUD 18BL search for  $H^0 \rightarrow \Upsilon(2S)\gamma$ ,  $\Upsilon(2S) \rightarrow \mu^+\mu^-$  with  $36.1 \text{ fb}^{-1}$  of  $pp$  collision data at  $E_{\text{cm}} = 13 \text{ TeV}$ .

<sup>2</sup> AAD 15i use  $19.7 \text{ fb}^{-1}$  of  $pp$  collision data at  $8 \text{ TeV}$ .

**$\Gamma(\Upsilon(3S)\gamma)/\Gamma_{\text{total}}$**   **$\Gamma_{14}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**$<5.7 \times 10^{-4}$**       95      <sup>1</sup> AABOUD      18BL ATLS      13 TeV, 36.1 fb<sup>-1</sup>

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.3 \times 10^{-3}$       95      <sup>2</sup> AAD      15l ATLS      8 TeV

<sup>1</sup> AABOUD 18BL search for  $H^0 \rightarrow \Upsilon(3S)\gamma$ ,  $\Upsilon(3S) \rightarrow \mu^+\mu^-$  with 36.1 fb<sup>-1</sup> of  $pp$  collision data at  $E_{\text{cm}} = 13$  TeV.

<sup>2</sup> AAD 15l use 19.7 fb<sup>-1</sup> of  $pp$  collision data at 8 TeV.

**$\Gamma(\rho(770)\gamma)/\Gamma_{\text{total}}$**   **$\Gamma_{15}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**$<8.8 \times 10^{-4}$**       95      <sup>1</sup> AABOUD      18AU ATLS       $pp$ , 13 TeV

<sup>1</sup> AABOUD 18AU use 35.6 fb<sup>-1</sup> of  $pp$  collision data at 13 TeV.

**$\Gamma(\phi(1020)\gamma)/\Gamma_{\text{total}}$**   **$\Gamma_{16}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**$<4.8 \times 10^{-4}$**       95      <sup>1</sup> AABOUD      18AU ATLS       $pp$ , 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.4 \times 10^{-3}$       95      <sup>2</sup> AABOUD      16K ATLS       $pp$ , 13 TeV

<sup>1</sup> AABOUD 18AU use 35.6 fb<sup>-1</sup> of  $pp$  collision data at 13 TeV.

<sup>2</sup> AABOUD 16K use 2.7 fb<sup>-1</sup> of  $pp$  collision data at 13 TeV.

**$\Gamma(e\mu)/\Gamma_{\text{total}}$**   **$\Gamma_{17}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**$<3.5 \times 10^{-4}$**       95      <sup>1</sup> KHACHATRY...16CD CMS       $pp$ , 8 TeV

<sup>1</sup> KHACHATRYAN 16CD search for  $H^0 \rightarrow e\mu$  in 19.7 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. The limit constrains the  $Y_{e\mu}$  Yukawa coupling to  $\sqrt{|Y_{e\mu}|^2 + |Y_{\mu e}|^2} < 5.4 \times 10^{-4}$  at 95% CL (see their Fig. 6).

**$\Gamma(e\tau)/\Gamma_{\text{total}}$**   **$\Gamma_{18}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**$<6.1 \times 10^{-3}$**       95      <sup>1</sup> SIRUNYAN      18BH CMS       $pp$ , 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.04 \times 10^{-2}$       95      <sup>2</sup> AAD      17 ATLS       $pp$ , 8 TeV

$<6.9 \times 10^{-3}$       95      <sup>3</sup> KHACHATRY...16CD CMS       $pp$ , 8 TeV

<sup>1</sup> SIRUNYAN 18BH search for  $H^0 \rightarrow e\tau$  in 35.9 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV.

The limit constrains the  $Y_{e\tau}$  Yukawa coupling to  $\sqrt{|Y_{e\tau}|^2 + |Y_{\tau e}|^2} < 2.26 \times 10^{-3}$  at 95% CL (see their Fig. 10).

<sup>2</sup> AAD 17 search for  $H^0 \rightarrow e\tau$  in 20.3 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV.

<sup>3</sup> KHACHATRYAN 16CD search for  $H^0 \rightarrow e\tau$  in 19.7 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. The limit constrains the  $Y_{e\tau}$  Yukawa coupling to  $\sqrt{|Y_{e\tau}|^2 + |Y_{\tau e}|^2} < 2.4 \times 10^{-3}$  at 95% CL (see their Fig. 6).



$\Gamma(\mu\tau)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.5 × 10<sup>-3</sup></b>	95	<sup>1</sup> SIRUNYAN	18BH CMS	$pp$ , 13 TeV
<0.26	95	<sup>2</sup> AAIJ	18AMLHCB	$pp$ , 8 TeV
<1.43 × 10 <sup>-2</sup>	95	<sup>3</sup> AAD	17 ATLS	$pp$ , 8 TeV
<1.51 × 10 <sup>-2</sup>	95	<sup>4</sup> KHACHATRYAN...15Q	CMS	$pp$ , 8 TeV

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup> SIRUNYAN 18BH search for  $H^0 \rightarrow \mu\tau$  in 35.9 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. The limit constrains the  $Y_{\mu\tau}$  Yukawa coupling to  $\sqrt{|Y_{\mu\tau}|^2 + |Y_{\tau\mu}|^2} < 1.43 \times 10^{-3}$  at 95% CL (see their Fig. 10).
- <sup>2</sup> AAIJ 18AM search for  $H^0 \rightarrow \mu\tau$  in 2.0 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. The limit constrains the  $Y_{\mu\tau}$  Yukawa coupling to  $\sqrt{|Y_{\mu\tau}|^2 + |Y_{\tau\mu}|^2} < 1.7 \times 10^{-2}$  at 95% CL assuming SM production cross sections.
- <sup>3</sup> AAD 17 search for  $H^0 \rightarrow \mu\tau$  in 20.3 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV.
- <sup>4</sup> KHACHATRYAN 15Q search for  $H^0 \rightarrow \mu\tau$  with  $\tau$  decaying electronically or hadronically in 19.7 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. The fit gives  $B(H^0 \rightarrow \mu\tau) = (0.84^{+0.39}_{-0.37})\%$  with a significance of 2.4  $\sigma$ .

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$   
 Invisible final states.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.24</b>	95	<sup>1</sup> KHACHATRYAN...17F	CMS	$pp$ , 7, 8, 13 TeV
<0.67	95	<sup>2</sup> AABOUD	18 ATLS	$pp \rightarrow H^0 ZX$ , $H^0 \rightarrow$ inv., 13 TeV
<0.83	95	<sup>3</sup> AABOUD	18CA ATLS	$pp \rightarrow H^0 W/Z$ , $W/Z \rightarrow jj$ , 13 TeV
<0.46	95	<sup>4</sup> AABOUD	17BD ATLS	$pp \rightarrow gH^0 X$ , $qqH^0 X$ , $H^0 \rightarrow$ inv, 13 TeV
<0.28	95	<sup>5</sup> AAD	16AF ATLS	$pp \rightarrow qqH^0 X$ , 8 TeV
<0.34	95	<sup>6</sup> AAD	16AN LHC	$pp$ , 7, 8 TeV
<0.78	95	<sup>7</sup> AAD	15BD ATLS	$pp \rightarrow H^0 W/Z X$ , 8 TeV
<0.75	95	<sup>8</sup> AAD	14O ATLS	$pp \rightarrow H^0 ZX$ , 7, 8 TeV
<0.58	95	<sup>9</sup> CHATRCHYAN 14B	CMS	$pp \rightarrow H^0 ZX$ , $qqH^0 X$
<0.81	95	<sup>10</sup> CHATRCHYAN 14B	CMS	$pp \rightarrow H^0 ZX$ , 7, 8 TeV
<0.65	95	<sup>11</sup> CHATRCHYAN 14B	CMS	$pp \rightarrow qqH^0 X$ , 8 TeV

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup> KHACHATRYAN 17F search for  $H^0$  decaying to invisible final states with gluon fusion, VBF,  $ZH$ , and  $WH$  productions using 2.3 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV, 19.7 fb<sup>-1</sup> at 8 TeV, and 5.1 fb<sup>-1</sup> at 7 TeV. The quoted limit is given for  $m_{H^0} = 125$  GeV and assumes the Standard Model rates for gluon fusion, VBF,  $ZH$ , and  $WH$  productions.
- <sup>2</sup> AABOUD 18 search for  $pp \rightarrow H^0 ZX$ ,  $Z \rightarrow ee, \mu\mu$  with  $H^0$  decaying to invisible final states in 36.1 fb<sup>-1</sup> at  $E_{\text{cm}} = 13$  TeV. The quoted limit on the branching ratio is given for  $m_{H^0} = 125$  GeV and assumes the Standard Model rate for  $H^0 Z$  production.
- <sup>3</sup> AABOUD 18CA search for  $H^0$  decaying to invisible final states using  $WH$ , and  $ZH$  productions, where  $W$  and  $Z$  hadronically decay. The data of 36.1 fb<sup>-1</sup> at  $E_{\text{cm}} = 13$  TeV is used. The quoted limit assumes SM production cross sections with combining the contributions from  $WH$ ,  $ZH$ , ggF and VBF production modes.

- <sup>4</sup> AABOUD 17BD search for  $H^0$  decaying to invisible final states with  $\geq 1$  jet and VBF events using  $3.2 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . A cross-section ratio  $R^{\text{miss}}$  is used in the measurement. The quoted limit is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>5</sup> AAD 16AF search for  $pp \rightarrow qqH^0X$  (VBF) with  $H^0$  decaying to invisible final states in  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted limit on the branching ratio is given for  $m_{H^0} = 125 \text{ GeV}$  and assumes the Standard Model rates for VBF and gluon-fusion production.
- <sup>6</sup> AAD 16AN perform fits to the ATLAS and CMS data at  $E_{\text{cm}} = 7$  and  $8 \text{ TeV}$ . The branching fraction of decays into BSM particles that are invisible or into undetected decay modes is measured for  $m_{H^0} = 125.09 \text{ GeV}$ .
- <sup>7</sup> AAD 15BD search for  $pp \rightarrow H^0WX$  and  $pp \rightarrow H^0ZX$  with  $W$  or  $Z$  decaying hadronically and  $H^0$  decaying to invisible final states using data at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted limit is given for  $m_{H^0} = 125 \text{ GeV}$ , assumes the Standard Model rates for the production processes and is based on a combination of the contributions from  $H^0W$ ,  $H^0Z$  and the gluon-fusion process.
- <sup>8</sup> AAD 14O search for  $pp \rightarrow H^0ZX$ ,  $Z \rightarrow \ell\ell$ , with  $H^0$  decaying to invisible final states in  $4.5 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted limit on the branching ratio is given for  $m_{H^0} = 125.5 \text{ GeV}$  and assumes the Standard Model rate for  $H^0Z$  production.
- <sup>9</sup> CHATRCHYAN 14B search for  $pp \rightarrow H^0ZX$ ,  $Z \rightarrow \ell\ell$  and  $Z \rightarrow b\bar{b}$ , and also  $pp \rightarrow qqH^0X$  with  $H^0$  decaying to invisible final states using data at  $E_{\text{cm}} = 7$  and  $8 \text{ TeV}$ . The quoted limit on the branching ratio is obtained from a combination of the limits from  $H^0Z$  and  $qqH^0$ . It is given for  $m_{H^0} = 125 \text{ GeV}$  and assumes the Standard Model rates for the two production processes.
- <sup>10</sup> CHATRCHYAN 14B search for  $pp \rightarrow H^0ZX$  with  $H^0$  decaying to invisible final states and  $Z \rightarrow \ell\ell$  in  $4.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ , and also with  $Z \rightarrow b\bar{b}$  in  $18.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted limit on the branching ratio is given for  $m_{H^0} = 125 \text{ GeV}$  and assumes the Standard Model rate for  $H^0Z$  production.
- <sup>11</sup> CHATRCHYAN 14B search for  $pp \rightarrow qqH^0X$  (vector boson fusion) with  $H^0$  decaying to invisible final states in  $19.5 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted limit on the branching ratio is given for  $m_{H^0} = 125 \text{ GeV}$  and assumes the Standard Model rate for  $qqH^0$  production.

## $H^0$ SIGNAL STRENGTHS IN DIFFERENT CHANNELS

The  $H^0$  signal strength in a particular final state  $xx$  is given by the cross section times branching ratio in this channel normalized to the Standard Model (SM) value,  $\sigma \cdot B(H^0 \rightarrow xx) / (\sigma \cdot B(H^0 \rightarrow xx))_{\text{SM}}$ , for the specified mass value of  $H^0$ . For the SM predictions, see DITTMAIER 11, DITTMAIER 12, and HEINEMEYER 13A. Results for fiducial and differential cross sections are also listed below.

### Combined Final States

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.10 ± 0.11 OUR AVERAGE</b>			
$1.09 \pm 0.07 \pm 0.04 \pm 0.03^{+0.07}_{-0.06}$	<sup>1,2</sup> AAD	16AN LHC	$pp$ , 7, 8 TeV
$1.44^{+0.59}_{-0.56}$	<sup>3</sup> AALTONEN	13M TEVA	$p\bar{p} \rightarrow H^0X$ , 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.20 \pm 0.10 \pm 0.06 \pm 0.04^{+0.08}_{-0.07}$	2 AAD	16AN ATLS	$pp$ , 7, 8 TeV
$0.97 \pm 0.09 \pm 0.05^{+0.04+0.07}_{-0.03-0.06}$	2 AAD	16AN CMS	$pp$ , 7, 8 TeV
$1.18 \pm 0.10 \pm 0.07^{+0.08}_{-0.07}$	4 AAD	16K ATLS	$pp$ , 7, 8 TeV
$0.75^{+0.28+0.13+0.08}_{-0.26-0.11-0.05}$	4 AAD	16K ATLS	$pp$ , 7 TeV
$1.28 \pm 0.11^{+0.08+0.10}_{-0.07-0.08}$	4 AAD	16K ATLS	$pp$ , 8 TeV
	5 AAD	15P ATLS	$pp$ , 8 TeV, cross section
$1.00 \pm 0.09 \pm 0.07^{+0.08}_{-0.07}$	6 KHACHATRYAN	15AM CMS	$pp$ , 7, 8 TeV
$1.33^{+0.14}_{-0.10} \pm 0.15$	7 AAD	13AK ATLS	$pp$ , 7 and 8 TeV
$1.54^{+0.77}_{-0.73}$	8 AALTONEN	13L CDF	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
$1.40^{+0.92}_{-0.88}$	9 ABAZOV	13L D0	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
$1.4 \pm 0.3$	10 AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 7, 8 TeV
$1.2 \pm 0.4$	10 AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 7 TeV
$1.5 \pm 0.4$	10 AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 8 TeV
$0.87 \pm 0.23$	11 CHATRCHYAN	12N CMS	$pp \rightarrow H^0 X$ , 7, 8 TeV

<sup>1</sup> AAD 16AN perform fits to the ATLAS and CMS data at  $E_{\text{cm}} = 7$  and 8 TeV. The signal strengths for individual production processes are  $1.03^{+0.16}_{-0.14}$  for gluon fusion,  $1.18^{+0.25}_{-0.23}$  for vector boson fusion,  $0.89^{+0.40}_{-0.38}$  for  $WH^0$  production,  $0.79^{+0.38}_{-0.36}$  for  $ZH^0$  production, and  $2.3^{+0.7}_{-0.6}$  for  $t\bar{t}H^0$  production.

<sup>2</sup> AAD 16AN: The uncertainties represent statistics, experimental systematics, theory systematics on the background, and theory systematics on the signal. The quoted signal strengths are given for  $m_{H^0} = 125.09$  GeV. In the fit, relative branching ratios and relative production cross sections are fixed to those in the Standard Model.

<sup>3</sup> AALTONEN 13M combine all Tevatron data from the CDF and D0 Collaborations with up to  $10.0 \text{ fb}^{-1}$  and  $9.7 \text{ fb}^{-1}$ , respectively, of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.

<sup>4</sup> AAD 16K use up to  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and up to  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The third uncertainty in the measurement is theory systematics. The signal strengths for individual production modes are  $1.23 \pm 0.14^{+0.09+0.16}_{-0.08-0.12}$  for gluon fusion,  $1.23^{+0.28+0.13+0.11}_{-0.27-0.12-0.09}$  for vector boson fusion,  $0.80^{+0.31}_{-0.30} \pm 0.17^{+0.10}_{-0.05}$  for  $W/ZH^0$  production, and  $1.81^{+0.52+0.58+0.31}_{-0.50-0.55-0.12}$  for  $t\bar{t}H^0$  production. The quoted signal strengths are given for  $m_{H^0} = 125.36$  GeV.

<sup>5</sup> AAD 15P measure total and differential cross sections of the process  $pp \rightarrow H^0 X$  at  $E_{\text{cm}} = 8$  TeV with  $20.3 \text{ fb}^{-1}$ .  $\gamma\gamma$  and  $4\ell$  final states are used.  $\sigma(pp \rightarrow H^0 X) = 33.0 \pm 5.3 \pm 1.6 \text{ pb}$  is given. See their Figs. 2 and 3 for data on differential cross sections.

<sup>6</sup> KHACHATRYAN 15AM use up to  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and up to  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The third uncertainty in the measurement is theory systematics. Fits to each production mode give the value of  $0.85^{+0.19}_{-0.16}$  for gluon fusion,  $1.16^{+0.37}_{-0.34}$  for vector boson fusion,  $0.92^{+0.38}_{-0.36}$  for  $WH^0$ ,  $ZH^0$  production, and  $2.90^{+1.08}_{-0.94}$  for  $t\bar{t}H^0$  production.

- <sup>7</sup> AAD 13AK use  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $20.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The combined signal strength is based on the  $\gamma\gamma$ ,  $ZZ^* \rightarrow 4\ell$ , and  $WW^* \rightarrow \ell\nu\ell\nu$  channels. The quoted signal strength is given for  $m_{H^0} = 125.5 \text{ GeV}$ . Reported statistical error value modified following private communication with the experiment.
- <sup>8</sup> AALTONEN 13L combine all CDF results with  $9.45\text{--}10.0 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>9</sup> ABAZOV 13L combine all D0 results with up to  $9.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>10</sup> AAD 12AI obtain results based on  $4.6\text{--}4.8 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $5.8\text{--}5.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . An excess of events over background with a local significance of  $5.9 \sigma$  is observed at  $m_{H^0} = 126 \text{ GeV}$ . The quoted signal strengths are given for  $m_{H^0} = 126 \text{ GeV}$ . See also AAD 12DA.
- <sup>11</sup> CHATRCHYAN 12N obtain results based on  $4.9\text{--}5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $5.1\text{--}5.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . An excess of events over background with a local significance of  $5.0 \sigma$  is observed at about  $m_{H^0} = 125 \text{ GeV}$ . The combined signal strength is based on the  $\gamma\gamma$ ,  $ZZ^*$ ,  $WW^*$ ,  $\tau^+\tau^-$ , and  $b\bar{b}$  channels. The quoted signal strength is given for  $m_{H^0} = 125.5 \text{ GeV}$ . See also CHATRCHYAN 13Y.

## WW\* Final State

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>1.08^{+0.18}_{-0.16}</math> OUR AVERAGE</b>			
$1.09^{+0.18}_{-0.16}$	1,2 AAD	16AN LHC	$pp$ , 7, 8 TeV
$0.94^{+0.85}_{-0.83}$	<sup>3</sup> AALTONEN	13M TEVA	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
	<sup>4</sup> AABOUD	19F ATLS	$pp$ , 13 TeV, cross sections
$1.22^{+0.23}_{-0.21}$	<sup>2</sup> AAD	16AN ATLS	$pp$ , 7, 8 TeV
$0.90^{+0.23}_{-0.21}$	<sup>2</sup> AAD	16AN CMS	$pp$ , 7, 8 TeV
	<sup>5</sup> AAD	16AO ATLS	$pp$ , 8 TeV, cross sections
$1.18 \pm 0.16^{+0.17}_{-0.14}$	<sup>6</sup> AAD	16K ATLS	$pp$ , 7, 8 TeV
$1.09^{+0.16+0.17}_{-0.15-0.14}$	<sup>7</sup> AAD	15AA ATLS	$pp$ , 7, 8 TeV
$3.0^{+1.3+1.0}_{-1.1-0.7}$	<sup>8</sup> AAD	15AQ ATLS	$pp \rightarrow H^0 W/Z X$ , 7, 8 TeV
$1.16^{+0.16+0.18}_{-0.15-0.15}$	<sup>9</sup> AAD	15AQ ATLS	$pp$ , 7, 8 TeV
$0.72 \pm 0.12 \pm 0.10^{+0.12}_{-0.10}$	<sup>10</sup> CHATRCHYAN 14G	CMS	$pp$ , 7, 8 TeV
$0.99^{+0.31}_{-0.28}$	<sup>11</sup> AAD	13AK ATLS	$pp$ , 7 and 8 TeV
$0.00^{+1.78}_{-0.00}$	<sup>12</sup> AALTONEN	13L CDF	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
$1.90^{+1.63}_{-1.52}$	<sup>13</sup> ABAZOV	13L D0	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV

1.3 ± 0.5	14 AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 7, 8 TeV
0.5 ± 0.6	14 AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 7 TeV
1.9 ± 0.7	14 AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 8 TeV
0.60 <sup>+0.42</sup> <sub>-0.37</sub>	15 CHATRCHYAN	12N CMS	$pp \rightarrow H^0 X$ , 7, 8 TeV

- <sup>1</sup> AAD 16AN perform fits to the ATLAS and CMS data at  $E_{\text{cm}} = 7$  and 8 TeV. The signal strengths for individual production processes are  $0.84 \pm 0.17$  for gluon fusion,  $1.2 \pm 0.4$  for vector boson fusion,  $1.6^{+1.2}_{-1.0}$  for  $WH^0$  production,  $5.9^{+2.6}_{-2.2}$  for  $ZH^0$  production, and  $5.0^{+1.8}_{-1.7}$  for  $t\bar{t}H^0$  production.
- <sup>2</sup> AAD 16AN: In the fit, relative production cross sections are fixed to those in the Standard Model. The quoted signal strength is given for  $m_{H^0} = 125.09$  GeV.
- <sup>3</sup> AALTONEN 13M combine all Tevatron data from the CDF and D0 Collaborations with up to  $10.0 \text{ fb}^{-1}$  and  $9.7 \text{ fb}^{-1}$ , respectively, of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.
- <sup>4</sup> AABOUD 19F measure cross-sections times the  $H^0 \rightarrow WW^*$  branching fraction in the  $H^0 \rightarrow WW^* \rightarrow e\nu\mu\nu$  channel using  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV:  $\sigma_{ggF} \times B(H^0 \rightarrow WW^*) = 11.4^{+1.2+1.8}_{-1.1-1.7}$  pb and  $\sigma_{VBF} \times B(H^0 \rightarrow WW^*) = 0.50^{+0.24}_{-0.22} \pm 0.17$  pb.
- <sup>5</sup> AAD 16AO measure fiducial total and differential cross sections of gluon fusion process at  $E_{\text{cm}} = 8$  TeV with  $20.3 \text{ fb}^{-1}$  using  $H^0 \rightarrow WW^* \rightarrow e\nu\mu\nu$ . The measured fiducial total cross section is  $36.0 \pm 9.7 \text{ fb}$  in their fiducial region (Table 7). See their Fig. 6 for fiducial differential cross sections. The results are given for  $m_{H^0} = 125$  GeV.
- <sup>6</sup> AAD 16K use up to  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and up to  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125.36$  GeV.
- <sup>7</sup> AAD 15AA use  $4.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The signal strength for the gluon fusion and vector boson fusion mode is  $1.02 \pm 0.19^{+0.22}_{-0.18}$  and  $1.27^{+0.44+0.30}_{-0.40-0.21}$ , respectively. The quoted signal strengths are given for  $m_{H^0} = 125.36$  GeV.
- <sup>8</sup> AAD 15AQ use  $4.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125.36$  GeV.
- <sup>9</sup> AAD 15AQ combine their result on  $W/ZH^0$  production with the results of AAD 15AA (gluon fusion and vector boson fusion, slightly updated). The quoted signal strength is given for  $m_{H^0} = 125.36$  GeV.
- <sup>10</sup> CHATRCHYAN 14G use  $4.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $19.4 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The last uncertainty in the measurement is theory systematics. The quoted signal strength is given for  $m_{H^0} = 125.6$  GeV.
- <sup>11</sup> AAD 13AK use  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $20.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125.5$  GeV. Superseded by AAD 15AA.
- <sup>12</sup> AALTONEN 13L combine all CDF results with  $9.45\text{--}10.0 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.
- <sup>13</sup> ABAZOV 13L combine all D0 results with up to  $9.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.
- <sup>14</sup> AAD 12AI obtain results based on  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $5.8 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The quoted signal strengths are given for  $m_{H^0} = 126$  GeV. See also AAD 12DA.
- <sup>15</sup> CHATRCHYAN 12N obtain results based on  $4.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $5.1 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125.5$  GeV. See also CHATRCHYAN 13Y.

**ZZ\* Final State**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.19<sup>+0.12</sup><sub>-0.11</sub> OUR AVERAGE</b>			
1.28 <sup>+0.21</sup> <sub>-0.19</sub>	1 AABOUD	18AJ ATLS	$pp$ , 13 TeV
1.05 <sup>+0.15+0.11</sup> <sub>-0.14-0.09</sub>	2 SIRUNYAN	17AV CMS	$pp$ , 13 TeV
1.29 <sup>+0.26</sup> <sub>-0.23</sub>	3,4 AAD	16AN LHC	$pp$ , 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.52 <sup>+0.40</sup> <sub>-0.34</sub>	4 AAD	16AN ATLS	$pp$ , 7, 8 TeV
1.04 <sup>+0.32</sup> <sub>-0.26</sub>	4 AAD	16AN CMS	$pp$ , 7, 8 TeV
1.46 <sup>+0.35+0.19</sup> <sub>-0.31-0.13</sub>	5 AAD	16K ATLS	$pp$ , 7, 8 TeV
	6 KHACHATRYAN	16AR CMS	$pp$ , 7, 8 TeV cross sections
1.44 <sup>+0.34+0.21</sup> <sub>-0.31-0.11</sub>	7 AAD	15F ATLS	$pp \rightarrow H^0 X$ , 7, 8 TeV
	8 AAD	14AR ATLS	$pp$ , 8 TeV, differential cross section
0.93 <sup>+0.26+0.13</sup> <sub>-0.23-0.09</sub>	9 CHATRCHYAN	14AA CMS	$pp$ , 7, 8 TeV
1.43 <sup>+0.40</sup> <sub>-0.35</sub>	10 AAD	13AK ATLS	$pp$ , 7 and 8 TeV
0.80 <sup>+0.35</sup> <sub>-0.28</sub>	11 CHATRCHYAN	13J CMS	$pp \rightarrow H^0 X$ , 7, 8 TeV
1.2 $\pm$ 0.6	12 AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 7, 8 TeV
1.4 $\pm$ 1.1	12 AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 7 TeV
1.1 $\pm$ 0.8	12 AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 8 TeV
0.73 <sup>+0.45</sup> <sub>-0.33</sub>	13 CHATRCHYAN	12N CMS	$pp \rightarrow H^0 X$ , 7, 8 TeV

<sup>1</sup> AABOUD 18AJ perform analyses using  $H^0 \rightarrow ZZ^* \rightarrow 4\ell$  ( $\ell = e, \mu$ ) with data of  $36.1 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . Results are given for  $m_{H^0} = 125.09 \text{ GeV}$ . The inclusive cross section times branching ratio for  $H^0 \rightarrow ZZ^*$  decay ( $|\eta(H^0)| < 2.5$ ) is measured to be  $1.73^{+0.26}_{-0.24} \text{ pb}$  (with  $1.34^{+0.09}_{-0.09} \text{ pb}$  expected in the SM).

<sup>2</sup> SIRUNYAN 17AV use  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . The quoted signal strength, obtained from the analysis of  $H^0 \rightarrow ZZ^* \rightarrow 4\ell$  ( $\ell = e, \mu$ ) decays, is given for  $m_{H^0} = 125.09 \text{ GeV}$ . The signal strengths for different production modes are given in their Table 3. The fiducial and differential cross sections are shown in their Fig. 10.

<sup>3</sup> AAD 16AN perform fits to the ATLAS and CMS data at  $E_{\text{cm}} = 7$  and 8 TeV. The signal strengths for individual production processes are  $1.13^{+0.34}_{-0.31}$  for gluon fusion and  $0.1^{+1.1}_{-0.6}$  for vector boson fusion.

<sup>4</sup> AAD 16AN: In the fit, relative production cross sections are fixed to those in the Standard Model. The quoted signal strength is given for  $m_{H^0} = 125.09 \text{ GeV}$ .

<sup>5</sup> AAD 16K use up to  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and up to  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125.36 \text{ GeV}$ .

<sup>6</sup> KHACHATRYAN 16AR use data of  $5.1 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at 8 TeV. The fiducial cross sections for the production of 4 leptons via  $H^0 \rightarrow 4\ell$  decays are measured to be  $0.56^{+0.67+0.21}_{-0.44-0.06} \text{ fb}$  at 7 TeV and  $1.11^{+0.41+0.14}_{-0.35-0.10} \text{ fb}$  at 8 TeV in

their fiducial region (Table 2). The differential cross sections at  $E_{\text{cm}} = 8$  TeV are also shown in Figs. 4 and 5. The results are given for  $m_{H^0} = 125$  GeV.

- <sup>7</sup> AAD 15F use  $4.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125.36$  GeV. The signal strength for the gluon fusion production mode is  $1.66^{+0.45+0.25}_{-0.41-0.15}$ , while the signal strength for the vector boson fusion production mode is  $0.26^{+1.60+0.36}_{-0.91-0.23}$ .
- <sup>8</sup> AAD 14AR measure the cross section for  $pp \rightarrow H^0 X$ ,  $H^0 \rightarrow ZZ^*$  using  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. They give  $\sigma \cdot B = 2.11^{+0.53}_{-0.47} \pm 0.08$  fb in their fiducial region, where  $1.30 \pm 0.13$  fb is expected in the Standard Model for  $m_{H^0} = 125.4$  GeV. Various differential cross sections are also given, which are in agreement with the Standard Model expectations.
- <sup>9</sup> CHATRCHYAN 14AA use  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125.6$  GeV. The signal strength for the gluon fusion and  $t\bar{t}H$  production mode is  $0.80^{+0.46}_{-0.36}$ , while the signal strength for the vector boson fusion and  $WH^0$ ,  $ZH^0$  production mode is  $1.7^{+2.2}_{-2.1}$ .
- <sup>10</sup> AAD 13AK use  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $20.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125.5$  GeV.
- <sup>11</sup> CHATRCHYAN 13J obtain results based on  $ZZ \rightarrow 4\ell$  final states in  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $12.2 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125.8$  GeV. Superseded by CHATRCHYAN 14AA.
- <sup>12</sup> AAD 12AI obtain results based on  $4.7\text{--}4.8 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $5.8 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The quoted signal strengths are given for  $m_{H^0} = 126$  GeV. See also AAD 12DA.
- <sup>13</sup> CHATRCHYAN 12N obtain results based on  $4.9\text{--}5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $5.1\text{--}5.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. An excess of events over background with a local significance of  $5.0 \sigma$  is observed at about  $m_{H^0} = 125$  GeV. The quoted signal strengths are given for  $m_{H^0} = 125.5$  GeV. See also CHATRCHYAN 12BY and CHATRCHYAN 13Y.

## $\gamma\gamma$ Final State

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>1.10^{+0.10}_{-0.09}</math> OUR AVERAGE</b>			
$0.99^{+0.15}_{-0.14}$	1 AABOUD	18BO ATLS	$pp$ , 13 TeV, $36.1 \text{ fb}^{-1}$
$1.18^{+0.17}_{-0.14}$	2 SIRUNYAN	18DS CMS	$pp$ , $H^0 \rightarrow \gamma\gamma$ , 13 TeV, floated $m_{H^0}$
$1.14^{+0.19}_{-0.18}$	3,4 AAD	16AN LHC	$pp$ , 7, 8 TeV
$5.97^{+3.39}_{-3.12}$	5 AALTONEN	13M TEVA	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.14^{+0.27}_{-0.25}$	6 SIRUNYAN	19L CMS	$pp$ , 13 TeV, diff. x-section
$1.11^{+0.25}_{-0.23}$	4 AAD	16AN ATLS	$pp$ , 7, 8 TeV
	4 AAD	16AN CMS	$pp$ , 7, 8 TeV
	7 KHACHATRY...	16G CMS	$pp$ , 8 TeV, diff. x-section
$1.17 \pm 0.23^{+0.10+0.12}_{-0.08-0.08}$	8 AAD	14BC ATLS	$pp \rightarrow H^0 X$ , 7, 8 TeV
	9 AAD	14BJ ATLS	$pp$ , 8 TeV, diff. x-section

$1.14 \pm 0.21^{+0.09+0.13}_{-0.05-0.09}$	<sup>10</sup> KHACHATRYAN 14P	CMS	$pp$ , 7, 8 TeV
$1.55^{+0.33}_{-0.28}$	<sup>11</sup> AAD	13AK ATLS	$pp$ , 7 and 8 TeV
$7.81^{+4.61}_{-4.42}$	<sup>12</sup> AALTONEN	13L CDF	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
$4.20^{+4.60}_{-4.20}$	<sup>13</sup> ABAZOV	13L D0	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
$1.8 \pm 0.5$	<sup>14</sup> AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 7, 8 TeV
$2.2 \pm 0.7$	<sup>14</sup> AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 7 TeV
$1.5 \pm 0.6$	<sup>14</sup> AAD	12AI ATLS	$pp \rightarrow H^0 X$ , 8 TeV
$1.54^{+0.46}_{-0.42}$	<sup>15</sup> CHATRCHYAN 12N	CMS	$pp \rightarrow H^0 X$ , 7, 8 TeV

<sup>1</sup> AABOUD 18B0 use  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . The signal strengths for the individual production modes are:  $0.81^{+0.19}_{-0.18}$  for gluon fusion,  $2.0^{+0.6}_{-0.5}$  for vector boson fusion,  $0.7^{+0.9}_{-0.8}$  for  $VH^0$  production ( $V = W, Z$ ), and  $0.5 \pm 0.6$  for  $t\bar{t}H^0$  and  $tH^0$  production. Other measurements of cross sections and couplings are summarized in their Section 10. The quoted values are given for  $m_{H^0} = 125.09 \text{ GeV}$ .

<sup>2</sup> SIRUNYAN 18DS use  $35.9 \text{ fb}^{-1}$  of  $pp \rightarrow H^0$  collisions with  $H^0 \rightarrow \gamma\gamma$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The Higgs mass is floated in the measurement of a signal strength. The result is  $1.18^{+0.12}_{-0.11}(\text{stat.})^{+0.09}_{-0.07}(\text{syst.})^{+0.07}_{-0.06}(\text{theory})$ , which is largely insensitive to the Higgs mass around 125 GeV.

<sup>3</sup> AAD 16AN perform fits to the ATLAS and CMS data at  $E_{\text{cm}} = 7$  and 8 TeV. The signal strengths for individual production processes are  $1.10^{+0.23}_{-0.22}$  for gluon fusion,  $1.3 \pm 0.5$  for vector boson fusion,  $0.5^{+1.3}_{-1.2}$  for  $WH^0$  production,  $0.5^{+3.0}_{-2.5}$  for  $ZH^0$  production, and  $2.2^{+1.6}_{-1.3}$  for  $t\bar{t}H^0$  production.

<sup>4</sup> AAD 16AN: In the fit, relative production cross sections are fixed to those in the Standard Model. The quoted signal strength is given for  $m_{H^0} = 125.09 \text{ GeV}$ .

<sup>5</sup> AALTONEN 13M combine all Tevatron data from the CDF and D0 Collaborations with up to  $10.0 \text{ fb}^{-1}$  and  $9.7 \text{ fb}^{-1}$ , respectively, of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .

<sup>6</sup> SIRUNYAN 19L measure fiducial and differential cross sections of the process  $pp \rightarrow H^0 \rightarrow \gamma\gamma$  at  $E_{\text{cm}} = 13 \text{ TeV}$  with  $35.9 \text{ fb}^{-1}$ . See their Figs. 4–11.

<sup>7</sup> KHACHATRYAN 16G measure fiducial and differential cross sections of the process  $pp \rightarrow H^0 X$ ,  $H^0 \rightarrow \gamma\gamma$  at  $E_{\text{cm}} = 8 \text{ TeV}$  with  $19.7 \text{ fb}^{-1}$ . See their Figs. 4–6 and Table 1 for data.

<sup>8</sup> AAD 14BC use  $4.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The last uncertainty in the measurement is theory systematics. The quoted signal strength is given for  $m_{H^0} = 125.4 \text{ GeV}$ . The signal strengths for the individual production modes are:  $1.32 \pm 0.38$  for gluon fusion,  $0.8 \pm 0.7$  for vector boson fusion,  $1.0 \pm 1.6$  for  $WH^0$  production,  $0.1^{+3.7}_{-0.1}$  for  $ZH^0$  production, and  $1.6^{+2.7}_{-1.8}$  for  $t\bar{t}H^0$  production.

<sup>9</sup> AAD 14BJ measure fiducial and differential cross sections of the process  $pp \rightarrow H^0 X$ ,  $H^0 \rightarrow \gamma\gamma$  at  $E_{\text{cm}} = 8 \text{ TeV}$  with  $20.3 \text{ fb}^{-1}$ . See their Table 3 and Figs. 3–12 for data.

<sup>10</sup> KHACHATRYAN 14P use  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The last uncertainty in the measurement is theory systematics. The quoted signal strength is given for  $m_{H^0} = 124.7 \text{ GeV}$ . The signal strength for the gluon fusion and  $t\bar{t}H$  production mode is  $1.13^{+0.37}_{-0.31}$ , while the signal strength for the vector boson fusion and  $WH^0$ ,  $ZH^0$  production mode is  $1.16^{+0.63}_{-0.58}$ .



- <sup>11</sup> AAD 13AK use  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $20.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125.5 \text{ GeV}$ .
- <sup>12</sup> AALTONEN 13L combine all CDF results with  $9.45\text{--}10.0 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>13</sup> ABAZOV 13L combine all D0 results with up to  $9.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>14</sup> AAD 12AI obtain results based on  $4.8 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $5.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted signal strengths are given for  $m_{H^0} = 126 \text{ GeV}$ . See also AAD 12DA.
- <sup>15</sup> CHATRCHYAN 12N obtain results based on  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}}=7 \text{ TeV}$  and  $5.3 \text{ fb}^{-1}$  at  $E_{\text{cm}}=8 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0}=125.5 \text{ GeV}$ . See also CHATRCHYAN 13Y.

### $c\bar{c}$ Final State

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;110</b>	95	<sup>1</sup> AABOUD	18M ATLS	$pp$ , 13 TeV

<sup>1</sup> AABOUD 18M use  $36.1 \text{ fb}^{-1}$  at of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . The upper limit on  $\sigma(pp \rightarrow ZH^0) \cdot B(H^0 \rightarrow c\bar{c})$  is 2.7 pb at 95% CL. The quoted values are given for  $m_{H^0} = 125 \text{ GeV}$ .

### $b\bar{b}$ Final State

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>1.02 \pm 0.15</math> OUR AVERAGE</b>			
$1.16^{+0.27}_{-0.25}$	<sup>1</sup> AABOUD	18BN ATLS	$pp \rightarrow H^0 W / H^0 Z, H^0 \rightarrow b\bar{b}$ , 13 TeV, $79.8 \text{ fb}^{-1}$
$1.06 \pm 0.26$	<sup>2</sup> SIRUNYAN	18DB CMS	$pp \rightarrow H^0 W / H^0 Z, H^0 \rightarrow b\bar{b}$ , 13 TeV, $77.2 \text{ fb}^{-1}$
$0.70^{+0.29}_{-0.27}$	<sup>3,4</sup> AAD	16AN LHC	$pp$ , 7, 8 TeV
$1.59^{+0.69}_{-0.72}$	<sup>5</sup> AALTONEN	13M TEVA	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.98^{+0.22}_{-0.21}$	<sup>6</sup> AABOUD	18BN ATLS	$pp \rightarrow H^0 W / H^0 Z, H^0 \rightarrow b\bar{b}$ , 7, 8, 13 TeV
$1.01 \pm 0.20$	<sup>7</sup> AABOUD	18BN ATLS	$pp \rightarrow H^0 X$ , ggF, VBF, $VH^0, t\bar{t}H^0$ 7, 8, 13 TeV
$2.5^{+1.4}_{-1.3}$	<sup>8,9</sup> AABOUD	18BQ ATLS	$pp \rightarrow H^0 X$ , VBF, ggF, $VH^0, t\bar{t}H^0$ , 13 TeV
$3.0^{+1.7}_{-1.6}$	<sup>8,10</sup> AABOUD	18BQ ATLS	$pp \rightarrow H^0 X$ , VBF, 13 TeV
	<sup>11</sup> AALTONEN	18C CDF	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
$1.19^{+0.40}_{-0.38}$	<sup>12</sup> SIRUNYAN	18AE CMS	$pp \rightarrow H^0 W / H^0 Z, H^0 \rightarrow b\bar{b}$ , 13 TeV
$1.06^{+0.31}_{-0.29}$	<sup>13</sup> SIRUNYAN	18AE CMS	$pp \rightarrow H^0 W / H^0 Z, H^0 \rightarrow b\bar{b}$ , 7, 8, 13 TeV
$1.01 \pm 0.22$	<sup>14</sup> SIRUNYAN	18DB CMS	$pp \rightarrow H^0 W / H^0 Z, H^0 \rightarrow b\bar{b}$ , 7, 8, 13 TeV

$1.04 \pm 0.20$	<sup>15</sup> SIRUNYAN	18DB CMS	$pp \rightarrow H^0 X$ , ggF, VBF, $VH^0$ , $t\bar{t}H^0$ 7, 8, 13 TeV
$2.3^{+1.8}_{-1.6}$	<sup>16</sup> SIRUNYAN	18E CMS	$pp \rightarrow H^0 X$ , boosted, 13 TeV
$1.20^{+0.24+0.34}_{-0.23-0.28}$	<sup>17</sup> AABOUD	17BA ATLS	$pp \rightarrow H^0 W/ZX$ , $H^0 \rightarrow$ $b\bar{b}$ , 13 TeV, 36.1 fb <sup>-1</sup>
$0.90 \pm 0.18^{+0.21}_{-0.19}$	<sup>18</sup> AABOUD	17BA ATLS	$pp \rightarrow H^0 W/ZX$ , $H^0 \rightarrow$ $b\bar{b}$ , 7, 8, 13 TeV
$-0.8 \pm 1.3^{+1.8}_{-1.9}$	<sup>19</sup> AABOUD	16X ATLS	$pp \rightarrow H^0 X$ , VBF, 8 TeV
$0.62 \pm 0.37$	<sup>4</sup> AAD	16AN ATLS	$pp$ , 7, 8 TeV
$0.81^{+0.45}_{-0.43}$	<sup>4</sup> AAD	16AN CMS	$pp$ , 7, 8 TeV
$0.63^{+0.31+0.24}_{-0.30-0.23}$	<sup>20</sup> AAD	16K ATLS	$pp$ , 7, 8 TeV
$0.52 \pm 0.32 \pm 0.24$	<sup>21</sup> AAD	15G ATLS	$pp \rightarrow H^0 W/ZX$ , 7, 8 TeV
$2.8^{+1.6}_{-1.4}$	<sup>22</sup> KHACHATRY...15Z	CMS	$pp \rightarrow H^0 X$ , VBF, 8 TeV
$1.03^{+0.44}_{-0.42}$	<sup>23</sup> KHACHATRY...15Z	CMS	$pp$ , 8 TeV, combined
$1.0 \pm 0.5$	<sup>24</sup> CHATRCHYAN14AI	CMS	$pp \rightarrow H^0 W/ZX$ , 7, 8 TeV
$1.72^{+0.92}_{-0.87}$	<sup>25</sup> AALTONEN	13L CDF	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
$1.23^{+1.24}_{-1.17}$	<sup>26</sup> ABAZOV	13L D0	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
$0.5 \pm 2.2$	<sup>27</sup> AAD	12AI ATLS	$pp \rightarrow H^0 W/ZX$ , 7 TeV
	<sup>28</sup> AALTONEN	12T TEVA	$p\bar{p} \rightarrow H^0 W/ZX$ , 1.96 TeV
$0.48^{+0.81}_{-0.70}$	<sup>29</sup> CHATRCHYAN12N	CMS	$pp \rightarrow H^0 W/ZX$ , 7, 8 TeV

<sup>1</sup> AABOUD 18BN search for  $VH^0$ ,  $H^0 \rightarrow b\bar{b}$  ( $V = W, Z$ ) using 79.8 fb<sup>-1</sup> of  $pp$  collision data at  $E_{\text{cm}} = 13$  TeV. The quoted signal strength corresponds to a significance of 4.9 standard deviations and is given for  $m_{H^0} = 125$  GeV.

<sup>2</sup> SIRUNYAN 18DB search for  $VH^0$ ,  $H^0 \rightarrow b\bar{b}$  ( $V = W, Z$ ) using 77.2 fb<sup>-1</sup> of  $pp$  collision data at  $E_{\text{cm}} = 13$  TeV. The quoted signal strength corresponds to a significance of 4.4 standard deviations and is given for  $m_{H^0} = 125.09$  GeV.

<sup>3</sup> AAD 16AN perform fits to the ATLAS and CMS data at  $E_{\text{cm}} = 7$  and 8 TeV. The signal strengths for individual production processes are  $1.0 \pm 0.5$  for  $WH^0$  production,  $0.4 \pm 0.4$  for  $ZH^0$  production, and  $1.1 \pm 1.0$  for  $t\bar{t}H^0$  production.

<sup>4</sup> AAD 16AN: In the fit, relative production cross sections are fixed to those in the Standard Model. The quoted signal strength is given for  $m_{H^0} = 125.09$  GeV.

<sup>5</sup> AALTONEN 13M combine all Tevatron data from the CDF and D0 Collaborations with up to 10.0 fb<sup>-1</sup> and 9.7 fb<sup>-1</sup>, respectively, of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.

<sup>6</sup> AABOUD 18BN combine results of 79.8 fb<sup>-1</sup> at  $E_{\text{cm}} = 13$  TeV with results of  $VH^0$  at  $E_{\text{cm}} = 7$  and 8 TeV.

<sup>7</sup> AABOUD 18BN combine results of  $VH^0$  at  $E_{\text{cm}} = 7, 8$  and 13 TeV with results of VBF (+gluon fusion) and  $t\bar{t}H^0$  at  $E_{\text{cm}} = 7, 8$ , and 13 TeV to perform a search for the  $H^0 \rightarrow b\bar{b}$  decay. The quoted signal strength assumes a SM production strength and corresponds to a significance of 5.4 standard deviations.

- <sup>8</sup> AABOUD 18BQ search for  $H^0 \rightarrow b\bar{b}$  produced through vector-boson fusion (VBF) and VBF+ $\gamma$  with  $30.6 \text{ fb}^{-1}$   $pp$  collision data at  $E_{\text{cm}} = 13 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>9</sup> The signal strength is measured including all production modes (VBF, ggF,  $VH^0$ ,  $t\bar{t}H^0$ ).
- <sup>10</sup> The signal strength is measured for VBF-only and others (ggF,  $VH^0$ ,  $t\bar{t}H^0$ ) are constrained to Standard Model expectations with uncertainties described in their Section VIII B.
- <sup>11</sup> AALTONEN 18C use  $5.4 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . The upper limit at 95% CL on  $p\bar{p} \rightarrow H^0 \rightarrow b\bar{b}$  is 33 times the SM prediction, which corresponds to a cross section of  $40.6 \text{ pb}$ .
- <sup>12</sup> SIRUNYAN 18AE use  $35.9 \text{ fb}^{-1}$  of  $pp$  collision data at  $E_{\text{cm}} = 13 \text{ TeV}$ . The quoted signal strength corresponds to 3.3 standard deviations and is given for  $m_{H^0} = 125.09 \text{ GeV}$ .
- <sup>13</sup> SIRUNYAN 18AE combine the result of  $35.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$  with the results obtained from data of up to  $5.1 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 7 \text{ TeV}$  and up to  $18.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$  (CHATRCHYAN 14AI and KHACHATRYAN 15Z). The quoted signal strength corresponds to 3.8 standard deviations and is given for  $m_{H^0} = 125.09 \text{ GeV}$ .
- <sup>14</sup> SIRUNYAN 18DB combine the result of  $77.2 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$  with the results obtained from data of up to  $5.1 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 7 \text{ TeV}$  and up to  $18.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted signal strength corresponds to a significance of 4.8 standard deviations and is given for  $m_{H^0} = 125.09 \text{ GeV}$ .
- <sup>15</sup> SIRUNYAN 18DB combine results of  $77.2 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$  with results of gluon fusion (ggF), VBF and  $t\bar{t}H^0$  at  $E_{\text{cm}} = 7 \text{ TeV}$ ,  $8 \text{ TeV}$  and  $13 \text{ TeV}$  to perform a search for the  $H^0 \rightarrow b\bar{b}$  decay. The quoted signal strength assumes a SM production strength and corresponds to a significance of 5.6 standard deviations and is given for  $m_{H^0} = 125.09 \text{ GeV}$ .
- <sup>16</sup> SIRUNYAN 18E use  $35.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ . They measure  $\sigma \cdot B$  for gluon fusion production of  $H^0 \rightarrow b\bar{b}$  with  $p_T > 450 \text{ GeV}$ ,  $|\eta| < 2.5$  to be  $74 \pm 48^{+17}_{-10} \text{ fb}$ .
- <sup>17</sup> AABOUD 17BA use  $36.1 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ . They give  $\sigma(WH) \cdot B(H^0 \rightarrow b\bar{b}) = 1.08^{+0.54}_{-0.47} \text{ pb}$  and  $\sigma(ZH) \cdot B(H^0 \rightarrow b\bar{b}) = 0.57^{+0.26}_{-0.23} \text{ pb}$ .
- <sup>18</sup> AABOUD 17BA combine 7, 8 and 13 TeV analyses. The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>19</sup> AABOUD 16X search for vector-boson fusion production of  $H^0$  decaying to  $b\bar{b}$  in  $20.2 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>20</sup> AAD 16K use up to  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and up to  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125.36 \text{ GeV}$ .
- <sup>21</sup> AAD 15G use  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125.36 \text{ GeV}$ .
- <sup>22</sup> KHACHATRYAN 15Z search for vector-boson fusion production of  $H^0$  decaying to  $b\bar{b}$  in up to  $19.8 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>23</sup> KHACHATRYAN 15Z combined vector boson fusion,  $WH^0$ ,  $ZH^0$  production, and  $t\bar{t}H^0$  production results. The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>24</sup> CHATRCHYAN 14AI use up to  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and up to  $18.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ . See also CHATRCHYAN 14AJ.

- <sup>25</sup> AALTONEN 13L combine all CDF results with 9.45–10.0 fb<sup>-1</sup> of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.
- <sup>26</sup> ABAZOV 13L combine all D0 results with up to 9.7 fb<sup>-1</sup> of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.
- <sup>27</sup> AAD 12AI obtain results based on 4.6–4.8 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV. The quoted signal strengths are given in their Fig. 10 for  $m_{H^0} = 126$  GeV. See also Fig. 13 of AAD 12DA.
- <sup>28</sup> AALTONEN 12T combine AALTONEN 12Q, AALTONEN 12R, AALTONEN 12S, ABAZOV 12O, ABAZOV 12P, and ABAZOV 12K. An excess of events over background is observed which is most significant in the region  $m_{H^0} = 120$ –135 GeV, with a local significance of up to 3.3  $\sigma$ . The local significance at  $m_{H^0} = 125$  GeV is 2.8  $\sigma$ , which corresponds to  $(\sigma(H^0 W) + \sigma(H^0 Z)) \cdot \text{B}(H^0 \rightarrow b\bar{b}) = (0.23^{+0.09}_{-0.08})$  pb, compared to the Standard Model expectation at  $m_{H^0} = 125$  GeV of  $0.12 \pm 0.01$  pb. Superseded by AALTONEN 13M.
- <sup>29</sup> CHATRCHYAN 12N obtain results based on 5.0 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}}=7$  TeV and 5.1 fb<sup>-1</sup> at  $E_{\text{cm}}=8$  TeV. The quoted signal strength is given for  $m_{H^0}=125.5$  GeV. See also CHATRCHYAN 13Y.

### $\mu^+ \mu^-$ Final State

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>0.6±0.8 OUR AVERAGE</b>				
1.0±1.0±0.1		1 SIRUNYAN	19E CMS	$pp$ , 7, 8, 13 TeV
-0.1±1.4		2 AABOUD	17Y ATLS	$pp$ , 7, 8, 13 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.7±1.0 <sup>+0.2</sup> <sub>-0.1</sub>		1 SIRUNYAN	19E CMS	$pp$ , 13 TeV, 35.9 fb <sup>-1</sup>
-0.1±1.5		2 AABOUD	17Y ATLS	$pp$ , 13 TeV
0.1±2.5		3 AAD	16AN LHC	$pp$ , 7, 8 TeV
-0.6±3.6		3 AAD	16AN ATLS	$pp$ , 7, 8 TeV
0.9 <sup>+3.6</sup> <sub>-3.5</sub>		3 AAD	16AN CMS	$pp$ , 7, 8 TeV
< 7.4	95	4 KHACHATRY...15H	CMS	$pp \rightarrow H^0 X$ , 7, 8 TeV
< 7.0	95	5 AAD	14AS ATLS	$pp \rightarrow H^0 X$ , 7, 8 TeV

<sup>1</sup> SIRUNYAN 19E search for  $H^0 \rightarrow \mu^+ \mu^-$  using 35.9 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV and combine with results of 7 TeV (5.0 fb<sup>-1</sup>) and 8 TeV (19.7 fb<sup>-1</sup>). The upper limit at 95% CL on the signal strength is 2.9, which corresponds to the SM Higgs boson branching fraction to a muon pair of  $6.4 \times 10^{-4}$ .

<sup>2</sup> AABOUD 17Y use 36.1 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV, 20.3 fb<sup>-1</sup> at 8 TeV and 4.5 fb<sup>-1</sup> at 7 TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.

<sup>3</sup> AAD 16AN: In the fit, relative production cross sections are fixed to those in the Standard Model. The quoted signal strength is given for  $m_{H^0} = 125.09$  GeV.

<sup>4</sup> KHACHATRYAN 15H use 5.0 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and 19.7 fb<sup>-1</sup> at 8 TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.

<sup>5</sup> AAD 14AS search for  $H^0 \rightarrow \mu^+ \mu^-$  in 4.5 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and 20.3 fb<sup>-1</sup> at  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125.5$  GeV.

$\tau^+\tau^-$  Final State

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.11±0.17 OUR AVERAGE</b>			
1.09 <sup>+0.27</sup> <sub>-0.26</sub>	<sup>1</sup> SIRUNYAN	18Y CMS	$p\bar{p}$ , 13 TeV
1.11 <sup>+0.24</sup> <sub>-0.22</sub>	<sup>2,3</sup> AAD	16AN LHC	$p\bar{p}$ , 7, 8 TeV
1.68 <sup>+2.28</sup> <sub>-1.68</sub>	<sup>4</sup> AALTONEN	13M TEVA	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.98±0.18	<sup>5</sup> SIRUNYAN	18Y CMS	$p\bar{p}$ , 7, 8, 13 TeV
2.3 ±1.6	<sup>6</sup> AAD	16AC ATLS	$p\bar{p} \rightarrow H^0 W/Z X$ , 8 TeV
1.41 <sup>+0.40</sup> <sub>-0.36</sub>	<sup>3</sup> AAD	16AN ATLS	$p\bar{p}$ , 7, 8 TeV
0.88 <sup>+0.30</sup> <sub>-0.28</sub>	<sup>3</sup> AAD	16AN CMS	$p\bar{p}$ , 7, 8 TeV
1.44 <sup>+0.30+0.29</sup> <sub>-0.29-0.23</sub>	<sup>7</sup> AAD	16K ATLS	$p\bar{p}$ , 7, 8 TeV
1.43 <sup>+0.27+0.32</sup> <sub>-0.26-0.25</sub> ±0.09	<sup>8</sup> AAD	15AH ATLS	$p\bar{p} \rightarrow H^0 X$ , 7, 8 TeV
0.78±0.27	<sup>9</sup> CHATRCHYAN	14K CMS	$p\bar{p} \rightarrow H^0 X$ , 7, 8 TeV
0.00 <sup>+8.44</sup> <sub>-0.00</sub>	<sup>10</sup> AALTONEN	13L CDF	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
3.96 <sup>+4.11</sup> <sub>-3.38</sub>	<sup>11</sup> ABAZOV	13L D0	$p\bar{p} \rightarrow H^0 X$ , 1.96 TeV
0.4 <sup>+1.6</sup> <sub>-2.0</sub>	<sup>12</sup> AAD	12AI ATLS	$p\bar{p} \rightarrow H^0 X$ , 7 TeV
0.09 <sup>+0.76</sup> <sub>-0.74</sub>	<sup>13</sup> CHATRCHYAN	12N CMS	$p\bar{p} \rightarrow H^0 X$ , 7, 8 TeV

<sup>1</sup> SIRUNYAN 18Y use 35.9 fb<sup>-1</sup> of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 13$  TeV. The quoted signal strength is given for  $m_{H^0} = 125.09$  GeV and corresponds to 4.9 standard deviations.

<sup>2</sup> AAD 16AN perform fits to the ATLAS and CMS data at  $E_{\text{cm}} = 7$  and 8 TeV. The signal strengths for individual production processes are  $1.0 \pm 0.6$  for gluon fusion,  $1.3 \pm 0.4$  for vector boson fusion,  $-1.4 \pm 1.4$  for  $WH^0$  production,  $2.2^{+2.2}_{-1.8}$  for  $ZH^0$  production, and  $-1.9^{+3.7}_{-3.3}$  for  $t\bar{t}H^0$  production.

<sup>3</sup> AAD 16AN: In the fit, relative production cross sections are fixed to those in the Standard Model. The quoted signal strength is given for  $m_{H^0} = 125.09$  GeV.

<sup>4</sup> AALTONEN 13M combine all Tevatron data from the CDF and D0 Collaborations with up to 10.0 fb<sup>-1</sup> and 9.7 fb<sup>-1</sup>, respectively, of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.

<sup>5</sup> SIRUNYAN 18Y combine the result of 35.9 fb<sup>-1</sup> at  $E_{\text{cm}} = 13$  TeV with the results obtained from data of 4.9 fb<sup>-1</sup> at  $E_{\text{cm}} = 7$  TeV and 19.7 fb<sup>-1</sup> at  $E_{\text{cm}} = 8$  TeV (KHACHATRYAN 15AM). The quoted signal strength is given for  $m_{H^0} = 125.09$  GeV and corresponds to 5.9 standard deviations.

<sup>6</sup> AAD 16AC measure the signal strength with  $p\bar{p} \rightarrow H^0 W/Z X$  processes using 20.3 fb<sup>-1</sup> of  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.

<sup>7</sup> AAD 16K use up to 4.7 fb<sup>-1</sup> of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 7$  TeV and up to 20.3 fb<sup>-1</sup> at  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125.36$  GeV.

<sup>8</sup> AAD 15AH use 4.5 fb<sup>-1</sup> of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 7$  TeV and 20.3 fb<sup>-1</sup> at  $E_{\text{cm}} = 8$  TeV. The third uncertainty in the measurement is theory systematics. The signal strength for the gluon fusion mode is  $2.0 \pm 0.8^{+1.2}_{-0.8} \pm 0.3$  and that for vector boson

fusion and  $W/ZH^0$  production modes is  $1.24^{+0.49+0.31}_{-0.45-0.29} \pm 0.08$ . The quoted signal strength is given for  $m_{H^0} = 125.36$  GeV.

- <sup>9</sup> CHATRCHYAN 14K use  $4.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV. See also CHATRCHYAN 14AJ.
- <sup>10</sup> AALTONEN 13L combine all CDF results with  $9.45\text{--}10.0 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.
- <sup>11</sup> ABAZOV 13L combine all D0 results with up to  $9.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV. The quoted signal strength is given for  $m_{H^0} = 125$  GeV.
- <sup>12</sup> AAD 12AI obtain results based on  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV. The quoted signal strengths are given in their Fig. 10 for  $m_{H^0} = 126$  GeV. See also Fig. 13 of AAD 12DA.
- <sup>13</sup> CHATRCHYAN 12N obtain results based on  $4.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}}=7$  TeV and  $5.1 \text{ fb}^{-1}$  at  $E_{\text{cm}}=8$  TeV. The quoted signal strength is given for  $m_{H^0}=125.5$  GeV. See also CHATRCHYAN 13Y .

## Z $\gamma$ Final State

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 6.6</b>	95	<sup>1</sup> AABOUD	17AW ATLS	$pp \rightarrow H^0 X$ , 13 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 7.4	95	<sup>2</sup> SIRUNYAN	18DQ CMS	$pp \rightarrow H^0 X$ , 13 TeV, $H^0 \rightarrow Z\gamma$
<11	95	<sup>3</sup> AAD	14J ATLS	$pp \rightarrow H^0 X$ , 7, 8 TeV
< 9.5	95	<sup>4</sup> CHATRCHYAN 13BK	CMS	$pp \rightarrow H^0 X$ , 7, 8 TeV
<sup>1</sup> AABOUD 17AW search for $H^0 \rightarrow Z\gamma$ , $Z \rightarrow ee, \mu\mu$ in $36.1 \text{ fb}^{-1}$ of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. The quoted signal strength is given for $m_{H^0} = 125.09$ GeV. The upper limit on the branching ratio of $H^0 \rightarrow Z\gamma$ is 1.0% at 95% CL assuming the SM Higgs boson production.				
<sup>2</sup> SIRUNYAN 18DQ search for $H^0 \rightarrow Z\gamma$ , $Z \rightarrow ee, \mu\mu$ in $35.9 \text{ fb}^{-1}$ of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. The quoted signal strength (see their Figs. 6 and 7) is given for $m_{H^0} = 125$ GeV.				
<sup>3</sup> AAD 14J search for $H^0 \rightarrow Z\gamma \rightarrow \ell\ell\gamma$ in $4.5 \text{ fb}^{-1}$ of $pp$ collisions at $E_{\text{cm}} = 7$ TeV and $20.3 \text{ fb}^{-1}$ at $E_{\text{cm}} = 8$ TeV. The quoted signal strength is given for $m_{H^0} = 125.5$ GeV.				
<sup>4</sup> CHATRCHYAN 13BK search for $H^0 \rightarrow Z\gamma \rightarrow \ell\ell\gamma$ in $5.0 \text{ fb}^{-1}$ of $pp$ collisions at $E_{\text{cm}} = 7$ TeV and $19.6 \text{ fb}^{-1}$ at $E_{\text{cm}} = 8$ TeV. A limit on cross section times branching ratio which corresponds to (4–25) times the expected Standard Model cross section is given in the range $m_{H^0} = 120\text{--}160$ GeV at 95% CL. The quoted limit is given for $m_{H^0} = 125$ GeV, where 10 is expected for no signal.				

## $\gamma^*\gamma$ Final State

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<4.0	95	<sup>1</sup> SIRUNYAN	18DQ CMS	$pp \rightarrow H^0 X$ , 13 TeV, $H^0 \rightarrow \gamma^*\gamma$
<6.7	95	<sup>2</sup> KHACHATRY...16B	CMS	$pp$ , 8 TeV, $ee\gamma, \mu\mu\gamma$
<sup>1</sup> SIRUNYAN 18DQ search for $H^0 \rightarrow \gamma^*\gamma, \gamma^* \rightarrow \mu\mu$ in $35.9 \text{ fb}^{-1}$ of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. The mass of $\gamma^*$ is smaller than 50 GeV except in $J/\psi$ and $\Upsilon$ mass regions. The quoted signal strength (see their Figs. 6 and 7) is given for $m_{H^0} = 125$ GeV.				

<sup>2</sup> KHACHATRYAN 16B search for  $H^0 \rightarrow \gamma^* \gamma \rightarrow e^+ e^- \gamma$  and  $\mu^+ \mu^- \gamma$  (with  $m(e^+ e^-) < 3.5$  GeV and  $m(\mu^+ \mu^-) < 20$  GeV) in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 6 for limits on individual channels.

## OTHER $H^0$ PRODUCTION PROPERTIES

### $t\bar{t}H^0$ Production

Signal strength relative to the Standard Model cross section.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>1.28 ± 0.20 OUR AVERAGE</b>				
1.2 ± 0.3		<sup>1</sup> AABOUD	18AC ATLS	$pp, 13 \text{ TeV}, H^0 \rightarrow b\bar{b} \tau\tau, \gamma\gamma, WW^*, ZZ^*$
1.26 <sup>+0.31</sup> <sub>-0.26</sub>		<sup>2</sup> SIRUNYAN	18L CMS	$pp, 7, 8, 13 \text{ TeV}, H^0 \rightarrow b\bar{b}, \tau\tau, \gamma\gamma, WW^*, ZZ^*$
1.9 <sup>+0.8</sup> <sub>-0.7</sub>		<sup>3</sup> AAD	16AN ATLS	$pp, 7, 8 \text{ TeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.6 <sup>+0.5</sup> <sub>-0.4</sub>		<sup>4</sup> AABOUD	18AC ATLS	$pp, 13 \text{ TeV}, H^0 \rightarrow \tau\tau, WW^*, ZZ^*$
		<sup>5</sup> AABOUD	18BK ATLS	$pp, 13 \text{ TeV}, H^0 \rightarrow b\bar{b} \tau\tau, \gamma\gamma, WW^*, ZZ^*$
0.84 <sup>+0.64</sup> <sub>-0.61</sub>		<sup>6</sup> AABOUD	18T ATLS	$pp, 13 \text{ TeV}, H^0 \rightarrow b\bar{b}$
0.9 ± 1.5		<sup>7</sup> SIRUNYAN	18BD CMS	$pp, 13 \text{ TeV}, H^0 \rightarrow b\bar{b}$
1.23 <sup>+0.45</sup> <sub>-0.43</sub>		<sup>8</sup> SIRUNYAN	18BQ CMS	$pp, 13 \text{ TeV}, H^0 \rightarrow \tau\tau, WW^*, ZZ^*$
		<sup>9</sup> SIRUNYAN	18BU CMS	$pp, 13 \text{ TeV}$
1.7 ± 0.8		<sup>10</sup> AAD	16AL ATLS	$pp, 7, 8 \text{ TeV}, H^0 \rightarrow b\bar{b}, \tau\tau, \gamma\gamma, WW^*, \text{ and } ZZ^*$
2.3 <sup>+0.7</sup> <sub>-0.6</sub>		<sup>3,11</sup> AAD	16AN LHC	$pp, 7, 8 \text{ TeV}$
2.9 <sup>+1.0</sup> <sub>-0.9</sub>		<sup>3</sup> AAD	16AN CMS	$pp, 7, 8 \text{ TeV}$
1.81 <sup>+0.52 +0.58 +0.31</sup> <sub>-0.50 -0.55 -0.12</sub>		<sup>12</sup> AAD	16K ATLS	$pp, 7, 8 \text{ TeV}$
1.4 <sup>+2.1 +0.6</sup> <sub>-1.4 -0.3</sub>		<sup>13</sup> AAD	15 ATLS	$pp, 7, 8 \text{ TeV}$
1.5 ± 1.1		<sup>14</sup> AAD	15BC ATLS	$pp, 8 \text{ TeV}$
2.1 <sup>+1.4</sup> <sub>-1.2</sub>		<sup>15</sup> AAD	15T ATLS	$pp, 8 \text{ TeV}$
1.2 <sup>+1.6</sup> <sub>-1.5</sub>		<sup>16</sup> KHACHATRYAN	15AN CMS	$pp, 8 \text{ TeV}$
2.8 <sup>+1.0</sup> <sub>-0.9</sub>		<sup>17</sup> KHACHATRYAN	14H CMS	$pp, 7, 8 \text{ TeV}$
9.49 <sup>+6.60</sup> <sub>-6.28</sub>		<sup>18</sup> AALTONEN	13L CDF	$p\bar{p}, 1.96 \text{ TeV}$
<5.8	95	<sup>19</sup> CHATRCHYAN	13X CMS	$pp, 7, 8 \text{ TeV}, H^0 \rightarrow b\bar{b}$

- <sup>1</sup> AABOUD 18AC combine results of  $t\bar{t}H^0$ ,  $H^0 \rightarrow \tau\tau$ ,  $WW^*(\rightarrow \ell\nu\ell\nu, \ell\nu q\bar{q})$ ,  $ZZ^*(\rightarrow \ell\ell\nu\nu, \ell\ell q\bar{q})$  with results of  $t\bar{t}H^0$ ,  $H^0 \rightarrow b\bar{b}$  (AABOUD 18T),  $\gamma\gamma$  (AABOUD 18BO),  $ZZ^*(\rightarrow 4\ell)$  (AABOUD 18AJ) in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ . See their Table 14.
- <sup>2</sup> SIRUNYAN 18L use up to 5.1, 19.7 and  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7, 8,$  and  $13 \text{ TeV}$ , respectively. The quoted signal strength corresponds to a significance of 5.2 standard deviations and is given for  $m_{H^0} = 125.09 \text{ GeV}$ .  $H^0$  decay channels of  $WW^*$ ,  $ZZ^*$ ,  $\gamma\gamma$ ,  $\tau\tau$ , and  $b\bar{b}$  are used. See their Table 1 and Fig. 2 for results on individual channels.
- <sup>3</sup> AAD 16AN: In the fit, relative branching ratios are fixed to those in the Standard Model. The quoted signal strength is given for  $m_{H^0} = 125.09 \text{ GeV}$ .
- <sup>4</sup> AABOUD 18AC search for  $t\bar{t}H^0$  production with  $H^0$  decaying to  $\tau\tau$ ,  $WW^*(\rightarrow \ell\nu\ell\nu, \ell\nu q\bar{q})$ ,  $ZZ^*(\rightarrow \ell\ell\nu\nu, \ell\ell q\bar{q})$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ . See their Table 13 and Fig. 13.
- <sup>5</sup> AABOUD 18BK use  $79.8 \text{ fb}^{-1}$  data for  $t\bar{t}H^0$  production with  $H^0 \rightarrow \gamma\gamma$  and  $ZZ^* \rightarrow 4\ell$  ( $\ell = e, \mu$ ) and  $36.1 \text{ fb}^{-1}$  for other decay channels at  $E_{\text{cm}} = 13 \text{ TeV}$ . A significance of 5.8 standard deviations is observed for  $m_{H^0} = 125.09 \text{ GeV}$  and its signal strength without the uncertainty of the  $t\bar{t}H^0$  cross section is  $1.32^{+0.28}_{-0.26}$ . Combining with results of 7 and 8 TeV (AAD 16K), the significance is 6.3 standard deviations. Assuming Standard Model branching fractions, the total  $t\bar{t}H^0$  production cross section at 13 TeV is measured to be  $670 \pm 90^{+110}_{-100} \text{ fb}$ .
- <sup>6</sup> AABOUD 18T search for  $t\bar{t}H^0$  production with  $H^0$  decaying to  $b\bar{b}$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>7</sup> SIRUNYAN 18BD search for  $t\bar{t}H^0$ ,  $H^0 \rightarrow b\bar{b}$  in the all-jet final state with  $35.9 \text{ fb}^{-1}$   $pp$  collision data at  $E_{\text{cm}} = 13 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>8</sup> SIRUNYAN 18BQ search for  $t\bar{t}H^0$  in final states with electrons, muons and hadronically decaying  $\tau$  leptons ( $H^0 \rightarrow WW^*, ZZ^*, \tau\tau$ ) with  $35.9 \text{ fb}^{-1}$  of  $pp$  collision data at  $E_{\text{cm}} = 13 \text{ TeV}$ . The quoted signal strength corresponds to a significance of 3.2 standard deviations and is given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>9</sup> SIRUNYAN 18BU search for the production of four top quarks with same-sign and multilepton final states with  $35.9 \text{ fb}^{-1}$   $pp$  collision data at  $E_{\text{cm}} = 13 \text{ TeV}$ . The results constrain the ratio of the top quark Yukawa coupling  $y_t$  to its the Standard Model by comparing to the central value of a theoretical prediction (see their Ref. [16]), yielding  $|y_t/y_t^{SM}| < 2.1$  at 95% CL.
- <sup>10</sup> AAD 16AL search for  $t\bar{t}H^0$  production with  $H^0$  decaying to  $\gamma\gamma$  in  $4.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $b\bar{b}, \tau\tau, \gamma\gamma, WW^*$ , and  $ZZ^*$  in  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ . This paper combines the results of previous papers, and the new result of this paper only is:  $\mu = 1.6 \pm 2.6$ .
- <sup>11</sup> AAD 16AN perform fits to the ATLAS and CMS data at  $E_{\text{cm}} = 7$  and  $8 \text{ TeV}$ .
- <sup>12</sup> AAD 16K use up to  $4.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and up to  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The third uncertainty in the measurement is theory systematics. The quoted signal strength is given for  $m_{H^0} = 125.36 \text{ GeV}$ .
- <sup>13</sup> AAD 15 search for  $t\bar{t}H^0$  production with  $H^0$  decaying to  $\gamma\gamma$  in  $4.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted result on the signal strength is equivalent to an upper limit of 6.7 at 95% CL and is given for  $m_{H^0} = 125.4 \text{ GeV}$ .
- <sup>14</sup> AAD 15BC search for  $t\bar{t}H^0$  production with  $H^0$  decaying to  $b\bar{b}$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . The corresponding upper limit is 3.4 at 95% CL. The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .



- 15 AAD 15T search for  $t\bar{t}H^0$  production with  $H^0$  resulting in multilepton final states (mainly from  $WW^*$ ,  $\tau\tau$ ,  $ZZ^*$ ) in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted result on the signal strength is given for  $m_{H^0} = 125 \text{ GeV}$  and corresponds to an upper limit of 4.7 at 95% CL. The data sample is independent from AAD 15 and AAD 15BC.
- 16 KHACHATRYAN 15AN search for  $t\bar{t}H^0$  production with  $H^0$  decaying to  $b\bar{b}$  in  $19.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted result on the signal strength is equivalent to an upper limit of 4.2 at 95% CL and is given for  $m_{H^0} = 125 \text{ GeV}$ .
- 17 KHACHATRYAN 14H search for  $t\bar{t}H^0$  production with  $H^0$  decaying to  $b\bar{b}$ ,  $\tau\tau$ ,  $\gamma\gamma$ ,  $WW^*$ , and  $ZZ^*$ , in  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125.6 \text{ GeV}$ .
- 18 AALTONEN 13L combine all CDF results with  $9.45\text{--}10.0 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . The quoted signal strength is given for  $m_{H^0} = 125 \text{ GeV}$ .
- 19 CHATRCHYAN 13X search for  $t\bar{t}H^0$  production followed by  $H^0 \rightarrow b\bar{b}$ , one top decaying to  $\ell\nu$  and the other to either  $\ell\nu$  or  $q\bar{q}$  in  $5.0 \text{ fb}^{-1}$  and  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  and  $8 \text{ TeV}$ . A limit on cross section times branching ratio which corresponds to (4.0–8.6) times the expected Standard Model cross section is given for  $m_{H^0} = 110\text{--}140 \text{ GeV}$  at 95% CL. The quoted limit is given for  $m_{H^0} = 125 \text{ GeV}$ , where 5.2 is expected for no signal.

## $H^0 H^0$ Production

The 95% CL limits are for the cross section (CS) and Higgs self coupling ( $\kappa_\lambda$ ) scaling factors both relative to the SM predictions.

CS	$\kappa_\lambda$	DOCUMENT ID	TECN	COMMENT
<b>&lt; 12.7</b>		1 AABOUD	18CQ ATLS	13 TeV, $b\bar{b}\tau\tau$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 12.9		2 AABOUD	19A ATLS	13 TeV, $b\bar{b}b\bar{b}$
< 24	–11 to 17	3 SIRUNYAN	19 CMS	13 TeV, $\gamma\gamma b\bar{b}$
<179		4 SIRUNYAN	19H CMS	13 TeV, $b\bar{b}b\bar{b}$
<230		5 AABOUD	18BU ATLS	13 TeV, $\gamma\gamma WW^*$
< 22	–8.2 to 13.2	6 AABOUD	18CWATLS	13 TeV, $\gamma\gamma b\bar{b}$
< 30		7 SIRUNYAN	18A CMS	13 TeV, $b\bar{b}\tau\tau$
< 79		8 SIRUNYAN	18F CMS	13 TeV, $b\bar{b}\ell\nu\ell\nu$
< 43		9 SIRUNYAN	17CN CMS	8 TeV, $b\bar{b}\tau\tau$ , $\gamma\gamma b\bar{b}$ , $b\bar{b}b\bar{b}$
<108		10 AABOUD	16I ATLS	13 TeV, $b\bar{b}b\bar{b}$
< 74		11 KHACHATRY...	16BQ CMS	8 TeV, $\gamma\gamma b\bar{b}$
< 70		12 AAD	15CE ATLS	8 TeV, $b\bar{b}b\bar{b}$ , $b\bar{b}\tau\tau$ , $\gamma\gamma b\bar{b}$ , $\gamma\gamma WW$

1 AABOUD 18CQ search for  $H^0 H^0$  production using  $H^0 H^0 \rightarrow b\bar{b}\tau\tau$  with data of  $36.1 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The upper limit on the  $pp \rightarrow H^0 H^0 \rightarrow b\bar{b}\tau\tau$  production cross section at 95% is measured to be  $30.9 \text{ fb}$ , which corresponds to about 12.7 times the SM prediction.

2 AABOUD 19A search for  $H^0 H^0$  production using  $H^0 H^0 \rightarrow b\bar{b}b\bar{b}$  with data of  $36.1 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The upper limit on the  $pp \rightarrow H^0 H^0 \rightarrow b\bar{b}b\bar{b}$  production cross section at 95% is measured to be  $147 \text{ fb}$ , which corresponds to about 12.9 times the SM prediction.

3 SIRUNYAN 19 search for  $H^0 H^0$  production using  $H^0 H^0 \rightarrow \gamma\gamma b\bar{b}$  with data of  $35.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The upper limit on the  $pp \rightarrow H^0 H^0 \rightarrow \gamma\gamma b\bar{b}$  production cross section at 95% CL is measured to be  $2.0 \text{ fb}$ , which corresponds to about 24 times the SM prediction. The effective Higgs boson self-coupling  $\kappa_\lambda (= \lambda_{HHH} / \lambda_{HHH}^{SM})$  is constrained to be  $-11 < \kappa_\lambda < 17$  at 95% CL assuming all other Higgs boson couplings are at their SM value.

- <sup>4</sup> SIRUNYAN 19H search for  $H^0 H^0$  production using  $H^0 H^0 \rightarrow b\bar{b}b\bar{b}$ , where one of  $b\bar{b}$  pairs is highly boosted and the other one is resolved, with data of  $35.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The upper limit on the  $pp \rightarrow H^0 H^0 \rightarrow b\bar{b}b\bar{b}$  production cross section at 95% is measured to be 1980 fb, which corresponds to about 179 times the SM prediction.
- <sup>5</sup> AABOUD 18BU search for  $H^0 H^0$  production using  $\gamma\gamma WW^*$  with the final state of  $\gamma\gamma\ell\nu jj$  using data of  $36.1 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The upper limit on the  $pp \rightarrow H^0 H^0$  production cross section at 95% CL is measured to be 7.7 pb, which corresponds to about 230 times the SM prediction. The upper limit on the  $pp \rightarrow H^0 H^0 \rightarrow \gamma\gamma WW^*$  at 95% CL is measured to be 7.5 fb (see their Table 6).
- <sup>6</sup> AABOUD 18CW search for  $H^0 H^0$  production using  $H^0 H^0 \rightarrow \gamma\gamma b\bar{b}$  with data of  $36.1 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The upper limit on the  $pp \rightarrow H^0 H^0$  production cross section at 95% is measured to be 0.73 pb, which corresponds to about 22 times the SM prediction. The effective Higgs boson self-coupling  $\kappa_\lambda$  is constrained to be  $-8.2 < \kappa_\lambda < 13.2$  at 95% CL assuming all other Higgs boson couplings are at their SM value.
- <sup>7</sup> SIRUNYAN 18A search for  $H^0 H^0$  production using  $H^0 H^0 \rightarrow b\bar{b}\tau\tau$  with data of  $35.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The upper limit on the  $gg \rightarrow H^0 H^0 \rightarrow b\bar{b}\tau\tau$  production cross section is measured to be 75.4 fb, which corresponds to about 30 times the SM prediction. Limits on Higgs-boson trilinear coupling  $\lambda_{HHH}$  and top Yukawa coupling  $y_t$  are also given (see their Fig. 6).
- <sup>8</sup> SIRUNYAN 18F search non-resonant for  $H^0 H^0$  production using  $H^0 H^0 \rightarrow b\bar{b}\ell\nu\ell\nu$ , where  $\ell\nu\ell\nu$  is either  $WW \rightarrow \ell\nu\ell\nu$  or  $ZZ \rightarrow \ell\nu\ell\nu$  ( $\ell$  is  $e, \mu$  or a leptonically decaying  $\tau$ ), with data of  $35.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The upper limit on the  $H^0 H^0 \rightarrow b\bar{b}\ell\nu\ell\nu$  production cross section at 95% CL is measured to be 72 fb, which corresponds to about 79 times the SM prediction.
- <sup>9</sup> SIRUNYAN 17CN search for  $H^0 H^0$  production using  $H^0 H^0 \rightarrow b\bar{b}\tau\tau$  with data of  $18.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . Results are then combined with the published results of the  $H^0 H^0 \rightarrow \gamma\gamma b\bar{b}$  and  $H^0 H^0 \rightarrow b\bar{b}b\bar{b}$ , which use data of up to  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The upper limit on the  $gg \rightarrow H^0 H^0$  production cross section is measured to be 0.59 pb from  $b\bar{b}\tau\tau$ , which corresponds to about 59 times the SM prediction (gluon fusion). The combined upper limit is 0.43 pb, which is about 43 times the SM prediction. The quoted values are given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>10</sup> AABOUD 16I search for  $H^0 H^0$  production using  $H^0 H^0 \rightarrow b\bar{b}b\bar{b}$  with data of  $3.2 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . The upper limit on the  $pp \rightarrow H^0 H^0 \rightarrow b\bar{b}b\bar{b}$  production cross section is measured to be 1.22 pb. This result corresponds to about 108 times the SM prediction (gluon fusion), which is  $11.3^{+0.9}_{-1.0} \text{ fb}$  (NNLO+NNLL) including top quark mass effects. The quoted values are given for  $m_{H^0} = 125 \text{ GeV}$ .
- <sup>11</sup> KHACHATRYAN 16BQ search for  $H^0 H^0$  production using  $H^0 H^0 \rightarrow \gamma\gamma b\bar{b}$  with data of  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The upper limit on the  $gg \rightarrow H^0 H^0 \rightarrow \gamma\gamma b\bar{b}$  production is measured to be 1.85 fb, which corresponds to about 74 times the SM prediction and is translated into 0.71 pb for  $gg \rightarrow H^0 H^0$  production cross section. Limits on Higgs-boson trilinear coupling  $\lambda$  are also given.
- <sup>12</sup> AAD 15CE search for  $H^0 H^0$  production using  $H^0 H^0 \rightarrow b\bar{b}\tau\tau$  and  $H^0 H^0 \rightarrow \gamma\gamma WW$  with data of  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . These results are then combined with the published results of the  $H^0 H^0 \rightarrow \gamma\gamma b\bar{b}$  and  $H^0 H^0 \rightarrow b\bar{b}b\bar{b}$ , which use data of up to  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The upper limits on the  $gg \rightarrow H^0 H^0$  production cross section are measured to be 1.6 pb, 11.4 pb, 2.2 pb and 0.62 pb from  $b\bar{b}\tau\tau, \gamma\gamma WW, \gamma\gamma b\bar{b}$  and  $b\bar{b}b\bar{b}$ , respectively. The combined upper limit is 0.69 pb, which corresponds to about 70 times the SM prediction. The quoted results are given for  $m_{H^0} = 125.4 \text{ GeV}$ . See their Table 4.

## $tH^0$ associated production cross section

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

95 <sup>1</sup> KHACHATRY...16AU CMS  $pp$ , 8 TeV

<sup>1</sup> KHACHATRYAN 16AU search for the  $tH^0$  associated production in  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . The 95% CL upper limits on the  $tH^0$  associated production cross section is measured to be 600–1000 fb depending on the assumed  $\gamma\gamma$  branching ratios of the Higgs boson. The  $\gamma\gamma$  branching ratio is varied to be by a factor of 0.5–3.0 of the Standard Model Higgs boson ( $m_{H^0} = 125 \text{ GeV}$ ). The results of the signal strengths for a negative Higgs-boson trilinear coupling are given. The results are given for  $m_{H^0} = 125 \text{ GeV}$ .

## $H^0$ Production Cross Section in $pp$ Collisions at $\sqrt{s} = 13 \text{ TeV}$

Assumes  $m_{H^0} = 125 \text{ GeV}$

VALUE (pb)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

**57.0<sup>+</sup><sub>-</sub> 6.0<sup>+</sup><sub>-</sub> 4.0<sup>+</sup><sub>-</sub> 5.9<sup>+</sup><sub>-</sub> 3.3** <sup>1</sup> AABOUD 18CG ATLS  $pp$ , 13 TeV,  $\gamma\gamma, ZZ^* \rightarrow 4\ell$  ( $\ell = e, \mu$ )

• • • We do not use the following data for averages, fits, limits, etc. • • •

47.9<sup>+</sup><sub>-</sub> 9.1<sup>+</sup><sub>-</sub> 8.6 <sup>1</sup> AABOUD 18CG ATLS  $pp$ , 13 TeV,  $\gamma\gamma$

68 <sup>+</sup><sub>-</sub> 11<sup>+</sup><sub>-</sub> 10 <sup>1</sup> AABOUD 18CG ATLS  $pp$ , 13 TeV,  $ZZ^* \rightarrow 4\ell$  ( $\ell = e, \mu$ )

69 <sup>+</sup><sub>-</sub> 10<sup>+</sup><sub>-</sub> 9  $\pm 5$  <sup>2</sup> AABOUD 17CO ATLS  $pp$ , 13 TeV,  $ZZ^* \rightarrow 4\ell$

<sup>1</sup> AABOUD 18CG use  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ .

<sup>2</sup> AABOUD 17CO use  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$  with  $H^0 \rightarrow ZZ^* \rightarrow 4\ell$  where  $\ell = e, \mu$  for  $m_{H^0} = 125 \text{ GeV}$ . Differential cross sections for the Higgs boson transverse momentum, Higgs boson rapidity, and other related quantities are measured as shown in their Figs. 8 and 9.

## $H^0$ REFERENCES

AABOUD 19A	JHEP 1901 030	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 19F	PL B789 508	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
SIRUNYAN 19	PL B788 7	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN 19E	PRL 122 021801	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN 19H	JHEP 1901 040	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN 19L	JHEP 1901 183	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AABOUD 18	PL B776 318	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18AC	PR D97 072003	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18AJ	JHEP 1803 095	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18AU	JHEP 1807 127	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18BK	PL B784 173	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18BL	PL B786 134	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18BM	PL B784 345	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18BN	PL B786 59	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18BO	PR D98 052005	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18BP	PL B786 223	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18BQ	PR D98 052003	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18BU	EPJ C78 1007	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18CA	JHEP 1810 180	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18CG	PL B786 114	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18CQ	PRL 121 191801	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18CW	JHEP 1811 040	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18M	PRL 120 211802	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD 18T	PR D97 072016	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAIJ 18AM	EPJ C78 1008	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN 18C	PR D98 072002	T. Aaltonen <i>et al.</i>	(CDF Collab.)
SIRUNYAN 18A	PL B778 101	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)

SIRUNYAN	18AE	PL B780 501	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18BD	JHEP 1806 101	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18BH	JHEP 1806 001	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18BQ	JHEP 1808 066	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18BU	EPJ C78 140	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18DB	PRL 121 121801	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18DQ	JHEP 1811 152	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18DS	JHEP 1811 185	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18E	PRL 120 071802	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18F	JHEP 1801 054	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18L	PRL 120 231801	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18Y	PL B779 283	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AABOUD	17AW	JHEP 1710 112	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	17BA	JHEP 1712 024	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	17BD	EPJ C77 765	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	17CO	JHEP 1710 132	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	17Y	PRL 119 051802	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAD	17	EPJ C77 70	G. Aad <i>et al.</i>	(ATLAS Collab.)
KHACHATRY...	17F	JHEP 1702 135	V. Khachatryan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17AM	PL B775 1	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17AV	JHEP 1711 047	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	17CN	PR D96 072004	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AABOUD	16I	PR D94 052002	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	16K	PRL 117 111802	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	16X	JHEP 1611 112	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAD	16	PL B753 69	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16AC	PR D93 092005	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16AF	JHEP 1601 172	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16AL	JHEP 1605 160	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16AN	JHEP 1608 045	G. Aad <i>et al.</i>	(ATLAS and CMS Collabs.)
AAD	16AO	JHEP 1608 104	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16BL	EPJ C76 658	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16K	EPJ C76 6	G. Aad <i>et al.</i>	(ATLAS Collab.)
KHACHATRY...	16AB	PL B759 672	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16AR	JHEP 1604 005	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16AU	JHEP 1606 177	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16B	PL B753 341	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16BA	JHEP 1609 051	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16BQ	PR D94 052012	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16CD	PL B763 472	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16G	EPJ C76 13	V. Khachatryan <i>et al.</i>	(CMS Collab.)
AAD	15	PL B740 222	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15AA	PR D92 012006	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15AH	JHEP 1504 117	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15AQ	JHEP 1508 137	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15AX	EPJ C75 231	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15B	PRL 114 191803	G. Aad <i>et al.</i>	(ATLAS and CMS Collabs.)
AAD	15BC	EPJ C75 349	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15BD	EPJ C75 337	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15BE	EPJ C75 335	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15CE	PR D92 092004	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15CI	EPJ C75 476	G. Aad <i>et al.</i>	(ATLAS Collab.)
Also		EPJ C76 152 (errat.)	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15F	PR D91 012006	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15G	JHEP 1501 069	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15I	PRL 114 121801	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15P	PRL 115 091801	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15T	PL B749 519	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	15	PRL 114 151802	T. Aaltonen <i>et al.</i>	(CDF and D0 Collabs.)
AALTONEN	15B	PRL 114 141802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
KHACHATRY...	15AM	EPJ C75 212	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15AN	EPJ C75 251	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15BA	PR D92 072010	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15H	PL B744 184	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15Q	PL B749 337	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15Y	PR D92 012004	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	15Z	PR D92 032008	V. Khachatryan <i>et al.</i>	(CMS Collab.)
AAD	14AR	PL B738 234	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	14AS	PL B738 68	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	14BC	PR D90 112015	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	14BJ	JHEP 1409 112	G. Aad <i>et al.</i>	(ATLAS Collab.)

AAD	14J	PL B732 8	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	14O	PRL 112 201802	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	14W	PR D90 052004	G. Aad <i>et al.</i>	(ATLAS Collab.)
ABAZOV	14F	PRL 113 161802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	14AA	PR D89 092007	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	14AI	PR D89 012003	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	14AJ	NATP 10 557	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	14B	EPJ C74 2980	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	14G	JHEP 1401 096	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	14K	JHEP 1405 104	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	14D	PL B736 64	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	14H	JHEP 1409 087	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	14P	EPJ C74 3076	V. Khachatryan <i>et al.</i>	(CMS Collab.)
AAD	13AJ	PL B726 120	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	13AK	PL B726 88	G. Aad <i>et al.</i>	(ATLAS Collab.)
Also		PL B734 406 (errat.)	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	13L	PR D88 052013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	13M	PR D88 052014	T. Aaltonen <i>et al.</i>	(CDF and D0 Collabs.)
ABAZOV	13L	PR D88 052011	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	13BK	PL B726 587	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13J	PRL 110 081803	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13X	JHEP 1305 145	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13Y	JHEP 1306 081	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
HEINEMEYER	13A	arXiv:1307.1347	S. Heinemeyer <i>et al.</i>	(LHC Higgs CS Working Group)
AAD	12AI	PL B716 1	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12DA	SCI 338 1576	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	12Q	PRL 109 111803	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	12R	PRL 109 111804	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	12S	PRL 109 111805	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	12T	PRL 109 071804	T. Aaltonen <i>et al.</i>	(CDF and D0 Collabs.)
ABAZOV	12K	PL B716 285	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	12O	PRL 109 121803	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	12P	PRL 109 121804	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	12BY	SCI 338 1569	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	12N	PL B716 30	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
DITTMAYER	12	arXiv:1201.3084	S. Dittmaier <i>et al.</i>	(LHC Higgs CS Working Group)
DITTMAYER	11	arXiv:1101.0593	S. Dittmaier <i>et al.</i>	(LHC Higgs CS Working Group)