

$$I^G(J^{PC}) = 0^-(1^{--})$$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

$\psi(2S)$ MASS

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------------------|------|--|
| 3686.097 ± 0.025 OUR FIT | | Error includes scale factor of 2.6. | | |
| 3686.097 ± 0.010 OUR AVERAGE | | | | |
| 3686.099 ± 0.004 ± 0.009 | | ¹ ANASHIN | 15 | KEDR $e^+e^- \rightarrow$ hadrons |
| 3686.12 ± 0.06 ± 0.10 | 4k | AAIJ | 12H | LHCB $pp \rightarrow J/\psi \pi^+ \pi^- X$ |
| 3685.95 ± 0.10 | 413 | ² ARTAMONOV | 00 | OLYA $e^+e^- \rightarrow$ hadrons |
| 3685.98 ± 0.09 ± 0.04 | | ³ ARMSTRONG | 93B | E760 $\bar{p}p \rightarrow e^+e^-$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 3686.114 ± 0.007 ^{+0.011} _{-0.016} | | ⁴ ANASHIN | 12 | KEDR $e^+e^- \rightarrow$ hadrons |
| 3686.111 ± 0.025 ± 0.009 | | AULCHENKO | 03 | KEDR $e^+e^- \rightarrow$ hadrons |
| 3686.00 ± 0.10 | 413 | ⁵ ZHOLENTZ | 80 | OLYA e^+e^- |

¹ Supersedes AULCHENKO 03 and ANASHIN 12.

² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $J/\psi(1S)$ mass from AULCHENKO 03.

⁴ From the scans in 2004 and 2006. ANASHIN 12 reports the value $3686.114 \pm 0.007 \pm 0.011^{+0.002}_{-0.012}$ MeV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

⁵ Superseded by ARTAMONOV 00.

$m_{\psi(2S)} - m_{J/\psi(1S)}$

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|------|---|
| 589.188 ± 0.028 OUR AVERAGE | | | |
| 589.194 ± 0.027 ± 0.011 | ¹ AULCHENKO | 03 | KEDR $e^+e^- \rightarrow$ hadrons |
| 589.7 ± 1.2 | LEMOIGNE | 82 | GOLI $185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$ |
| 589.07 ± 0.13 | ¹ ZHOLENTZ | 80 | OLYA e^+e^- |
| 588.7 ± 0.8 | LUTH | 75 | MRK1 |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 588 ± 1 | ² BAI | 98E | BES e^+e^- |

¹ Redundant with data in mass above.

² Systematic errors not evaluated.

$\psi(2S)$ WIDTH

| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|------|------------------------|----------|--|
| 294 ± 8 OUR FIT | | | | |
| 286 ± 16 OUR AVERAGE | | | | |
| 358 ± 88 ± 4 | | ABLIKIM | 08B BES2 | $e^+e^- \rightarrow \text{hadrons}$ |
| 290 ± 25 ± 4 | 2.7k | ANDREOTTI | 07 E835 | $\rho\bar{p} \rightarrow e^+e^-, J/\psi X$ |
| 331 ± 58 ± 2 | | ABLIKIM | 06L BES2 | $e^+e^- \rightarrow \text{hadrons}$ |
| 264 ± 27 | | ¹ BAI | 02B BES2 | e^+e^- |
| 287 ± 37 ± 16 | | ² ARMSTRONG | 93B E760 | $\bar{p}p \rightarrow e^+e^-$ |

¹ From a simultaneous fit to the hadronic and $\mu^+\mu^-$ cross section, assuming $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$ and lepton universality. Does not include vacuum polarization correction.

² The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

$\psi(2S)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|---|------------------------------------|-----------------------------------|
| Γ_1 hadrons | (97.85 ± 0.13) % | |
| Γ_2 virtual $\gamma \rightarrow$ hadrons | (1.73 ± 0.14) % | S=1.5 |
| Γ_3 ggg | (10.6 ± 1.6) % | |
| Γ_4 γgg | (1.03 ± 0.29) % | |
| Γ_5 light hadrons | (15.4 ± 1.5) % | |
| Γ_6 e^+e^- | (7.93 ± 0.17) × 10 ⁻³ | |
| Γ_7 $\mu^+\mu^-$ | (8.0 ± 0.6) × 10 ⁻³ | |
| Γ_8 $\tau^+\tau^-$ | (3.1 ± 0.4) × 10 ⁻³ | |

Decays into $J/\psi(1S)$ and anything

| | |
|--------------------------------------|--------------------------------------|
| Γ_9 $J/\psi(1S)$ anything | (61.4 ± 0.6) % |
| Γ_{10} $J/\psi(1S)$ neutrals | (25.38 ± 0.32) % |
| Γ_{11} $J/\psi(1S)\pi^+\pi^-$ | (34.68 ± 0.30) % |
| Γ_{12} $J/\psi(1S)\pi^0\pi^0$ | (18.24 ± 0.31) % |
| Γ_{13} $J/\psi(1S)\eta$ | (3.37 ± 0.05) % |
| Γ_{14} $J/\psi(1S)\pi^0$ | (1.268 ± 0.032) × 10 ⁻³ |

Hadronic decays

| | | |
|--|------------------------------------|--------|
| Γ_{15} $\pi^0 h_c(1P)$ | (8.6 ± 1.3) × 10 ⁻⁴ | |
| Γ_{16} $3(\pi^+\pi^-\pi^0)$ | (3.5 ± 1.6) × 10 ⁻³ | |
| Γ_{17} $2(\pi^+\pi^-\pi^0)$ | (2.9 ± 1.0) × 10 ⁻³ | S=4.7 |
| Γ_{18} $\rho a_2(1320)$ | (2.6 ± 0.9) × 10 ⁻⁴ | |
| Γ_{19} $\pi^+\pi^-\pi^0\pi^0\pi^0$ | (5.3 ± 0.9) × 10 ⁻³ | |
| Γ_{20} $\rho^\pm\pi^\mp\pi^0\pi^0$ | < 2.7 × 10 ⁻³ | CL=90% |
| Γ_{21} $\rho\bar{p}$ | (2.94 ± 0.08) × 10 ⁻⁴ | |
| Γ_{22} $n\bar{n}$ | (3.06 ± 0.15) × 10 ⁻⁴ | |
| Γ_{23} $\Delta^{++}\bar{\Delta}^{--}$ | (1.28 ± 0.35) × 10 ⁻⁴ | |
| Γ_{24} $\Lambda\bar{\Lambda}\pi^0$ | < 2.9 × 10 ⁻⁶ | CL=90% |
| Γ_{25} $\Lambda\bar{\Lambda}\eta$ | (2.5 ± 0.4) × 10 ⁻⁵ | |

| | | | |
|---------------|--|--|--------|
| Γ_{26} | $\Lambda \bar{p} K^+$ | $(1.00 \pm 0.14) \times 10^{-4}$ | |
| Γ_{27} | $\Lambda \bar{p} K^+ \pi^+ \pi^-$ | $(1.8 \pm 0.4) \times 10^{-4}$ | |
| Γ_{28} | $\Lambda \bar{\Lambda} \pi^+ \pi^-$ | $(2.8 \pm 0.6) \times 10^{-4}$ | |
| Γ_{29} | $\Lambda \bar{\Lambda}$ | $(3.81 \pm 0.13) \times 10^{-4}$ | S=1.4 |
| Γ_{30} | $\Lambda \bar{\Sigma}^+ \pi^- + \text{c.c.}$ | $(1.40 \pm 0.13) \times 10^{-4}$ | |
| Γ_{31} | $\Lambda \bar{\Sigma}^- \pi^+ + \text{c.c.}$ | $(1.54 \pm 0.14) \times 10^{-4}$ | |
| Γ_{32} | $\Lambda \bar{\Sigma}^0$ | $(1.23 \pm 0.24) \times 10^{-5}$ | |
| Γ_{33} | $\Sigma^0 \bar{p} K^+ + \text{c.c.}$ | $(1.67 \pm 0.18) \times 10^{-5}$ | |
| Γ_{34} | $\Sigma^+ \bar{\Sigma}^-$ | $(2.32 \pm 0.12) \times 10^{-4}$ | |
| Γ_{35} | $\Sigma^0 \bar{\Sigma}^0$ | $(2.35 \pm 0.09) \times 10^{-4}$ | S=1.1 |
| Γ_{36} | $\Sigma(1385)^+ \bar{\Sigma}(1385)^-$ | $(8.5 \pm 0.7) \times 10^{-5}$ | |
| Γ_{37} | $\Sigma(1385)^- \bar{\Sigma}(1385)^+$ | $(8.5 \pm 0.8) \times 10^{-5}$ | |
| Γ_{38} | $\Sigma(1385)^0 \bar{\Sigma}(1385)^0$ | $(6.9 \pm 0.7) \times 10^{-5}$ | |
| Γ_{39} | $\Xi^- \bar{\Xi}^+$ | $(2.87 \pm 0.11) \times 10^{-4}$ | S=1.1 |
| Γ_{40} | $\Xi^0 \bar{\Xi}^0$ | $(2.3 \pm 0.4) \times 10^{-4}$ | S=4.2 |
| Γ_{41} | $\Xi(1530)^0 \bar{\Xi}(1530)^0$ | $(5.2 \begin{smallmatrix} +3.2 \\ -1.2 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{42} | $K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$ | $(3.9 \pm 0.4) \times 10^{-5}$ | |
| Γ_{43} | $\Xi(1690)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ +$ | $(5.2 \pm 1.6) \times 10^{-6}$ | |
| Γ_{44} | $\begin{smallmatrix} \text{c.c.} \\ \Xi(1820)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ + \end{smallmatrix}$ | $(1.20 \pm 0.32) \times 10^{-5}$ | |
| Γ_{45} | $K^- \bar{\Sigma}^0 \bar{\Xi}^+ + \text{c.c.}$ | $(3.7 \pm 0.4) \times 10^{-5}$ | |
| Γ_{46} | $\Omega^- \bar{\Omega}^+$ | $(5.2 \pm 0.4) \times 10^{-5}$ | |
| Γ_{47} | $\pi^0 \rho \bar{p}$ | $(1.53 \pm 0.07) \times 10^{-4}$ | |
| Γ_{48} | $N(940) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(6.4 \begin{smallmatrix} +1.8 \\ -1.3 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{49} | $N(1440) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(7.3 \begin{smallmatrix} +1.7 \\ -1.5 \end{smallmatrix}) \times 10^{-5}$ | S=2.5 |
| Γ_{50} | $N(1520) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(6.4 \begin{smallmatrix} +2.3 \\ -1.8 \end{smallmatrix}) \times 10^{-6}$ | |
| Γ_{51} | $N(1535) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(2.5 \pm 1.0) \times 10^{-5}$ | |
| Γ_{52} | $N(1650) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(3.8 \begin{smallmatrix} +1.4 \\ -1.7 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{53} | $N(1720) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(1.79 \begin{smallmatrix} +0.26 \\ -0.70 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{54} | $N(2300) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(2.6 \begin{smallmatrix} +1.2 \\ -0.7 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{55} | $N(2570) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(2.13 \begin{smallmatrix} +0.40 \\ -0.31 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{56} | $\pi^0 f_0(2100) \rightarrow \pi^0 \rho \bar{p}$ | $(1.1 \pm 0.4) \times 10^{-5}$ | |
| Γ_{57} | $\eta \rho \bar{p}$ | $(6.0 \pm 0.4) \times 10^{-5}$ | |
| Γ_{58} | $\eta f_0(2100) \rightarrow \eta \rho \bar{p}$ | $(1.2 \pm 0.4) \times 10^{-5}$ | |
| Γ_{59} | $N(1535) \bar{p} \rightarrow \eta \rho \bar{p}$ | $(4.4 \pm 0.7) \times 10^{-5}$ | |
| Γ_{60} | $\omega \rho \bar{p}$ | $(6.9 \pm 2.1) \times 10^{-5}$ | |
| Γ_{61} | $\eta' \rho \bar{p}$ | $(1.10 \pm 0.13) \times 10^{-5}$ | |
| Γ_{62} | $\phi \rho \bar{p}$ | $< 2.4 \times 10^{-5}$ | CL=90% |
| Γ_{63} | $\pi^+ \pi^- \rho \bar{p}$ | $(6.0 \pm 0.4) \times 10^{-4}$ | |

| | | | |
|----------------|---|------------------------------------|--------|
| Γ_{64} | $\rho\bar{n}\pi^-$ or c.c. | $(2.48 \pm 0.17) \times 10^{-4}$ | |
| Γ_{65} | $\rho\bar{n}\pi^-\pi^0$ | $(3.2 \pm 0.7) \times 10^{-4}$ | |
| Γ_{66} | $2(\pi^+\pi^-\pi^0)$ | $(4.8 \pm 1.5) \times 10^{-3}$ | |
| Γ_{67} | $\eta\pi^+\pi^-$ | $< 1.6 \times 10^{-4}$ | CL=90% |
| Γ_{68} | $\eta\pi^+\pi^-\pi^0$ | $(9.5 \pm 1.7) \times 10^{-4}$ | |
| Γ_{69} | $2(\pi^+\pi^-\eta)$ | $(1.2 \pm 0.6) \times 10^{-3}$ | |
| Γ_{70} | $\pi^+\pi^-\pi^0\pi^0\eta$ | $< 4 \times 10^{-4}$ | CL=90% |
| Γ_{71} | $\eta'\pi^+\pi^-\pi^0$ | $(4.5 \pm 2.1) \times 10^{-4}$ | |
| Γ_{72} | $\omega\pi^+\pi^-$ | $(7.3 \pm 1.2) \times 10^{-4}$ | S=2.1 |
| Γ_{73} | $b_1^\pm\pi^\mp$ | $(4.0 \pm 0.6) \times 10^{-4}$ | S=1.1 |
| Γ_{74} | $b_1^0\pi^0$ | $(2.4 \pm 0.6) \times 10^{-4}$ | |
| Γ_{75} | $\omega f_2(1270)$ | $(2.2 \pm 0.4) \times 10^{-4}$ | |
| Γ_{76} | $\omega\pi^0\pi^0$ | $(1.11 \pm 0.35) \times 10^{-3}$ | |
| Γ_{77} | $\pi^0\pi^0 K^+ K^-$ | $(2.6 \pm 1.3) \times 10^{-4}$ | |
| Γ_{78} | $\pi^+\pi^- K^+ K^-$ | $(7.3 \pm 0.5) \times 10^{-4}$ | |
| Γ_{79} | $\pi^0\pi^0 K_S^0 K_L^0$ | $(1.3 \pm 0.6) \times 10^{-3}$ | |
| Γ_{80} | $\rho^0 K^+ K^-$ | $(2.2 \pm 0.4) \times 10^{-4}$ | |
| Γ_{81} | $K^*(892)^0 \bar{K}_2^*(1430)^0$ | $(1.9 \pm 0.5) \times 10^{-4}$ | |
| Γ_{82} | $K^+ K^- \pi^+ \pi^- \eta$ | $(1.3 \pm 0.7) \times 10^{-3}$ | |
| Γ_{83} | $K^+ K^- 2(\pi^+ \pi^-) \pi^0$ | $(1.00 \pm 0.31) \times 10^{-3}$ | |
| Γ_{84} | $K^+ K^- 2(\pi^+ \pi^-)$ | $(1.9 \pm 0.9) \times 10^{-3}$ | |
| Γ_{85} | $K_1(1270)^\pm K^\mp$ | $(1.00 \pm 0.28) \times 10^{-3}$ | |
| Γ_{86} | $K_S^0 K_S^0 \pi^+ \pi^-$ | $(2.2 \pm 0.4) \times 10^{-4}$ | |
| Γ_{87} | $\rho^0 p\bar{p}$ | $(5.0 \pm 2.2) \times 10^{-5}$ | |
| Γ_{88} | $K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$ | $(6.7 \pm 2.5) \times 10^{-4}$ | |
| Γ_{89} | $2(\pi^+ \pi^-)$ | $(2.4 \pm 0.6) \times 10^{-4}$ | S=2.2 |
| Γ_{90} | $\rho^0 \pi^+ \pi^-$ | $(2.2 \pm 0.6) \times 10^{-4}$ | S=1.4 |
| Γ_{91} | $K^+ K^- \pi^+ \pi^- \pi^0$ | $(1.26 \pm 0.09) \times 10^{-3}$ | |
| Γ_{92} | $\omega f_0(1710) \rightarrow \omega K^+ K^-$ | $(5.9 \pm 2.2) \times 10^{-5}$ | |
| Γ_{93} | $K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}$ | $(8.6 \pm 2.2) \times 10^{-4}$ | |
| Γ_{94} | $K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}$ | $(9.6 \pm 2.8) \times 10^{-4}$ | |
| Γ_{95} | $K^*(892)^+ K^- \rho^0 + \text{c.c.}$ | $(7.3 \pm 2.6) \times 10^{-4}$ | |
| Γ_{96} | $K^*(892)^0 K^- \rho^+ + \text{c.c.}$ | $(6.1 \pm 1.8) \times 10^{-4}$ | |
| Γ_{97} | $\eta K^+ K^-$, no $\eta\phi$ | $(3.1 \pm 0.4) \times 10^{-5}$ | |
| Γ_{98} | $\omega K^+ K^-$ | $(1.62 \pm 0.11) \times 10^{-4}$ | S=1.1 |
| Γ_{99} | $\omega K^*(892)^+ K^- + \text{c.c.}$ | $(2.07 \pm 0.26) \times 10^{-4}$ | |
| Γ_{100} | $\omega K_2^*(1430)^+ K^- + \text{c.c.}$ | $(6.1 \pm 1.2) \times 10^{-5}$ | |
| Γ_{101} | $\omega \bar{K}^*(892)^0 K^0$ | $(1.68 \pm 0.30) \times 10^{-4}$ | |
| Γ_{102} | $\omega \bar{K}_2^*(1430)^0 K^0$ | $(5.8 \pm 2.2) \times 10^{-5}$ | |
| Γ_{103} | $\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$ | $(1.6 \pm 0.4) \times 10^{-5}$ | |
| Γ_{104} | $\omega X(1440) \rightarrow \omega K^+ K^- \pi^0$ | $(1.09 \pm 0.26) \times 10^{-5}$ | |
| Γ_{105} | $\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$ | $(3.0 \pm 1.0) \times 10^{-6}$ | |

| | | | |
|----------------|---|---|--------|
| Γ_{106} | $\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0$ | $(1.2 \pm 0.7) \times 10^{-6}$ | |
| Γ_{107} | $3(\pi^+ \pi^-)$ | $(3.5 \pm 2.0) \times 10^{-4}$ | S=2.8 |
| Γ_{108} | $\rho \bar{\rho} \pi^+ \pi^- \pi^0$ | $(7.3 \pm 0.7) \times 10^{-4}$ | |
| Γ_{109} | $K^+ K^-$ | $(7.5 \pm 0.5) \times 10^{-5}$ | |
| Γ_{110} | $K_S^0 K_L^0$ | $(5.34 \pm 0.33) \times 10^{-5}$ | |
| Γ_{111} | $\pi^+ \pi^- \pi^0$ | $(2.01 \pm 0.17) \times 10^{-4}$ | S=1.7 |
| Γ_{112} | $\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$ | $(1.9 \begin{smallmatrix} +1.2 \\ -0.4 \end{smallmatrix}) \times 10^{-4}$ | |
| Γ_{113} | $\rho(770)\pi \rightarrow \pi^+ \pi^- \pi^0$ | $(3.2 \pm 1.2) \times 10^{-5}$ | S=1.8 |
| Γ_{114} | $\pi^+ \pi^-$ | $(7.8 \pm 2.6) \times 10^{-6}$ | |
| Γ_{115} | $K_1(1400)^\pm K^\mp$ | $< 3.1 \times 10^{-4}$ | CL=90% |
| Γ_{116} | $K_2^*(1430)^\pm K^\mp$ | $(7.1 \begin{smallmatrix} +1.3 \\ -0.9 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{117} | $K^+ K^- \pi^0$ | $(4.07 \pm 0.31) \times 10^{-5}$ | |
| Γ_{118} | $K_S^0 K_L^0 \pi^0$ | $< 3.0 \times 10^{-4}$ | CL=90% |
| Γ_{119} | $K_S^0 K_L^0 \eta$ | $(1.3 \pm 0.5) \times 10^{-3}$ | |
| Γ_{120} | $K^+ K^*(892)^- + \text{c.c.}$ | $(2.9 \pm 0.4) \times 10^{-5}$ | S=1.2 |
| Γ_{121} | $K^*(892)^0 \bar{K}^0 + \text{c.c.}$ | $(1.09 \pm 0.20) \times 10^{-4}$ | |
| Γ_{122} | $\phi \pi^+ \pi^-$ | $(1.18 \pm 0.26) \times 10^{-4}$ | S=1.5 |
| Γ_{123} | $\phi f_0(980) \rightarrow \pi^+ \pi^-$ | $(7.5 \pm 3.3) \times 10^{-5}$ | S=1.6 |
| Γ_{124} | $2(K^+ K^-)$ | $(6.3 \pm 1.3) \times 10^{-5}$ | |
| Γ_{125} | $\phi K^+ K^-$ | $(7.0 \pm 1.6) \times 10^{-5}$ | |
| Γ_{126} | $2(K^+ K^-) \pi^0$ | $(1.10 \pm 0.28) \times 10^{-4}$ | |
| Γ_{127} | $\phi \eta$ | $(3.10 \pm 0.31) \times 10^{-5}$ | |
| Γ_{128} | $\eta \phi(2170), \phi(2170) \rightarrow$ $\phi f_0(980), f_0 \rightarrow \pi^+ \pi^-$ | $< 2.2 \times 10^{-6}$ | CL=90% |
| Γ_{129} | $\phi \eta'$ | $(3.1 \pm 1.6) \times 10^{-5}$ | |
| Γ_{130} | $\omega \eta'$ | $(3.2 \begin{smallmatrix} +2.5 \\ -2.1 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{131} | $\omega \pi^0$ | $(2.1 \pm 0.6) \times 10^{-5}$ | |
| Γ_{132} | $\rho \eta'$ | $(1.9 \begin{smallmatrix} +1.7 \\ -1.2 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{133} | $\rho \eta$ | $(2.2 \pm 0.6) \times 10^{-5}$ | S=1.1 |
| Γ_{134} | $\omega \eta$ | $< 1.1 \times 10^{-5}$ | CL=90% |
| Γ_{135} | $\phi \pi^0$ | $< 4 \times 10^{-7}$ | CL=90% |
| Γ_{136} | $\eta_c \pi^+ \pi^- \pi^0$ | $< 1.0 \times 10^{-3}$ | CL=90% |
| Γ_{137} | $\rho \bar{\rho} K^+ K^-$ | $(2.7 \pm 0.7) \times 10^{-5}$ | |
| Γ_{138} | $\bar{\Lambda} n K_S^0 + \text{c.c.}$ | $(8.1 \pm 1.8) \times 10^{-5}$ | |
| Γ_{139} | $\phi f_2'(1525)$ | $(4.4 \pm 1.6) \times 10^{-5}$ | |
| Γ_{140} | $\Theta(1540) \bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + \text{c.c.}$ | $< 8.8 \times 10^{-6}$ | CL=90% |
| Γ_{141} | $\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$ | $< 1.0 \times 10^{-5}$ | CL=90% |
| Γ_{142} | $\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$ | $< 7.0 \times 10^{-6}$ | CL=90% |
| Γ_{143} | $\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$ | $< 2.6 \times 10^{-5}$ | CL=90% |

| | | | | |
|----------------|--|---------|------------------|--------|
| Γ_{144} | $\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$ | < 6.0 | $\times 10^{-6}$ | CL=90% |
| Γ_{145} | $K_S^0 K_S^0$ | < 4.6 | $\times 10^{-6}$ | |
| Γ_{146} | $\Lambda_c^+ \bar{p} e^+ e^- + \text{c.c.}$ | < 1.7 | $\times 10^{-6}$ | CL=90% |

Radiative decays

| | | | | |
|----------------|--|--|------------------|--------|
| Γ_{147} | $\gamma \chi_{c0}(1P)$ | $(9.79 \pm 0.20) \%$ | | |
| Γ_{148} | $\gamma \chi_{c1}(1P)$ | $(9.75 \pm 0.24) \%$ | | |
| Γ_{149} | $\gamma \chi_{c2}(1P)$ | $(9.52 \pm 0.20) \%$ | | |
| Γ_{150} | $\gamma \eta_c(1S)$ | $(3.4 \pm 0.5) \times 10^{-3}$ | | S=1.3 |
| Γ_{151} | $\gamma \eta_c(2S)$ | $(7 \pm 5) \times 10^{-4}$ | | |
| Γ_{152} | $\gamma \pi^0$ | $(1.04 \pm 0.22) \times 10^{-6}$ | | S=1.4 |
| Γ_{153} | $\gamma \eta'(958)$ | $(1.24 \pm 0.04) \times 10^{-4}$ | | |
| Γ_{154} | $\gamma f_2(1270)$ | $(2.73 \begin{smallmatrix} +0.29 \\ -0.25 \end{smallmatrix}) \times 10^{-4}$ | | S=1.8 |
| Γ_{155} | $\gamma f_0(1370) \rightarrow \gamma K \bar{K}$ | $(3.1 \pm 1.7) \times 10^{-5}$ | | |
| Γ_{156} | $\gamma f_0(1500)$ | $(9.3 \pm 1.9) \times 10^{-5}$ | | |
| Γ_{157} | $\gamma f_2'(1525)$ | $(3.3 \pm 0.8) \times 10^{-5}$ | | |
| Γ_{158} | $\gamma f_0(1710)$ | | | |
| Γ_{159} | $\gamma f_0(1710) \rightarrow \gamma \pi \pi$ | $(3.5 \pm 0.6) \times 10^{-5}$ | | |
| Γ_{160} | $\gamma f_0(1710) \rightarrow \gamma K \bar{K}$ | $(6.6 \pm 0.7) \times 10^{-5}$ | | |
| Γ_{161} | $\gamma f_0(2100) \rightarrow \gamma \pi \pi$ | $(4.8 \pm 1.0) \times 10^{-6}$ | | |
| Γ_{162} | $\gamma f_0(2200) \rightarrow \gamma K \bar{K}$ | $(3.2 \pm 1.0) \times 10^{-6}$ | | |
| Γ_{163} | $\gamma f_J(2220) \rightarrow \gamma \pi \pi$ | < 5.8 | $\times 10^{-6}$ | CL=90% |
| Γ_{164} | $\gamma f_J(2220) \rightarrow \gamma K \bar{K}$ | < 9.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{165} | $\gamma \gamma$ | < 1.5 | $\times 10^{-4}$ | CL=90% |
| Γ_{166} | $\gamma \eta$ | $(9.2 \pm 1.8) \times 10^{-7}$ | | |
| Γ_{167} | $\gamma \eta \pi^+ \pi^-$ | $(8.7 \pm 2.1) \times 10^{-4}$ | | |
| Γ_{168} | $\gamma \eta(1405)$ | | | |
| Γ_{169} | $\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi$ | < 9 | $\times 10^{-5}$ | CL=90% |
| Γ_{170} | $\gamma \eta(1405) \rightarrow \eta \pi^+ \pi^-$ | $(3.6 \pm 2.5) \times 10^{-5}$ | | |
| Γ_{171} | $\gamma \eta(1405) \rightarrow \gamma f_0(980) \pi^0 \rightarrow \gamma \pi^+ \pi^- \pi^0$ | < 5.0 | $\times 10^{-7}$ | CL=90% |
| Γ_{172} | $\gamma \eta(1475)$ | | | |
| Γ_{173} | $\gamma \eta(1475) \rightarrow K \bar{K} \pi$ | < 1.4 | $\times 10^{-4}$ | CL=90% |
| Γ_{174} | $\gamma \eta(1475) \rightarrow \eta \pi^+ \pi^-$ | < 8.8 | $\times 10^{-5}$ | CL=90% |
| Γ_{175} | $\gamma 2(\pi^+ \pi^-)$ | $(4.0 \pm 0.6) \times 10^{-4}$ | | |
| Γ_{176} | $\gamma K^{*0} K^+ \pi^- + \text{c.c.}$ | $(3.7 \pm 0.9) \times 10^{-4}$ | | |
| Γ_{177} | $\gamma K^{*0} \bar{K}^{*0}$ | $(2.4 \pm 0.7) \times 10^{-4}$ | | |
| Γ_{178} | $\gamma K_S^0 K^+ \pi^- + \text{c.c.}$ | $(2.6 \pm 0.5) \times 10^{-4}$ | | |
| Γ_{179} | $\gamma K^+ K^- \pi^+ \pi^-$ | $(1.9 \pm 0.5) \times 10^{-4}$ | | |
| Γ_{180} | $\gamma p \bar{p}$ | $(3.9 \pm 0.5) \times 10^{-5}$ | | S=2.0 |
| Γ_{181} | $\gamma f_2(1950) \rightarrow \gamma p \bar{p}$ | $(1.20 \pm 0.22) \times 10^{-5}$ | | |
| Γ_{182} | $\gamma f_2(2150) \rightarrow \gamma p \bar{p}$ | $(7.2 \pm 1.8) \times 10^{-6}$ | | |
| Γ_{183} | $\gamma X(1835) \rightarrow \gamma p \bar{p}$ | $(4.6 \begin{smallmatrix} +1.8 \\ -4.0 \end{smallmatrix}) \times 10^{-6}$ | | |

| | | | | |
|----------------|---|--|------------------|--------|
| Γ_{184} | $\gamma X \rightarrow \gamma p \bar{p}$ | [a] < 2 | $\times 10^{-6}$ | CL=90% |
| Γ_{185} | $\gamma \pi^+ \pi^- p \bar{p}$ | (2.8 \pm 1.4) | $\times 10^{-5}$ | |
| Γ_{186} | $\gamma 2(\pi^+ \pi^-) K^+ K^-$ | < 2.2 | $\times 10^{-4}$ | CL=90% |
| Γ_{187} | $\gamma 3(\pi^+ \pi^-)$ | < 1.7 | $\times 10^{-4}$ | CL=90% |
| Γ_{188} | $\gamma K^+ K^- K^+ K^-$ | < 4 | $\times 10^{-5}$ | CL=90% |
| Γ_{189} | $\gamma \gamma J/\psi$ | (3.1 $\begin{smallmatrix} +1.0 \\ -1.2 \end{smallmatrix}$) | $\times 10^{-4}$ | |
| Γ_{190} | $e^+ e^- \eta'$ | (1.90 \pm 0.26) | $\times 10^{-6}$ | |
| Γ_{191} | $e^+ e^- \chi_{c0}(1P)$ | (1.06 \pm 0.24) | $\times 10^{-3}$ | |
| Γ_{192} | $e^+ e^- \chi_{c1}(1P)$ | (8.5 \pm 0.6) | $\times 10^{-4}$ | |
| Γ_{193} | $e^+ e^- \chi_{c2}(1P)$ | (7.0 \pm 0.8) | $\times 10^{-4}$ | |

Weak decays

| | | | | |
|----------------|-----------------------------|-------|------------------|--------|
| Γ_{194} | $D^0 e^+ e^- + \text{c.c.}$ | < 1.4 | $\times 10^{-7}$ | CL=90% |
|----------------|-----------------------------|-------|------------------|--------|

Other decays

| | | | | |
|----------------|-----------|-------|---|--------|
| Γ_{195} | invisible | < 1.6 | % | CL=90% |
|----------------|-----------|-------|---|--------|

[a] For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 378.1$ for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

| | | | | | | | | | | |
|-----------|-------|-------|-------|----------|----------|----------|----------|-----------|-----------|-----------|
| x_7 | 3 | | | | | | | | | |
| x_8 | 1 | 0 | | | | | | | | |
| x_{11} | 29 | 11 | 2 | | | | | | | |
| x_{12} | 28 | 6 | 1 | 48 | | | | | | |
| x_{13} | 13 | 4 | 1 | 36 | 15 | | | | | |
| x_{21} | 0 | 0 | 0 | 4 | 3 | 2 | | | | |
| x_{147} | 1 | 0 | 0 | 2 | 1 | 1 | 0 | | | |
| x_{148} | 1 | 0 | 0 | 2 | 1 | 1 | 0 | 0 | | |
| x_{149} | 1 | 0 | 0 | 3 | 1 | 1 | 0 | 0 | 0 | |
| Γ | -81 | -4 | -1 | -38 | -34 | -16 | -7 | -1 | -1 | -1 |
| | x_6 | x_7 | x_8 | x_{11} | x_{12} | x_{13} | x_{21} | x_{147} | x_{148} | x_{149} |

$\psi(2S)$ PARTIAL WIDTHS $\Gamma(\text{hadrons})$ Γ_1

| VALUE (keV) | DOCUMENT ID | TECN | COMMENT |
|--------------|-------------|------|---------------|
| 258 ± 26 | BAI | 02B | BES2 e^+e^- |
| 224 ± 56 | LUTH | 75 | MRK1 e^+e^- |

 $\Gamma(e^+e^-)$ Γ_6

| VALUE (keV) | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|------|---|
| 2.33 ± 0.04 OUR FIT | | | |
| 2.29 ± 0.06 OUR AVERAGE | | | |
| $2.23 \pm 0.10 \pm 0.02$ | ¹ ABLIKIM | 15V | BES3 4.0–4.4 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ |
| $2.338 \pm 0.037 \pm 0.096$ | ABLIKIM | 08B | BES2 $e^+e^- \rightarrow \text{hadrons}$ |
| $2.330 \pm 0.036 \pm 0.110$ | ABLIKIM | 06L | BES2 $e^+e^- \rightarrow \text{hadrons}$ |
| 2.44 ± 0.21 | ² BAI | 02B | BES2 e^+e^- |
| 2.14 ± 0.21 | ALEXANDER | 89 | RVUE See Υ mini-review |
| $2.279 \pm 0.015 \pm 0.042$ | ³ ANASHIN | 18 | KEDR e^+e^- |
| $2.282 \pm 0.015 \pm 0.042$ | ⁴ ANASHIN | 18 | KEDR e^+e^- |
| 2.0 ± 0.3 | BRANDELIK | 79C | DASP e^+e^- |
| 2.1 ± 0.3 | ⁵ LUTH | 75 | MRK1 e^+e^- |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ ABLIKIM 15V reports $2.213 \pm 0.018 \pm 0.099$ keV from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.95 \pm 0.45) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channel, assuming $\Gamma_e = \Gamma_\mu = \Gamma_\tau/0.38847$.

³ Combining $\Gamma_{e^+e^-} \cdot B(\mu^+\mu^-)$ from ANASHIN 18 with $\Gamma_{e^+e^-} \cdot B(\text{hadrons})$ from ANASHIN 12 and assuming lepton universality.

⁴ From the sum of $\Gamma_{e^+e^-} \cdot B(\text{hadrons})$ from ANASHIN 12, $\Gamma_{e^+e^-} \cdot B(e^+e^-)$ and $\Gamma_{e^+e^-} \cdot B(\mu^+\mu^-)$ from ANASHIN 18, and $\Gamma_{e^+e^-} \cdot B(\tau^+\tau^-)$ from ANASHIN 07.

⁵ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

 $\Gamma(\gamma\gamma)$ Γ_{165}

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------|-----|-------------|------|---------------|
| <43 | 90 | BRANDELIK | 79C | DASP e^+e^- |

$\psi(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(i) in the e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(i)$ or the branching ratio $\Gamma(i)/\text{total}$.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1\Gamma_6/\Gamma$

| VALUE (keV) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|----------------------|------|--|
| 2.233±0.015±0.042 | ¹ ANASHIN | 12 | KEDR $e^+e^- \rightarrow \text{hadrons}$ |
| 2.2 ± 0.4 | ABRAMS | 75 | MRK1 e^+e^- |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ ANASHIN 12 reports the value $2.233 \pm 0.015 \pm 0.037 \pm 0.020$ keV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

$\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_6\Gamma_6/\Gamma$

| VALUE (eV) | DOCUMENT ID | TECN | COMMENT |
|---------------------|----------------------|------|---------------|
| 21.2±0.7±1.2 | ¹ ANASHIN | 18 | KEDR e^+e^- |

¹ From the average of nine scans of the $\psi(2S)$.

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_7\Gamma_6/\Gamma$

| VALUE (eV) | DOCUMENT ID | TECN | COMMENT |
|---------------------|----------------------|------|--|
| 19.3±0.3±0.5 | ¹ ANASHIN | 18 | KEDR $\psi(2S) \rightarrow \mu^+\mu^-$ |

¹ From the average of nine scans of the $\psi(2S)$.

$\Gamma(\tau^+\tau^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_8\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|----------------------|------|---|
| 9.0±2.6 | 79 | ¹ ANASHIN | 07 | KEDR $e^+e^- \rightarrow \psi(2S) \rightarrow \tau^+\tau^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Using $\psi(2S)$ total width of 337 ± 13 keV. Systematic errors not evaluated.

$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{11}\Gamma_6/\Gamma$


| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------|-------------------|------|---|
| 0.808±0.013 OUR FIT | | | | |
| 0.837±0.025 OUR AVERAGE | | | | Error includes scale factor of 1.3. See the ideogram below. |
| 0.837±0.028±0.005 | | ¹ LEES | 12E | BABR $10.6 e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$ |
| 0.852±0.010±0.026 | 19.5k | ADAM | 06 | CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$ |
| 0.68 ± 0.09 | | ² BAI | 98E | BES e^+e^- |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------|-----|---------------------|------|---|
| 0.88 ± 0.08 ± 0.03 | 256 | ³ AUBERT | 07AU | BABR $10.6 e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$ |
| 0.755±0.048±0.004 | 544 | ⁴ AUBERT | 05D | BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$ |

¹ LEES 12E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (49.9 \pm 1.3 \pm 1.0) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

- ²The value of $\Gamma(e^+e^-)$ quoted in BAI 98E is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6) \times 10^{-2}$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$. Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.
- ³AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0186 \pm 0.0012 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.10 \pm 0.08) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴AUBERT 05D reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = 0.0450 \pm 0.0018 \pm 0.0022$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E.



| $\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{12}\Gamma_6/\Gamma$ |
|---|------|-------------------|------|---|------------------------------|
| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 0.425±0.009 OUR FIT | | | | | |
| 0.411±0.008±0.018 | 3.6k | ADAM | 06 | CLEO 3.773 $e^+e^- \rightarrow \gamma\psi(2S)$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 0.48 ±0.09 ±0.02 | 142 | ¹ LEES | 18E | BABR 10.6 $e^+e^- \rightarrow J/\psi\pi^0\pi^0\gamma$ | |

¹LEES 18E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0101 \pm 0.0015 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.10 \pm 0.08) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(J/\psi(1S)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{13}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|----------|---------------------|-----------|---|
| 78.6 ± 1.6 OUR FIT | | | | |
| 87 ± 9 OUR AVERAGE | | | | |
| 83 ± 25 ± 5 | 14 | ¹ AUBERT | 07AU BABR | 10.6 e ⁺ e ⁻ → J/ψ π ⁺ π ⁻ π ⁰ γ |
| 88 ± 6 ± 7 | 291 ± 24 | ADAM | 06 CLEO | 3.773 e ⁺ e ⁻ → γ ψ(2S) |
| ¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow J/\psi \eta) \cdot B(J/\psi \rightarrow \mu^+ \mu^-) \cdot B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 1.11 \pm 0.33 \pm 0.07$ eV. | | | | |

$\Gamma(J/\psi(1S)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{14}\Gamma_6/\Gamma$

| VALUE (eV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------|-----|------|-------------|---------|---|
| <8 | 90 | <37 | ADAM | 06 CLEO | 3.773 e ⁺ e ⁻ → γ ψ(2S) |

$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{21}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------|----------|---|
| 0.686 ± 0.019 OUR FIT | | | | |
| 0.63 ± 0.05 OUR AVERAGE | | | | Error includes scale factor of 1.2. |
| 0.67 ± 0.12 ± 0.02 | 43 | ¹ LEES | 130 BABR | e ⁺ e ⁻ → p \bar{p} γ |
| 0.74 ± 0.07 ± 0.04 | 142 | ² LEES | 13Y BABR | e ⁺ e ⁻ → p \bar{p} γ |
| 0.579 ± 0.038 ± 0.036 | 2.7k | ANDREOTTI | 07 E835 | p \bar{p} → e ⁺ e ⁻ , J/ψ X |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.70 ± 0.17 ± 0.03 | 22 | ³ AUBERT | 06B BABR | e ⁺ e ⁻ → p \bar{p} γ |
| ¹ ISR photon reconstructed in the detector | | | | |
| ² ISR photon undetected | | | | |
| ³ Superseded by LEES 130 | | | | |

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{29}\Gamma_6/\Gamma$

| VALUE (eV) | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|-----------|--|
| 1.5 ± 0.4 ± 0.1 | AUBERT | 07BD BABR | 10.6 e ⁺ e ⁻ → Λ $\bar{\Lambda}$ γ |

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{66}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------|------|-------------|----------|---|
| 11.2 ± 3.3 ± 1.3 | 43 | AUBERT | 06D BABR | 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻ π ⁰) γ |

$\Gamma(\pi^0\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{77}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|-------------|----------|--|
| 0.60 ± 0.31 ± 0.03 | 17 | LEES | 12F BABR | 10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ |

$\Gamma(K^+K^-2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{84}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------|------|-------------|----------|--|
| 4.4 ± 2.1 ± 0.3 | 26 | AUBERT | 06D BABR | 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ 2(π ⁺ π ⁻) γ |

$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{78}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------|-----------|--|
| 1.92 ± 0.30 ± 0.06 | 133 | LEES | 12F BABR | 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 2.56 ± 0.42 ± 0.16 | 85 | ¹ AUBERT | 07AK BABR | 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ |

¹Superseded by LEES 12F.

$$\Gamma(\pi^0\pi^0 K_S^0 K_L^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{79}\Gamma_6/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|---|
| 2.92±1.27±0.15 | 14 | LEES | 17A BABR | $e^+e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$ |

$$\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{118}\Gamma_6/\Gamma$$

| VALUE (eV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------|-----|------|-------------|----------|---|
| <0.7 | 90 | 8 | LEES | 17A BABR | $e^+e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$ |

$$\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{119}\Gamma_6/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 3.14±1.08±0.16 | 16 | LEES | 17A BABR | $e^+e^- \rightarrow K_S^0 K_L^0 \eta \gamma$ |

$$\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{123}\Gamma_6/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------------|----------|--|
| 0.345±0.128±0.004 | 12 | ¹ LEES | 12F BABR | $10.6 e^+e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------|-------|---------------------|-----------|--|
| 0.345±0.168±0.004 | 6 ± 3 | ² AUBERT | 07AK BABR | $10.6 e^+e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |
|-------------------|-------|---------------------|-----------|--|

¹LEES 12F reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.17 \pm 0.06 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.17 \pm 0.08 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(2(K^+ K^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{124}\Gamma_6/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 0.22±0.10±0.02 | 13 | LEES | 12F BABR | $10.6 e^+e^- \rightarrow K^+ K^- K^+ K^- \gamma$ |

$$\Gamma(\phi \pi^+ \pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{122}\Gamma_6/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------------|----------|--|
| 0.55±0.19±0.01 | 19 | ¹ LEES | 12F BABR | $10.6 e^+e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|----------------|----|-------------------------|----------|--|
| 0.57±0.23±0.01 | 10 | ² AUBERT, BE | 06D BABR | $10.6 e^+e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$ |
|----------------|----|-------------------------|----------|--|

¹LEES 12F reports $[\Gamma(\psi(2S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.27 \pm 0.09 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Superseded by LEES 12F. AUBERT, BE 06D reports $[\Gamma(\psi(2S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.28 \pm 0.11 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{17}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------|------|-------------|-----------|--|
| 29.7±2.2±1.8 | 410 | AUBERT | 07AU BABR | 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)π ⁰ γ |

$\Gamma(\pi^+\pi^-\pi^0\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{19}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------|------|-------------|----------|--|
| 12.4±1.8±1.2 | 177 | LEES | 18E BABR | 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ 3π ⁰ γ |

$\Gamma(\rho^\pm\pi^\mp\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{20}\Gamma_6/\Gamma$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------|-----|-------------|----------|--|
| <6.2 | 90 | LEES | 18E BABR | 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ 3π ⁰ γ |

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{72}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|---------------------|-----------|---|
| 3.01±0.84±0.02 | 37 | ¹ AUBERT | 07AU BABR | 10.6 e ⁺ e ⁻ → ωπ ⁺ π ⁻ γ |

¹AUBERT 07AU reports [$\Gamma(\psi(2S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(ω(782) → π⁺π⁻π⁰)] = 2.69 ± 0.73 ± 0.16 eV which we divide by our best value B(ω(782) → π⁺π⁻π⁰) = (89.3 ± 0.6) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{76}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------------|----------|--|
| 2.58±0.82±0.02 | 33 | ¹ LEES | 18E BABR | 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ 3π ⁰ γ |

¹LEES 18E reports [$\Gamma(\psi(2S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(ω(782) → π⁺π⁻π⁰)] = 2.3 ± 0.7 ± 0.2 eV which we divide by our best value B(ω(782) → π⁺π⁻π⁰) = (89.3 ± 0.6) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+\pi^-\eta)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{69}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|---------------------|-----------|--|
| 2.87±1.41±0.01 | 16 | ¹ AUBERT | 07AU BABR | 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)ηγ |

¹AUBERT 07AU reports [$\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-\eta)) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(η → 2γ)] = 1.13 ± 0.55 ± 0.08 eV which we divide by our best value B(η → 2γ) = (39.41 ± 0.20) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{70}\Gamma_6/\Gamma$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------|-----|-------------|----------|---|
| <0.85 | 90 | LEES | 18E BABR | 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ π ⁰ ηγ |

$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{91}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------|------|-------------|-----------|---|
| 4.4±1.3±0.3 | 32 | AUBERT | 07AU BABR | 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ π ⁰ γ |

$\Gamma(K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ **$\Gamma_{82} \Gamma_6 / \Gamma$**

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|--------------------------|------|--|
| 3.04 ± 1.79 ± 0.02 | 7 | ¹ AUBERT 07AU | BABR | 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$ |

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.2 \pm 0.7 \pm 0.1$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ **$\Gamma_{109} \Gamma_6 / \Gamma$**

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|----|-------------------|-----|---|
| 0.147 ± 0.035 ± 0.005 | 66 | ¹ LEES | 15J | BABR $e^+ e^- \rightarrow K^+ K^- \gamma$ |
| 0.197 ± 0.035 ± 0.005 | 66 | ² LEES | 15J | BABR $e^+ e^- \rightarrow K^+ K^- \gamma$ |
| 0.35 ± 0.14 ± 0.03 | 11 | ³ LEES | 13Q | BABR $e^+ e^- \rightarrow K^+ K^- \gamma$ |

¹ $\sin\phi > 0$.

² $\sin\phi < 0$.

³ Interference with non-resonant $K^+ K^-$ production not taken into account.

$\psi(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons}) / \Gamma_{\text{total}}$ **Γ_1 / Γ**

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.9785 ± 0.0013 OUR AVERAGE

| | | | |
|-----------------|-------------------|-----|----------------|
| 0.9779 ± 0.0015 | ¹ BAI | 02B | BES2 $e^+ e^-$ |
| 0.981 ± 0.003 | ¹ LUTH | 75 | MRK1 $e^+ e^-$ |

¹ Includes cascade decay into $J/\psi(1S)$.

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons}) / \Gamma_{\text{total}}$ **Γ_2 / Γ**

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.0173 ± 0.0014 OUR AVERAGE Error includes scale factor of 1.5.

| | | | |
|-----------------|---------------------|-----|----------------|
| 0.0166 ± 0.0010 | ^{1,2} SETH | 04 | RVUE $e^+ e^-$ |
| 0.0199 ± 0.0019 | ¹ BAI | 02B | BES2 $e^+ e^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------------|-------------------|----|----------------|
| 0.029 ± 0.004 | ¹ LUTH | 75 | MRK1 $e^+ e^-$ |
|---------------|-------------------|----|----------------|

¹ Included in $\Gamma(\text{hadrons}) / \Gamma_{\text{total}}$.

² Using $B(\psi(2S) \rightarrow \ell^+ \ell^-) = (0.73 \pm 0.04)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C.

$\Gamma(g g g) / \Gamma_{\text{total}}$ **Γ_3 / Γ**

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

10.58 ± 1.62 2.9 M ¹ LIBBY 09 CLEO $\psi(2S) \rightarrow \text{hadrons}$

¹ Calculated using $\Gamma(\gamma g g) / \Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09, $B(\psi(2S) \rightarrow X J/\psi)$ relative and absolute branching fractions from MENDEZ 08, $B(\psi(2S) \rightarrow \gamma \eta_c)$ from MITCHELL 09, and $B(\psi(2S) \rightarrow \text{virtual } \gamma \rightarrow \text{hadrons})$, $B(\psi(2S) \rightarrow \gamma \chi_{cJ})$, and $B(\psi(2S) \rightarrow \ell^+ \ell^-)$ from PDG 08. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(\gamma g g) / \Gamma_{\text{total}}$ LIBBY 09 measurement.

$\Gamma(\gamma g g)/\Gamma_{\text{total}}$ Γ_4/Γ

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 1.025 ± 0.288 | 200 k | ¹ LIBBY | 09 | CLEO $\psi(2S) \rightarrow \gamma + \text{hadrons}$ |

¹ Calculated using $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(g g g)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

$\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_4/Γ_3

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 9.7 ± 2.6 ± 1.6 | 2.9 M | LIBBY | 09 | CLEO $\psi(2S) \rightarrow (\gamma +) \text{hadrons}$ |

$\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$ Γ_5/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|---------------------|-------------|-------------------------------------|
| 0.154 ± 0.015 | ¹ MENDEZ | 08 | CLEO $e^+ e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------------|-------------------|-----|-------------------------------------|
| 0.169 ± 0.026 | ² ADAM | 05A | CLEO $e^+ e^- \rightarrow \psi(2S)$ |
|---------------|-------------------|-----|-------------------------------------|

¹ Uses $B(\psi(2S) \rightarrow J/\psi X)$ from MENDEZ 08 and other branching fractions from PDG 07.

² Uses $B(J/\psi X)$ from ADAM 05A, $B(\chi_{cJ} \gamma)$, $B(\eta_c \gamma)$ from ATHAR 04 and $B(\ell^+ \ell^-)$ from PDG 04. Superseded by MENDEZ 08.

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_6/Γ

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| 79.3 ± 1.7 OUR FIT | | | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------|----------------------|----|----------------|
| 88 ± 13 | ¹ FELDMAN | 77 | RVUE $e^+ e^-$ |
|---------|----------------------|----|----------------|

¹ From an overall fit assuming equal partial widths for $e^+ e^-$ and $\mu^+ \mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> |
|---|--------------------|
| 80 ± 6 OUR FIT | |

$\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$ Γ_7/Γ_6

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|--------------------|-------------|----------------|
| 1.00 ± 0.08 OUR FIT | | | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------------|----------|-----|----------------|
| 0.89 ± 0.16 | BOYARSKI | 75C | MRK1 $e^+ e^-$ |
|-------------|----------|-----|----------------|

$\Gamma(\tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_8/Γ

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|----------------------|-------------|------------------------------------|
| 31 ± 4 OUR FIT | | | |
| 30.8 ± 2.1 ± 3.8 | ¹ ABLIKIM | 06W | BES $e^+ e^- \rightarrow \psi(2S)$ |

¹ Computed using PDG 02 value of $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$ to estimate the total number of $\psi(2S)$ events.

————— **DECAYS INTO $J/\psi(1S)$ AND ANYTHING** —————

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$

Γ_9/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------|----------|--------------------------------------|
| 0.614 ± 0.006 | | | | OUR FIT |
| 0.55 ± 0.07 | | | | OUR AVERAGE |
| 0.51 ± 0.12 | | BRANDELIK | 79C DASP | $e^+e^- \rightarrow \mu^+\mu^-X$ |
| 0.57 ± 0.08 | | ABRAMS | 75B MRK1 | $e^+e^- \rightarrow \mu^+\mu^-X$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.6254 ± 0.0016 ± 0.0155 | 1.1M | ¹ MENDEZ | 08 CLEO | $\psi(2S) \rightarrow \ell^+\ell^-X$ |
| 0.5950 ± 0.0015 ± 0.0190 | 151k | ADAM | 05A CLEO | Repl. by MENDEZ 08 |

¹ Not independent from other measurements of MENDEZ 08.


$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_6/\Gamma_9 = \Gamma_6/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.343\Gamma_{148} + 0.190\Gamma_{149})$$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----------|-------------------------|----------|--|
| 1.291 ± 0.026 | | | | OUR FIT |
| 1.28 ± 0.04 | | | | OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below. |
| 1.22 ± 0.02 ± 0.05 | 5097 ± 73 | ¹ ANDREOTTI | 05 E835 | $p\bar{p} \rightarrow \psi(2S) \rightarrow e^+e^-$ |
| 1.28 ± 0.03 ± 0.02 | | ¹ AMBROGIANI | 00A E835 | $p\bar{p} \rightarrow \psi(2S)$ |
| 1.44 ± 0.08 ± 0.02 | | ¹ ARMSTRONG | 97 E760 | $\bar{p}p \rightarrow \psi(2S)$ |

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

WEIGHTED AVERAGE
1.28 ± 0.04 (Error scaled by 1.6)



$\Gamma(\mu^+ \mu^-)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_7/\Gamma_9 = \Gamma_7/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13}+0.343\Gamma_{148}+0.190\Gamma_{149})$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------|------|------------------------------------|
| 0.0130±0.0010 OUR FIT | | | |
| 0.014 ±0.003 | HILGER | 75 | SPEC e ⁺ e ⁻ |

$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma_{\text{total}}$

Γ_{10}/Γ

| VALUE | DOCUMENT ID |
|------------------------------|-------------|
| 0.2538±0.0032 OUR FIT | |

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{11}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|------|---------|
| 0.3468±0.0030 OUR FIT | | | | |

0.348 ±0.005 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.


| | | | | | |
|----------------------|------|---------|-----|------|--|
| 0.3498±0.0002±0.0045 | 20M | ABLIKIM | 13R | BES3 | $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$ |
| 0.3504±0.0007±0.0077 | 565k | MENDEZ | 08 | CLEO | $\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$ |
| 0.323 ±0.014 | | BAI | 02B | BES2 | e ⁺ e ⁻ |
| 0.32 ±0.04 | | ABRAMS | 75B | MRK1 | e ⁺ e ⁻ → J/ψπ ⁺ π ⁻ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|----------------------|-----|-------------------|-----|------|--------------------|
| 0.3354±0.0014±0.0110 | 60k | ¹ ADAM | 05A | CLEO | Repl. by MENDEZ 08 |
|----------------------|-----|-------------------|-----|------|--------------------|

¹Not independent from other values reported by ADAM 05A.

WEIGHTED AVERAGE
0.348±0.005 (Error scaled by 1.3)



$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_6/Γ_{11}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------|---------------------|-------------|----------------|
| 0.0229±0.0005 OUR FIT | | | |
| 0.0252±0.0028±0.0011 | ¹ AUBERT | 02B | BABR e^+e^- |

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

$\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_7/Γ_{11}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|------------------------|-------------|--|
| 0.0230±0.0017 OUR FIT | | | |
| 0.0228±0.0018 OUR AVERAGE | | | |
| 0.0230±0.0020±0.0012 | ¹ AAIJ | 16Y | LHCB $\Lambda_b^0 \rightarrow \psi(2S)X$ |
| 0.0216±0.0026±0.0014 | ² AUBERT | 02B | BABR e^+e^- |
| 0.0327±0.0077±0.0072 | ² GRIBUSHIN | 96 | FMPS 515 $\pi^-Be \rightarrow 2\mu X$ |

¹ Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$.
² Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10) \times 10^{-2}$.

$\Gamma(\tau^+\tau^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_8/Γ_{11}

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| 8.8 ±1.1 OUR FIT | | | |
| 8.73±1.39±1.57 | BAI | 02 | BES e^+e^- |


$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$ Γ_{11}/Γ_9

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|------------------------|-------------|---|
| 0.5645±0.0026 OUR FIT | | | | |
| 0.554 ±0.008 OUR AVERAGE | | | | Error includes scale factor of 1.3. See the ideogram below. |
| 0.5604±0.0009±0.0062 | 565k | MENDEZ | 08 | CLEO $\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$ |
| 0.525 ±0.009 ±0.022 | 4k | ANDREOTTI | 05 | E835 $\psi(2S) \rightarrow J/\psi X$ |
| 0.536 ±0.007 ±0.016 | 20k | ^{1,2} ABLIKIM | 04B | BES $\psi(2S) \rightarrow J/\psi X$ |
| 0.496 ±0.037 | | ARMSTRONG | 97 | E760 $\bar{p}p \rightarrow \psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.5637±0.0027±0.0046 | 60k | ADAM | 05A | CLEO Repl. by MENDEZ 08 |

¹ From a fit to the J/ψ recoil mass spectra.

² ABLIKIM 04B quotes $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)$.

WEIGHTED AVERAGE
 0.554 ± 0.008 (Error scaled by 1.3)



$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

$\Gamma_{10}/\Gamma_{11} = (0.9761\Gamma_{12} + 0.719\Gamma_{13} + 0.343\Gamma_{148} + 0.190\Gamma_{149})/\Gamma_{11}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|--------------|------|----------|
| 0.732 ± 0.008 OUR FIT | | | |
| 0.73 ± 0.09 | TANENBAUM 76 | MRK1 | e^+e^- |

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{total}$

Γ_{12}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-------------|------|---------|
| 0.1824 ± 0.0031 OUR FIT | | | | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------------------|-------|---------------------|----------|---|
| $0.1769 \pm 0.0008 \pm 0.0053$ | 61k | ¹ MENDEZ | 08 CLEO | $\psi(2S) \rightarrow \ell^+\ell^-2\pi^0$ |
| $0.1652 \pm 0.0014 \pm 0.0058$ | 13.4k | ² ADAM | 05A CLEO | Repl. by MENDEZ 08 |

¹ Not independent from other measurements of MENDEZ 08.

² Not independent from other values reported by ADAM 05A.

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{12}/Γ_9

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-------------|------|---------|
| 0.2968 ± 0.0031 OUR FIT | | | | |

0.320 ± 0.012 OUR AVERAGE

| | | | | |
|-----------------------------|---------------|------------|----------|---------------------------------|
| $0.300 \pm 0.008 \pm 0.022$ | 1655 ± 44 | ANDREOTTI | 05 E835 | $\psi(2S) \rightarrow J/\psi X$ |
| $0.328 \pm 0.013 \pm 0.008$ | | AMBROGIANI | 00A E835 | $p\bar{p} \rightarrow \psi(2S)$ |
| 0.323 ± 0.033 | | ARMSTRONG | 97 E760 | $p\bar{p} \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------------------|-------|--------|----------|---|
| $0.2829 \pm 0.0012 \pm 0.0056$ | 61k | MENDEZ | 08 CLEO | $\psi(2S) \rightarrow \ell^+\ell^-2\pi^0$ |
| $0.2776 \pm 0.0025 \pm 0.0043$ | 13.4k | ADAM | 05A CLEO | Repl. by MENDEZ 08 |

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-) \qquad \Gamma_{12}/\Gamma_{11}$$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|------------------------|-------------|---|
| 0.526 ± 0.008 OUR FIT | | | | |
| 0.513 ± 0.022 OUR AVERAGE | | | | Error includes scale factor of 2.2. |
| 0.5047 ± 0.0022 ± 0.0102 | 61k | MENDEZ | 08 CLEO | $\psi(2S) \rightarrow \ell^+\ell^-2\pi^0$ |
| 0.570 ± 0.009 ± 0.026 | 14k | ¹ ABLIKIM | 04B BES | $\psi(2S) \rightarrow J/\psi X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.4924 ± 0.0047 ± 0.0086 | 73k | ^{2,3} ADAM | 05A CLEO | Repl. by MENDEZ 08 |
| 0.571 ± 0.018 ± 0.044 | | ⁴ ANDREOTTI | 05 E835 | $\psi(2S) \rightarrow J/\psi X$ |
| 0.53 ± 0.06 | | TANENBAUM | 76 MRK1 | e^+e^- |
| 0.64 ± 0.15 | | ⁵ HILGER | 75 SPEC | e^+e^- |

¹ From a fit to the J/ψ recoil mass spectra.

² Not independent from other values reported by ADAM 05A.

³ Using 13,217 $J/\psi\pi^0\pi^0$ and 60,010 $J/\psi\pi^+\pi^-$ events.

⁴ Not independent from other values reported by ANDREOTTI 05.

⁵ Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.

$$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}} \qquad \Gamma_{13}/\Gamma$$

| <u>VALUE (units 10⁻³)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|------------------------|-------------|---|
| 33.7 ± 0.5 OUR FIT | | | | |
| 32.9 ± 1.7 OUR AVERAGE | | | | Error includes scale factor of 2.1. See the ideogram below. |
| 33.75 ± 0.17 ± 0.86 | 68.2k | ABLIKIM | 12M BES3 | $e^+e^- \rightarrow \ell^+\ell^-2\gamma$ |
| 29.8 ± 0.9 ± 2.3 | 5.7k | BAI | 04I BES2 | $\psi(2S) \rightarrow J/\psi\gamma\gamma$ |
| 25.5 ± 2.9 | 386 | ¹ OREGLIA | 80 CBAL | $e^+e^- \rightarrow J/\psi2\gamma$ |
| 45 ± 12 | 17 | ² BRANDELIK | 79B DASP | $e^+e^- \rightarrow J/\psi2\gamma$ |
| 42 ± 6 | 164 | ² BARTEL | 78B CNTR | e^+e^- |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 34.3 ± 0.4 ± 0.9 | 18.4k | ³ MENDEZ | 08 CLEO | $\psi(2S) \rightarrow \ell^+\ell^-\eta$ |
| 32.5 ± 0.6 ± 1.1 | 2.8k | ⁴ ADAM | 05A CLEO | Repl. by MENDEZ 08 |
| 43 ± 8 | 44 | TANENBAUM | 76 MRK1 | e^+e^- |


¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

³ Not independent from other measurements of MENDEZ 08.

⁴ Not independent from other values reported by ADAM 05A.

WEIGHTED AVERAGE
 32.9 ± 1.7 (Error scaled by 2.1)




$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{13}/Γ_9

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|----------|------------------------|------|--|
| 0.0549 ± 0.0008 | | | | OUR FIT |
| 0.058 ± 0.007 | | | | OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below. |
| 0.050 ± 0.006 ± 0.003 | 298 ± 20 | ANDREOTTI 05 | E835 | $\psi(2S) \rightarrow J/\psi X$ |
| 0.072 ± 0.009 | | AMBROGIANI 00A | E835 | $p\bar{p} \rightarrow \psi(2S)$ |
| 0.061 ± 0.015 | | ARMSTRONG 97 | E760 | $\bar{p}p \rightarrow \psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.0549 ± 0.0006 ± 0.0009 | 18.4k | ¹ MENDEZ 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- \eta$ |
| 0.0546 ± 0.0010 ± 0.0007 | 2.8k | ADAM 05A | CLEO | Repl. by MENDEZ 08 |

¹Not independent from other measurements of MENDEZ 08.



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_{13}/Γ_{11}

| VALUE | EVTs | DOCUMENT ID | TECN | COMMENT |
|---|-------|---------------------------|------|---|
| 0.0972 ± 0.0014 | | | | OUR FIT |
| 0.0979 ± 0.0018 | | | | OUR AVERAGE |
| $0.0979 \pm 0.0010 \pm 0.0015$ | 18.4k | MENDEZ 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- \eta$ |
| $0.098 \pm 0.005 \pm 0.010$ | 2k | ¹ ABLIKIM 04B | BES | $\psi(2S) \rightarrow J/\psi X$ |
| 0.091 ± 0.021 | | ² HIMEL 80 | MRK2 | $e^+ e^- \rightarrow \psi(2S) X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| $0.0968 \pm 0.0019 \pm 0.0013$ | 2.8k | ³ ADAM 05A | CLEO | Repl. by MENDEZ 08 |
| $0.095 \pm 0.007 \pm 0.007$ | | ⁴ ANDREOTTI 05 | E835 | $\psi(2S) \rightarrow J/\psi X$ |

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow J/\psi(1S)\eta)$ reported in HIMEL 80 is derived using $B(\psi(2S)) \rightarrow J/\psi(1S)\pi^+\pi^- = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$.

³ Not independent from other values reported by ADAM 05A.

⁴ Not independent from other values reported by ANDREOTTI 05.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$

Γ_{14}/Γ

| VALUE (units 10^{-4}) | EVTs | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------|------|--|
| 12.68 ± 0.32 | | | | OUR AVERAGE |
| $12.6 \pm 0.2 \pm 0.3$ | 4.1k | ABLIKIM 12M | BES3 | $e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$ |
| $13.3 \pm 0.8 \pm 0.3$ | 530 | MENDEZ 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- 2\gamma$ |
| $14.3 \pm 1.4 \pm 1.2$ | 280 | BAI 04I | BES2 | $\psi(2S) \rightarrow J/\psi \gamma \gamma$ |
| 14 ± 6 | 7 | HIMEL 80 | MRK2 | $e^+ e^-$ |
| $9 \pm 2 \pm 1$ | 23 | ¹ OREGLIA 80 | CBAL | $\psi(2S) \rightarrow J/\psi 2\gamma$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |

13 ± 1 ± 1 88 ADAM 05A CLEO Repl. by MENDEZ 08

¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_{14}/\Gamma_9 = \Gamma_{14}/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.343\Gamma_{148} + 0.190\Gamma_{149})$$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------------|-----|---------------------|-----|--|
| $0.213 \pm 0.012 \pm 0.003$ | 527 | ¹ MENDEZ | 08 | CLEO $e^+ e^- \rightarrow J/\psi \gamma \gamma$ |
| $0.22 \pm 0.02 \pm 0.01$ | | ² ADAM | 05A | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma \gamma$ |

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+ \pi^-)$

Γ_{14}/Γ_{11}

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------------|-----|---------------------|-----|--|
| $0.380 \pm 0.022 \pm 0.005$ | 527 | ¹ MENDEZ | 08 | CLEO $e^+ e^- \rightarrow J/\psi \gamma \gamma$ |
| $0.39 \pm 0.04 \pm 0.01$ | | ² ADAM | 05A | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma \gamma$ |

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

———— HADRONIC DECAYS ————

$\Gamma(\pi^0 h_c(1P))/\Gamma_{\text{total}}$

Γ_{15}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

8.6 ± 1.3 OUR AVERAGE

| | | | | |
|-----------------------|-----|-----------------|-----|--|
| $9.0 \pm 1.5 \pm 1.3$ | 3k | ¹ GE | 11 | CLEO $\psi(2S) \rightarrow \pi^0 \text{ anything}$ |
| $8.4 \pm 1.3 \pm 1.0$ | 11k | ABLIKIM | 10B | BES3 $\psi(2S) \rightarrow \pi^0 h_c$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|------------------|--------|-----|--|
| seen | 92^{+23}_{-22} | ADAMS | 09 | CLEO $\psi(2S) \rightarrow 2\pi^+ 2\pi^- 2\pi^0$ |
| seen | 1282 | DOBBS | 08A | CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$ |
| seen | 168 ± 40 | ROSNER | 05 | CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$ |

¹ Assuming a width $\Gamma(h_c(1P)) = 0.86 \text{ MeV} \equiv \Gamma_0$, a measured dependence of the central value of $B = (7.6 + 1.4 \times \Gamma(h_c(1P)/\Gamma_0) \times 10^{-4}$, and with a systematic error that accounts for the width variation range 0.43–1.29 MeV.

$\Gamma(3(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{16}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

| | | | | |
|----------------|---|----------|----|---|
| 35 ± 16 | 6 | FRANKLIN | 83 | MRK2 $e^+ e^- \rightarrow \text{hadrons}$ |
|----------------|---|----------|----|---|

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$


Γ_{17}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|--------------------------|------|--|
| 29 ± 10 OUR AVERAGE | | | | Error includes scale factor of 4.7. See the ideogram below. |
| 24.9 ± 0.7 ± 3.6 | 2173 | ABLIKIM 07D | BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| 127 ± 12 ± 2 | 410 | ¹ AUBERT 07AU | BABR | 10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0\gamma$ |
| 26.1 ± 0.7 ± 3.0 | 1703 | BRIERE 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |

30 ± 8 42 FRANKLIN 83 MRK2 e^+e^-

¹AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (297 \pm 22 \pm 18) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

WEIGHTED AVERAGE
29±10 (Error scaled by 4.7)



| | | | χ^2 |
|---|----------|-----------|----------|
| + | ABLIKIM | 07D BES2 | 1.4 |
| + | AUBERT | 07AU BABR | 63.2 |
| + | BRIERE | 05 CLEO | 1.0 |
| + | FRANKLIN | 83 MRK2 | 0.0 |
| | | | 65.7 |

(Confidence Level < 0.0001)

$\Gamma(\rho a_2(1320))/\Gamma_{\text{total}}$

Γ_{18}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-----|----------|-------------|------|---|
| 2.55 ± 0.73 ± 0.47 | | 112 ± 31 | BAI 04C | BES2 | $\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |
| <2.3 | 90 | | BAI 98J | BES | e^+e^- |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\rho\bar{p})/\Gamma_{\text{total}}$

Γ_{21}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-----------------------|------|---|
| 2.94 ± 0.08 OUR FIT | | | | |
| 3.02 ± 0.08 OUR AVERAGE | | | | |
| 3.05 ± 0.02 ± 0.12 | 19k | ABLIKIM 18T | BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}$ |
| 3.08 ± 0.05 ± 0.18 | 4.5k | ¹ DOBBS 14 | | $e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}$ |

| | | | | | |
|--------------------------|------|-----------|-----|------|--|
| $3.36 \pm 0.09 \pm 0.25$ | 1.6k | ABLIKIM | 07C | BES | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| $2.87 \pm 0.12 \pm 0.15$ | 557 | PEDLAR | 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 1.4 ± 0.8 | 4 | BRANDELIK | 79C | DASP | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 2.3 ± 0.7 | | FELDMAN | 77 | MRK1 | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(p\bar{p})/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{21}/Γ_{11}

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|--|
| 8.49 ± 0.23 | | | | OUR FIT |
| $6.98 \pm 0.49 \pm 0.97$ | | BAI | 01 | BES $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |

$\Gamma(n\bar{n})/\Gamma_{\text{total}}$ Γ_{22}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| $3.06 \pm 0.06 \pm 0.14$ | 6k | ABLIKIM | 18T | BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow n\bar{n}$ |

$\Gamma(\Delta^{++}\bar{\Delta}^{--})/\Gamma_{\text{total}}$ Γ_{23}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| $12.8 \pm 1.0 \pm 3.4$ | 157 | ¹ BAI | 01 | BES $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{24}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|----------------------|-------------|--|
| < 0.29 | 90 | ¹ ABLIKIM | 13F | BES3 $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|----|----------------------|-----|------------------------------------|
| < 12 | 90 | ² ABLIKIM | 07H | BES2 $e^+e^- \rightarrow \psi(2S)$ |
|--------|----|----------------------|-----|------------------------------------|

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{25}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|-------------|----------------------|-------------|--|
| $2.48 \pm 0.34 \pm 0.19$ | | 60 | ¹ ABLIKIM | 13F | BES3 $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---------|----|----------------------|-----|------------------------------------|
| < 4.9 | 90 | ² ABLIKIM | 07H | BES2 $e^+e^- \rightarrow \psi(2S)$ |
|---------|----|----------------------|-----|------------------------------------|

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

$\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$ Γ_{26}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| $1.0 \pm 0.1 \pm 0.1$ | 74.0 | BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ $p\bar{p}K^+\pi^-$ |

$\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{27}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| $1.8 \pm 0.3 \pm 0.3$ | 45.8 | BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ $p\bar{p}K^+\pi^+\pi^-\pi^-$ |

| $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ | | | | | Γ_{28}/Γ |
|--|------|-------------|------|--|----------------------|
| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $2.8 \pm 0.4 \pm 0.5$ | 73.4 | BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}2(\pi^+\pi^-)$ | |


| $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ | | | | | Γ_{29}/Γ |
|---|------|------|---|------|---|
| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| 3.81 ± 0.13 OUR AVERAGE | | | Error includes scale factor of 1.4. See the ideogram below. | | |
| $3.97 \pm 0.02 \pm 0.12$ | 31k | | ABLIKIM | 17L | BES3 $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ |
| $3.71 \pm 0.05 \pm 0.15$ | 6.5k | | ¹ DOBBS | 17 | $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ |
| $3.39 \pm 0.20 \pm 0.32$ | 337 | | ABLIKIM | 07C | BES $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
| $6.4 \pm 1.8 \pm 0.1$ | | | ² AUBERT | 07BD | BABR $10.6 e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$ |
| $3.28 \pm 0.23 \pm 0.25$ | 208 | | PEDLAR | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| $3.75 \pm 0.09 \pm 0.23$ | 1.9k | | ^{1,3} DOBBS | 14 | $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ |
| $1.81 \pm 0.20 \pm 0.27$ | 80 | | ⁴ BAI | 01 | BES $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
| < 4 | 90 | | FELDMAN | 77 | MRK1 $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² AUBERT 07BD reports $[\Gamma(\psi(2S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (15 \pm 4 \pm 1) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by DOBBS 17.

⁴ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.



$\Gamma(\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{30}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|---------------------------------------|
| 1.40±0.03±0.13 | 2.8k | ABLIKIM | 13W BES3 | $\psi(2S) \rightarrow \text{hadrons}$ |

$\Gamma(\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{31}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|---------------------------------------|
| 1.54±0.04±0.13 | 2.8k | ABLIKIM | 13W BES3 | $\psi(2S) \rightarrow \text{hadrons}$ |

$\Gamma(\Lambda\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{32}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|--------------------|------|--|
| 1.23±0.23±0.08 | 30 | ¹ DOBBS | 17 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\Sigma^0\bar{p}K^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{33}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|----------------------|----------|--|
| 1.67±0.13±0.12 | 276 | ¹ ABLIKIM | 13D BES3 | $\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$ |

¹ Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$, and $B(\Sigma^0 \rightarrow \Lambda\gamma) = 100\%$.

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{34}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|------|---------|
| 2.32±0.12 OUR AVERAGE | | | | |

2.31±0.06±0.10 1.9k ¹ DOBBS 17 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

2.57±0.44±0.68 35 PEDLAR 05 CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.51±0.15±0.16 281 ^{1,2} DOBBS 14 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by DOBBS 17.

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{35}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|------|-------------------------------------|
| 2.35±0.09 OUR AVERAGE | | | | Error includes scale factor of 1.1. |

2.44±0.03±0.11 7k ABLIKIM 17L BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

2.22±0.05±0.11 2.6k ¹ DOBBS 17 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

2.35±0.36±0.32 59 ABLIKIM 07C BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

2.63±0.35±0.21 58 PEDLAR 05 CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.25±0.11±0.16 439 ^{1,2} DOBBS 14 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

1.2 ±0.4 ±0.4 8 ³ BAI 01 BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by DOBBS 17.

³ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Sigma(1385)^+\bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ Γ_{36}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|---------|
| 8.5±0.7 OUR AVERAGE | | | | |

8.4±0.5±0.5 1.5k ABLIKIM 16L BES3 $\psi(2S) \rightarrow \Sigma(1385)^+\bar{\Sigma}(1385)^-$

11 ±3 ±3 14 ¹ BAI 01 BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Sigma(1385)^-\bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ Γ_{37}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $8.5 \pm 0.6 \pm 0.6$ | 1.4K | ABLIKIM 16L | BES3 | $\psi(2S) \rightarrow \Sigma(1385)^-\bar{\Sigma}(1385)^+$ |

$\Gamma(\Sigma(1385)^0\bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{38}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|--|
| $0.69 \pm 0.05 \pm 0.05$ | 2.2k | ABLIKIM 17E | BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |

$\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{39}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|-----|------|--|------|---|
| 2.87 ± 0.11 | | | OUR AVERAGE Error includes scale factor of 1.1. | | |
| $3.03 \pm 0.05 \pm 0.14$ | | 3.6k | ¹ DOBBS 17 | | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
| $2.78 \pm 0.05 \pm 0.14$ | | 5k | ABLIKIM 16L | BES3 | $\psi(2S) \rightarrow \Xi^-\bar{\Xi}^+$ |
| $3.03 \pm 0.40 \pm 0.32$ | | 67 | ABLIKIM 07C | BES | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
| $2.38 \pm 0.30 \pm 0.21$ | | 63 | PEDLAR 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------------------|--|-----|-------------------------|------|---|
| $2.66 \pm 0.12 \pm 0.20$ | | 548 | ^{1,2} DOBBS 14 | | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
| $0.94 \pm 0.27 \pm 0.15$ | | 12 | ³ BAI 01 | BES | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
| <2 | | 90 | FELDMAN 77 | MRK1 | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by DOBBS 17.

³ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{40}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|------|-----------------------|--|---|
| 2.3 ± 0.4 | | | OUR AVERAGE Error includes scale factor of 4.2. | |
| $2.73 \pm 0.03 \pm 0.13$ | 11k | ABLIKIM 17E | BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
| $1.97 \pm 0.06 \pm 0.11$ | 1.2k | ¹ DOBBS 17 | | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
| $2.75 \pm 0.64 \pm 0.61$ | 19 | PEDLAR 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------------|-----|-------------------------|--|---|
| $2.02 \pm 0.19 \pm 0.15$ | 112 | ^{1,2} DOBBS 14 | | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
|--------------------------|-----|-------------------------|--|---|

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by DOBBS 17.

$\Gamma(\Xi(1530)^0\bar{\Xi}(1530)^0)/\Gamma_{\text{total}}$ Γ_{41}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|--------------------------|------|---------------------------------------|
| $5.2 \pm 0.3^{+3.2}_{-1.2}$ | | 527 | ¹ ABLIKIM 13S | BES3 | $\psi(2S) \rightarrow \eta p \bar{p}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----|----|-----------|------|--|
| <32 | 90 | PEDLAR 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
|-----|----|-----------|------|--|

| | | | | |
|-------|----|---------------------|-----|--|
| < 8.1 | 90 | ² BAI 01 | BES | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |
|-------|----|---------------------|-----|--|

¹ With $N(1535)$ decaying to $p\eta$.

² Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{42}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|-------------|------|--|
| 3.86 ± 0.27 ± 0.32 | 236 | ABLIKIM 15I | BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \Xi^+ + \text{c.c.}$ |

$\Gamma(\Xi(1690)^- \Xi^+ \rightarrow K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{43}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|-------------|------|--|
| 5.21 ± 1.48 ± 0.57 | 74 | ABLIKIM 15I | BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \Xi^+ + \text{c.c.}$ |

$\Gamma(\Xi(1820)^- \Xi^+ \rightarrow K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{44}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|--|
| 12.03 ± 2.94 ± 1.22 | 136 | ABLIKIM 15I | BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \Xi^+ + \text{c.c.}$ |

$\Gamma(K^- \Sigma^0 \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{45}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|-------------|------|---|
| 3.67 ± 0.33 ± 0.28 | 142 | ABLIKIM 15I | BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Sigma^0 \Xi^+ + \text{c.c.}$ |

$\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{46}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-----|------|-----------------------|------|---|
| 0.52 ± 0.03 ± 0.03 | | 326 | ¹ DOBBS 17 | | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------------|----|----|-------------------------|------|---|
| 0.47 ± 0.09 ± 0.05 | | 27 | ^{1,2} DOBBS 14 | | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| <1.5 | 90 | | ABLIKIM 12Q | BES2 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| <1.6 | 90 | | PEDLAR 05 | CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| <0.73 | 90 | | ³ BAI 01 | BES | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by DOBBS 17.

³ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

$\Gamma(\pi^0 \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{47}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|--------------------------|------|--|
| 1.53 ± 0.07 OUR AVERAGE | | | | |
| 1.65 ± 0.03 ± 0.15 | 4.5k | ABLIKIM 13A | BES3 | $\psi(2S) \rightarrow \rho \bar{\rho} \pi^0$ |
| 1.54 ± 0.06 ± 0.06 | 948 | ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \pi^0 \rho \bar{\rho}$ |
| 1.32 ± 0.10 ± 0.15 | 256 | ¹ ABLIKIM 05E | BES2 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{\rho} \gamma \gamma$ |
| 1.4 ± 0.5 | 9 | FRANKLIN 83 | MRK2 | $e^+ e^-$ |

¹ Computed using $B(\pi^0 \rightarrow \gamma \gamma) = (98.80 \pm 0.03)\%$.

$\Gamma(N(940) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{48}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|--------------------------|------|--|
| 6.42 ± 0.20 ^{+1.78}/_{-1.28} | 1.9k | ¹ ABLIKIM 13A | BES3 | $\psi(2S) \rightarrow \rho \bar{\rho} \pi^0$ |

¹ From a fit of $\pi^0 \rho \bar{\rho}$ data to eight distinct intermediate $N \bar{p}$ resonant states.

$\Gamma(N(1440)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p})/\Gamma_{\text{total}}$ Γ_{49}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

7.3 $^{+1.7}_{-1.5}$ OUR AVERAGE Error includes scale factor of 2.5.

| | | | | |
|------------------------------------|------|------------------------|-----|--|
| 3.58 ± 0.25 $^{+1.59}_{-0.84}$ | 1.1k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho \bar{p} \pi^0$ |
| 8.1 ± 0.7 ± 0.3 | 474 | ² ALEXANDER | 10 | CLEO $\psi(2S) \rightarrow \pi^0 \rho \bar{p}$ |

¹ From a fit of $\pi^0 \rho \bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

² From a fit of the $\rho \bar{p}$ and $\rho \pi^0$ mass distributions to a combination of $N(1440)\bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances.

$\Gamma(N(1520)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p})/\Gamma_{\text{total}}$ Γ_{50}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

| | | | | |
|---|------|----------------------|-----|--|
| 0.64 ± 0.05 $^{+0.22}_{-0.17}$ | 0.2k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho \bar{p} \pi^0$ |
|---|------|----------------------|-----|--|

¹ From a fit of $\pi^0 \rho \bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

$\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p})/\Gamma_{\text{total}}$ Γ_{51}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

| | | | | |
|---|------|----------------------|-----|--|
| 2.47 ± 0.28 $^{+0.99}_{-0.97}$ | 0.7k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho \bar{p} \pi^0$ |
|---|------|----------------------|-----|--|

¹ From a fit of $\pi^0 \rho \bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

$\Gamma(N(1650)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p})/\Gamma_{\text{total}}$ Γ_{52}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

| | | | | |
|---|------|----------------------|-----|--|
| 3.76 ± 0.28 $^{+1.37}_{-1.66}$ | 1.1k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho \bar{p} \pi^0$ |
|---|------|----------------------|-----|--|

¹ From a fit of $\pi^0 \rho \bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

$\Gamma(N(1720)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p})/\Gamma_{\text{total}}$ Γ_{53}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

| | | | | |
|---|------|----------------------|-----|--|
| 1.79 ± 0.10 $^{+0.24}_{-0.71}$ | 0.5k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho \bar{p} \pi^0$ |
|---|------|----------------------|-----|--|

¹ From a fit of $\pi^0 \rho \bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

$\Gamma(N(2300)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p})/\Gamma_{\text{total}}$ Γ_{54}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

| | | | | |
|---|------|----------------------|-----|--|
| 2.62 ± 0.28 $^{+1.12}_{-0.64}$ | 0.9k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho \bar{p} \pi^0$ |
|---|------|----------------------|-----|--|

¹ From a fit of $\pi^0 \rho \bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

$\Gamma(N(2570)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p})/\Gamma_{\text{total}}$ Γ_{55}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

| | | | | |
|---|------|----------------------|-----|--|
| 2.13 ± 0.08 $^{+0.40}_{-0.30}$ | 0.8k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho \bar{p} \pi^0$ |
|---|------|----------------------|-----|--|

¹ From a fit of $\pi^0 \rho \bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

$\Gamma(\pi^0 f_0(2100) \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{56}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------------|------|---------------------------------------|
| 1.1±0.4±0.1 | 76 | ¹ ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \pi^0 p\bar{p}$ |
| ¹ From a fit of the $p\bar{p}$ and $p\pi^0$ mass distributions to a combination of $N_1^*(1440)\bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances. | | | | |

$\Gamma(\eta p\bar{p})/\Gamma_{\text{total}}$ Γ_{57}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------------|---------------------------|------|---|
| 6.0±0.4 OUR AVERAGE | | | | |
| 6.4±0.2±0.6 | 679 | ¹ ABLIKIM 13S | BES3 | $\psi(2S) \rightarrow \eta p\bar{p}$ |
| 5.6±0.6±0.3 | 154 | ¹ ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \eta p\bar{p}$ |
| 5.8±1.1±0.7 | 44.8 ± 8.5 | ² ABLIKIM 05E | BES2 | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$ |
| 8 ±3 ±3 | 9.8 | BRIERE 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ |

¹ With $N(1535)$ decaying to $p\eta$.
² Computed using $B(\eta \rightarrow \gamma\gamma) = (39.43 \pm 0.26)\%$.

$\Gamma(\eta f_0(2100) \rightarrow \eta p\bar{p})/\Gamma_{\text{total}}$ Γ_{58}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------------|------|--------------------------------------|
| 1.2±0.4±0.1 | 31 | ¹ ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \eta p\bar{p}$ |
| ¹ From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$. | | | | |

$\Gamma(N(1535)\bar{p} \rightarrow \eta p\bar{p})/\Gamma_{\text{total}}$ Γ_{59}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------------|------|--------------------------------------|
| 4.4±0.6±0.3 | 123 | ¹ ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \eta p\bar{p}$ |
| ¹ From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$. | | | | |

$\Gamma(\omega p\bar{p})/\Gamma_{\text{total}}$ Γ_{60}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------------|----------------------|------|---|
| 0.69±0.21 OUR AVERAGE | | | | |
| 0.6 ±0.2 ±0.2 | 21.2 | BRIERE 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ |
| 0.8 ±0.3 ±0.1 | 14.9 ± 0.1 | ¹ BAI 03B | BES | $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ |
| ¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$. | | | | |

$\Gamma(\eta' p\bar{p})/\Gamma_{\text{total}}$ Γ_{61}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|--------------------------|------|---------------------------------------|
| 1.10±0.10±0.08 | 491 | ¹ ABLIKIM 19N | BES3 | $\psi(2S) \rightarrow \eta' p\bar{p}$ |
| ¹ From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels. | | | | |

$\Gamma(\phi p\bar{p})/\Gamma_{\text{total}}$ Γ_{62}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|--|
| <0.24 | 90 | BRIERE 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •
 <0.26 90 ¹ BAI 03B BES $\psi(2S) \rightarrow K^+K^-p\bar{p}$
¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(\pi^+\pi^-\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{63}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------|------------------------|------|---|
| 6.0±0.4 OUR AVERAGE | | | | |
| 5.9±0.2±0.4 | 904.5 | BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}\pi^+\pi^-$ |
| 8 ±2 | | ¹ TANENBAUM | 78 | MRK1 e^+e^- |

¹ Assuming entirely strong decay.

$\Gamma(\rho\bar{\rho}\pi^-\text{ or c.c.})/\Gamma_{\text{total}}$ Γ_{64}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|------|--|
| 2.48±0.17 OUR AVERAGE | | | | |
| 2.45±0.11±0.21 | 851 | ABLIKIM | 06i | BES2 $e^+e^- \rightarrow \rho\pi^-X$ |
| 2.52±0.12±0.22 | 849 | ABLIKIM | 06i | BES2 $e^+e^- \rightarrow \bar{\rho}\pi^+X$ |

$\Gamma(\rho\bar{\rho}\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{65}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|----------|-------------|------|---|
| 3.18±0.50±0.50 | 135 ± 21 | ABLIKIM | 06i | BES2 $e^+e^- \rightarrow \rho\pi^-\pi^0X$ |

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{67}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---|
| <1.6 | 90 | BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |

$\Gamma(\eta\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{68}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------|---------------------|------|---|
| 9.5±0.7±1.5 | | ¹ BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 10.3±0.8±1.4 | 201.7 | ² BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow \gamma\gamma)$ |
| 8.1±1.4±1.6 | 50.0 | ² BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow 3\pi)$ |

¹ Average of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi$.

² Not independent from other values reported by BRIERE 05.

$\Gamma(2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$ Γ_{69}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|---------------------|-----------|---|
| 1.2±0.6±0.1 | 16 | ¹ AUBERT | 07AU BABR | 10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$ |

¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+\pi^-)\eta) \cdot B(\eta \rightarrow \gamma\gamma) = 1.2 \pm 0.7 \pm 0.1 \text{ eV}$.

$\Gamma(\eta'\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{71}/Γ


| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|--|
| 4.5±1.6±1.3 | 12.8 | BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$ |

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{72}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|----------|---------------------|-----------|--|
| 7.3±1.2 OUR AVERAGE | | | | Error includes scale factor of 2.1. See the ideogram below. |
| 8.4±0.5±1.2 | 386 | ABLIKIM | 07D BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| 12.2±2.2±0.7 | 37 | ¹ AUBERT | 07AU BABR | 10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$ |
| 8.2±0.5±0.7 | 391 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |
| 4.8±0.6±0.7 | 100 ± 22 | ² BAI | 03B BES | $\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |
| ¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16$ eV. | | | | |
| ² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$. | | | | |

WEIGHTED AVERAGE
7.3±1.2 (Error scaled by 2.1)



$\Gamma(b_1^\pm\pi^\mp)/\Gamma_{\text{total}}$

Γ_{73}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|---------|--------------------|----------|---|
| 4.0 ± 0.6 OUR AVERAGE | | | | Error includes scale factor of 1.1. |
| 5.1 ± 0.6 ± 0.8 | 202 | ABLIKIM | 07D BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| 4.18 ^{+0.43} _{-0.42} ± 0.92 | 170 | ADAM | 05 CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| 3.2 ± 0.6 ± 0.5 | 61 ± 11 | ^{1,2} BAI | 03B BES | $\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 5.2 ± 0.8 ± 1.0 | | ¹ BAI | 99C BES | Repl. by BAI 03B |
| ¹ Assuming $B(b_1 \rightarrow \omega\pi) = 1$. | | | | |
| ² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$. | | | | |

| $\Gamma(b_1^0 \pi^0)/\Gamma_{\text{total}}$ | | | | | Γ_{74}/Γ |
|---|------|-------------|------|---------|--------------------------------|
| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $2.35^{+0.47}_{-0.42} \pm 0.40$ | 45 | ADAM | 05 | CLEO | $e^+ e^- \rightarrow \psi(2S)$ |

| $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ | | | | | Γ_{75}/Γ |
|---|--------------------|-------------|------------------|------|--|
| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| 2.2 ± 0.4 | OUR AVERAGE | | | | |
| 2.3 ± 0.5 ± 0.4 | | 57 | ABLIKIM | 07D | BES2 $e^+ e^- \rightarrow \psi(2S)$ |
| 2.05 $\pm 0.41 \pm 0.38$ | | 62 ± 12 | BAI | 04C | BES2 $\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <1.5 | | 90 | ¹ BAI | 03B | BES $\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ |
| <1.7 | | 90 | BAI | 98J | BES Repl. by BAI 03B |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

| $\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$ | | | | | Γ_{78}/Γ |
|---|--------------------|------------------------|------|---------|--|
| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 7.3 ± 0.5 | OUR AVERAGE | | | | |
| 8.1 $\pm 1.3 \pm 0.3$ | 133 | LEES | 12F | BABR | 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |
| 7.1 $\pm 0.3 \pm 0.4$ | 817.2 | BRIERE | 05 | CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| 16 ± 4 | | ¹ TANENBAUM | 78 | MRK1 | $e^+ e^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 11.0 $\pm 1.9 \pm 0.2$ | 85 | ² AUBERT | 07AK | BABR | 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |

¹ Assuming entirely strong decay.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(\rho^0 K^+ K^-)/\Gamma_{\text{total}}$ | | | | | Γ_{80}/Γ |
|--|-------|-------------|------|---------|--|
| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $2.2 \pm 0.2 \pm 0.4$ | 223.8 | BRIERE | 05 | CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |

| $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$ | | | | | Γ_{81}/Γ |
|---|-----|-------------|-------------|------|--|
| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| $1.86 \pm 0.32 \pm 0.43$ | | 93 ± 16 | BAI | 04C | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <1.2 | | 90 | BAI | 98J | BES $e^+ e^-$ |

| $\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$ | | | | | Γ_{82}/Γ |
|--|------|---------------------|------|---------|--|
| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $1.3 \pm 0.7 \pm 0.1$ | 7 | ¹ AUBERT | 07AU | BABR | 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$ |
| ¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^-) \eta) \cdot B(\eta \rightarrow \gamma \gamma) = 1.2 \pm 0.7 \pm 0.1$ eV. | | | | | |

$\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0) / \Gamma_{\text{total}}$ Γ_{83} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|--------------------------------|
| 10.0 ± 2.5 ± 1.8 | 65 | ABLIKIM | 07D BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(K_1(1270)^\pm K^\mp) / \Gamma_{\text{total}}$ Γ_{85} / Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------------------|---------|-----------|
| 10.0 ± 1.8 ± 2.1 | ¹ BAI | 99C BES | $e^+ e^-$ |

¹ Assuming $B(K_1(1270) \rightarrow K \rho) = 0.42 \pm 0.06$

$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{86} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|--------|-------------|----------|--------------------------------|
| 2.20 ± 0.25 ± 0.37 | 83 ± 9 | ABLIKIM | 050 BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\rho^0 \rho^0) / \Gamma_{\text{total}}$ Γ_{87} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|---------|---|
| 0.5 ± 0.1 ± 0.2 | 61.1 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow \rho^0 \pi^+ \pi^-$ |

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{88} / Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|--------------|------|-----------|
| 6.7 ± 2.5 | TANENBAUM 78 | MRK1 | $e^+ e^-$ |

$\Gamma(2(\pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{89} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------------------------------|--------------|---------|---|
| 2.4 ± 0.6 OUR AVERAGE | Error includes scale factor of 2.2. | | | |
| 2.2 ± 0.2 ± 0.2 | 308 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$ |
| 4.5 ± 1.0 | | TANENBAUM 78 | MRK1 | $e^+ e^-$ |

$\Gamma(\rho^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{90} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------------------------------|--------------|---------|---|
| 2.2 ± 0.6 OUR AVERAGE | Error includes scale factor of 1.4. | | | |
| 2.0 ± 0.2 ± 0.4 | 285.5 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$ |
| 4.2 ± 1.5 | | TANENBAUM 78 | MRK1 | $e^+ e^-$ |

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_{91} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-------|---------------------|-----------|--|
| 12.6 ± 0.9 OUR AVERAGE | | | | |
| 18.9 ± 5.7 ± 0.3 | 32 | ¹ AUBERT | 07AU BABR | 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$ |
| 11.7 ± 1.0 ± 1.5 | 597 | ABLIKIM | 06G BES2 | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| 12.7 ± 0.5 ± 1.0 | 711.6 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (44 \pm 13 \pm 3) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{92}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $5.9 \pm 2.0 \pm 0.9$ | 19 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

$\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{93}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $8.6 \pm 1.3 \pm 1.8$ | 238 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

$\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{94}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $9.6 \pm 2.2 \pm 1.7$ | 133 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

$\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{95}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $7.3 \pm 2.2 \pm 1.4$ | 78 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

$\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{96}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $6.1 \pm 1.3 \pm 1.2$ | 125 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

$\Gamma(\eta K^+ K^-, \text{no } \eta\phi)/\Gamma_{\text{total}}$ Γ_{97}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|--------------|-------------|------|---|
| $3.08 \pm 0.29 \pm 0.25$ | 0.3k | ¹ | ABLIKIM | 12L | BES3 $\psi(2S) \rightarrow K^+ K^- \gamma \gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----|----|--------|----|------|--|
| <13 | 90 | BRIERE | 05 | CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
|-----|----|--------|----|------|--|

¹ Excluding $\eta\phi$.

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{98}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------------------|------|---|
| 1.62 ± 0.11 OUR AVERAGE | | Error includes scale factor of 1.1. | | |
| $1.56 \pm 0.04 \pm 0.11$ | 2.8k | ABLIKIM | 14G | BES3 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| $2.38 \pm 0.37 \pm 0.29$ | 78 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| $1.9 \pm 0.3 \pm 0.3$ | 76.8 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| $1.5 \pm 0.3 \pm 0.2$ | 23 | ¹ BAI | 03B | BES $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(\omega K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{99}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|--|
| 20.7 ± 2.6 OUR AVERAGE | | | | |
| $18.9 \pm 2.9 \pm 2.2$ | 396 | ABLIKIM | 13M | BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |
| $22.6 \pm 3.0 \pm 2.4$ | 535 | ABLIKIM | 13M | BES3 $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

$\Gamma(\omega K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{100}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 6.1 ± 1.2 OUR AVERAGE | | | | |
| 6.39 ± 1.50 ± 0.78 | 128 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |
| 5.86 ± 1.61 ± 0.83 | 143 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

$\Gamma(\omega \bar{K}^*(892)^0 K^0)/\Gamma_{\text{total}}$ Γ_{101}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 16.8 ± 2.5 ± 1.6 | 356 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

$\Gamma(\omega \bar{K}_2^*(1430)^0 K^0)/\Gamma_{\text{total}}$ Γ_{102}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 5.82 ± 2.08 ± 0.72 | 116 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

$\Gamma(\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{103}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| 1.60 ± 0.27 ± 0.24 | 109 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

¹ X(1440) compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

$\Gamma(\omega X(1440) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{104}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| 1.09 ± 0.20 ± 0.16 | 82 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

¹ X(1440) compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

$\Gamma(\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{105}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| 0.302 ± 0.098 ± 0.027 | 22 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

¹ Statistical significance 4.5 σ . This measurement is equivalent to a limit of $< 0.478 \times 10^{-5}$ at 90% C.L.

$\Gamma(\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{106}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| 0.125 ± 0.070 ± 0.013 | 10 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

¹ Statistical significance 3.2 σ . This measurement is equivalent to a limit of $< 0.221 \times 10^{-5}$ at 90% C.L.

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{107}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 3.5 ± 2.0 OUR AVERAGE | | | | Error includes scale factor of 2.8. |
| 5.45 ± 0.42 ± 0.87 | 671 | ABLIKIM | 05H BES2 | $e^+ e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+ \pi^-)$ |

| | | | | |
|-----------|--|------------------------|---------|-----------|
| 1.5 ± 1.0 | | ¹ TANENBAUM | 78 MRK1 | $e^+ e^-$ |
|-----------|--|------------------------|---------|-----------|

¹ Assuming entirely strong decay.

$\Gamma(\rho\bar{\rho}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ **Γ_{108}/Γ**

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| $7.3\pm 0.4\pm 0.6$ | 434.9 | BRIERE 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}\pi^+\pi^-\pi^0$ |

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$ **Γ_{109}/Γ**

| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------|----------------------------|-------------|-----------------------------------|
| $7.48\pm 0.23\pm 0.39$ | | 1.3k | ¹ METREVELI 12 | | $\psi(2S) \rightarrow K^+K^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| $6.2 \pm 1.5 \pm 0.2$ | | 66 | ^{2,3} LEES 15J | BABR | $e^+e^- \rightarrow K^+K^-\gamma$ |
| $8.3 \pm 1.5 \pm 0.2$ | | 66 | ^{3,4} LEES 15J | BABR | $e^+e^- \rightarrow K^+K^-\gamma$ |
| $6.3 \pm 0.6 \pm 0.3$ | | | ⁵ DOBBS 06A | CLEO | e^+e^- |
| 10 ± 7 | | | ⁵ BRANDELIK 79C | DASP | e^+e^- |
| < 5 | 90 | | FELDMAN 77 | MRK1 | e^+e^- |

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² $\sin\phi > 0$.

³ Using $\Gamma(\psi(2S) \rightarrow e^+e^-) = (2.37 \pm 0.04) \text{ keV}$.

⁴ $\sin\phi < 0$.

⁵ Interference with non-resonant K^+K^- production not taken into account.

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ **Γ_{110}/Γ**

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------|---------------------------|-------------|--|
| 5.34 ± 0.33 OUR AVERAGE | | | | |
| $5.28\pm 0.25\pm 0.34$ | 478 ± 23 | ¹ METREVELI 12 | | $\psi(2S) \rightarrow K_S^0 K_L^0$ |
| $5.8 \pm 0.8 \pm 0.4$ | | DOBBS 06A | CLEO | e^+e^- |
| $5.24\pm 0.47\pm 0.48$ | 156 ± 14 | ² BAI 04B | BES2 | $\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$ |

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6860 \pm 0.0027$.


$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ **Γ_{111}/Γ**

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------|--------------------------|-------------|---|
| 2.01 ± 0.17 OUR AVERAGE | | | | Error includes scale factor of 1.7. See the ideogram below. |
| $2.14\pm 0.03^{+0.12}_{-0.11}$ | 7k | ¹ ABLIKIM 12H | BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| $1.81\pm 0.18\pm 0.19$ | 260 ± 19 | ² ABLIKIM 05J | BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| $1.88^{+0.16}_{-0.15} \pm 0.28$ | 194 | ADAM 05 | CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| 0.85 ± 0.46 | 4 | FRANKLIN 83 | MRK2 | $e^+e^- \rightarrow \text{hadrons}$ |

¹ From $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$ events directly. The quoted systematic error includes a contribution of 4% (added in quadrature) from the uncertainty on the number of $\psi(2S)$ events.

² From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

WEIGHTED AVERAGE
 2.01 ± 0.17 (Error scaled by 1.7)



$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{112}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------------|----------------------|----------|--|
| 1.94 ± 0.25 $+1.15$ -0.34 | ¹ ABLIKIM | 05J BES2 | $\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$ |

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

$\Gamma(\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{113}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|----------------------|----------|---|
| 0.32 ± 0.12 OUR AVERAGE | | | | | Error includes scale factor of 1.8. |
| $0.51 \pm 0.07 \pm 0.11$ | | | ¹ ABLIKIM | 05J BES2 | $\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$ |
| $0.24^{+0.08}_{-0.07} \pm 0.02$ | | 22 | ADAM | 05 CLEO | $e^+e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|---|---------------------|---------|----------|
| <0.83 | 90 | 1 | FRANKLIN | 83 MRK2 | e^+e^- |
| <10 | 90 | | BARTEL | 76 CNTR | e^+e^- |
| <10 | 90 | | ² ABRAMS | 75 MRK1 | e^+e^- |

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

² Final state $\rho^0\pi^0$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{114}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|------------------------|----------|-----------------------------------|
| 0.78 ± 0.26 OUR AVERAGE | | | | | |
| $0.76 \pm 0.25 \pm 0.06$ | | 30 | ¹ METREVELI | 12 | $\psi(2S) \rightarrow \pi^+\pi^-$ |
| 8 ± 5 | | | BRANDELIK | 79C DASP | e^+e^- |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <2.1 | 90 | | DOBBS | 06A CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| <5 | 90 | | FELDMAN | 77 MRK1 | e^+e^- |

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. Using $\psi(3770) \rightarrow \pi^+\pi^-$ for continuum subtraction.

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{115}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------------------|---------|----------|
| <3.1 | 90 | ¹ BAI | 99C BES | e^+e^- |

¹ Assuming $B(K_1(1400) \rightarrow K^*\pi) = 0.94 \pm 0.06$

$\Gamma(K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{116}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|--------------|-------------|----------|-------------------------------|
| $7.12 \pm 0.62 \pm \frac{1.13}{-0.61}$ | 251 ± 22 | ABLIKIM | 12L BES3 | $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{117}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------|----------|-------------------------------------|
| $4.07 \pm 0.16 \pm 0.26$ | | 0.9k | ABLIKIM | 12L BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <8.9 | 90 | 1 | FRANKLIN | 83 MRK2 | $e^+e^- \rightarrow \text{hadrons}$ |

$\Gamma(K^+K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{120}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-----|---------------|-------------|----------|-------------------------------|
| 2.9 ± 0.4 OUR AVERAGE Error includes scale factor of 1.2. | | | | | |
| $3.18 \pm 0.30 \pm \frac{0.26}{-0.31}$ | | 0.2k | ABLIKIM | 12L BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| $2.9 \pm \frac{1.3}{-1.7} \pm 0.4$ | | 9.6 ± 4.2 | ABLIKIM | 05I BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| $1.3 \pm \frac{1.0}{-0.7} \pm 0.3$ | | 7 | ADAM | 05 CLEO | $e^+e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------|----|--|----------|---------|-------------------------------------|
| <5.4 | 90 | | FRANKLIN | 83 MRK2 | $e^+e^- \rightarrow \text{hadrons}$ |
|------|----|--|----------|---------|-------------------------------------|

$\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{121}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------|----------------|-------------|----------|-------------------------------|
| 10.9 ± 2.0 OUR AVERAGE | | | | |
| $13.3 \pm \frac{2.4}{-2.8} \pm 1.7$ | 65.6 ± 9.0 | ABLIKIM | 05I BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| $9.2 \pm \frac{2.7}{-2.2} \pm 0.9$ | 25 | ADAM | 05 CLEO | $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(K^+ K^{*(892)-} + c.c.) / \Gamma(K^{*(892)^0} \bar{K}^0 + c.c.)$ $\Gamma_{120} / \Gamma_{121}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|-------------------------------------|
| 0.16 ± 0.06 OUR AVERAGE | | | |
| 0.22 ^{+0.10} _{-0.14} | ABLIKIM | 05I | BES2 $e^+ e^- \rightarrow \psi(2S)$ |
| 0.14 ^{+0.08} _{-0.06} | ADAM | 05 | CLEO $e^+ e^- \rightarrow \psi(2S)$ |


$\Gamma(\phi \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{122} / Γ

| VALUE (units 10 ⁻⁴) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------------|-----------------------|------|---|
| 1.18 ± 0.26 OUR AVERAGE | | | | Error includes scale factor of 1.5. See the ideogram below. |
| 2.3 ± 0.8 ± 0.1 | 19 ± 6 | LEES | 12F | BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |
| 0.9 ± 0.2 ± 0.1 | 47.6 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| 1.5 ± 0.2 ± 0.2 | 51.5 ± 8.3 | ¹ BAI | 03B | BES $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 2.45 ± 0.96 ± 0.04 | 10 ± 4 | ^{2,3} AUBERT | 07AK | BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.



$\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{123} / Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-----------------------|-------------|---|
| 0.75 ± 0.33 OUR AVERAGE Error includes scale factor of 1.6. | | | | |
| 1.5 ± 0.5 ± 0.1 | 12 ± 4 | LEES | 12F BABR | 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |
| 0.6 ± 0.2 ± 0.1 | 18.4 ± 6.4 | ¹ BAI | 03B BES | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.46 ± 0.71 ± 0.02 | 6 ± 3 | ^{2,3} AUBERT | 07AK BABR | 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.34 \pm 0.16 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

$\Gamma(2(K^+ K^-)) / \Gamma_{\text{total}}$ Γ_{124} / Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 0.63 ± 0.13 OUR AVERAGE | | | | |
| 0.9 ± 0.4 ± 0.1 | 13 | LEES | 12F BABR | 10.6 $e^+ e^- \rightarrow 2(K^+ K^-) \gamma$ |
| 0.6 ± 0.1 ± 0.1 | 59.2 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$ |

$\Gamma(\phi K^+ K^-) / \Gamma_{\text{total}}$ Γ_{125} / Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 0.70 ± 0.16 OUR AVERAGE | | | | |
| 0.8 ± 0.2 ± 0.1 | 36.8 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$ |
| 0.6 ± 0.2 ± 0.1 | 16.1 ± 5.0 | ¹ BAI | 03B BES | $\psi(2S) \rightarrow 2(K^+ K^-)$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(2(K^+ K^-) \pi^0) / \Gamma_{\text{total}}$ Γ_{126} / Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 1.1 ± 0.2 ± 0.2 | | | | |
| | 44.7 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-) \pi^0$ |

$\Gamma(\phi \eta) / \Gamma_{\text{total}}$ Γ_{127} / Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--------------------------------|
| 3.10 ± 0.31 OUR AVERAGE | | | | |
| 3.14 ± 0.23 ± 0.23 | 0.2k | ABLIKIM | 12L BES3 | $e^+ e^- \rightarrow \psi(2S)$ |
| 2.0 $^{+1.5}_{-1.1}$ ± 0.4 | 6 | ADAM | 05 CLEO | $e^+ e^- \rightarrow \psi(2S)$ |
| 3.3 ± 1.1 ± 0.5 | 17 | ABLIKIM | 04k BES | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\eta \phi(2170), \phi(2170) \rightarrow \phi f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{128} / Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------------|------------|--------------------|-------------|--|
| < 2.2 × 10⁻⁶ | | | | |
| | 90 | ABLIKIM | 19i BES3 | $e^+ e^- \rightarrow \eta \phi f_0(980)$ |

$\Gamma(\phi\eta')/\Gamma_{\text{total}}$ Γ_{129}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|------|-----------------------------------|
| $3.1 \pm 1.4 \pm 0.7$ | 8 | ¹ ABLIKIM | 04K | BES $e^+e^- \rightarrow \psi(2S)$ |

¹ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.

$\Gamma(\omega\eta')/\Gamma_{\text{total}}$ Γ_{130}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|------|-----------------------------------|
| $3.2^{+2.4}_{-2.0} \pm 0.7$ | 4 | ¹ ABLIKIM | 04K | BES $e^+e^- \rightarrow \psi(2S)$ |

¹ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{131}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|------------------------------------|
| 2.1 ± 0.6 OUR AVERAGE | | | | |
| $2.5^{+1.2}_{-1.0} \pm 0.2$ | 14 | ADAM | 05 | CLEO $e^+e^- \rightarrow \psi(2S)$ |
| $1.87^{+0.68}_{-0.62} \pm 0.28$ | 14 | ABLIKIM | 04L | BES $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(\rho\eta')/\Gamma_{\text{total}}$ Γ_{132}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|-----------------------------------|
| $1.87^{+1.64}_{-1.11} \pm 0.33$ | 2 | ABLIKIM | 04L | BES $e^+e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------------|-----|----------------------|------|------------------------------------|
| $1.02 \pm 0.11 \pm 0.24$ | 143 | ¹ ABLIKIM | 17AK | BES3 $e^+e^- \rightarrow \psi(2S)$ |
| $0.569 \pm 0.128 \pm 0.236$ | 80 | ² ABLIKIM | 17AK | BES3 $e^+e^- \rightarrow \psi(2S)$ |

¹ Destructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+\pi^-\eta'$.

² Constructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+\pi^-\eta'$.

$\Gamma(\rho\eta)/\Gamma_{\text{total}}$ Γ_{133}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|-------------------------------------|
| 2.2 ± 0.6 OUR AVERAGE | | | | Error includes scale factor of 1.1. |
| $3.0^{+1.1}_{-0.9} \pm 0.2$ | 18 | ADAM | 05 | CLEO $e^+e^- \rightarrow \psi(2S)$ |
| $1.78^{+0.67}_{-0.62} \pm 0.17$ | 13 | ABLIKIM | 04L | BES $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$ Γ_{134}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-----|-------------|------|------------------------------------|
| <1.1 | 90 | ADAM | 05 | CLEO $e^+e^- \rightarrow \psi(2S)$ |
| <3.1 | 90 | ABLIKIM | 04K | BES $e^+e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ Γ_{135}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|-------------------------------|
| <0.04 | 90 | ABLIKIM 12L | BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.7 | 90 | ADAM 05 | CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| <0.4 | 90 | ABLIKIM 04K | BES | $e^+e^- \rightarrow \psi(2S)$ |

 $\Gamma(\eta_c\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{136}/Γ

| VALUE (units 10^{-3}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|-------------------------------|
| <1.0 | 90 | PEDLAR 07 | CLEO | $e^+e^- \rightarrow \psi(2S)$ |

 $\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$ Γ_{137}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|--|
| $2.7 \pm 0.6 \pm 0.4$ | 30.1 | BRIERE 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$ |

 $\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{138}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|--------------------------|------|-----------------------------|
| $0.81 \pm 0.11 \pm 0.14$ | 50 | ¹ ABLIKIM 08C | BES2 | $e^+e^- \rightarrow J/\psi$ |
| ¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$. | | | | |

 $\Gamma(\phi f_2'(1525))/\Gamma_{\text{total}}$ Γ_{139}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------------|------|-------------|------|----------------------------------|
| $0.44 \pm 0.12 \pm 0.11$ | 20 ± 6 | | BAI 04C | | $\psi(2S) \rightarrow 2(K^+K^-)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <0.45 | 90 | | BAI 98J | BES | $e^+e^- \rightarrow 2(K^+K^-)$ |

 $\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{140}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|----------|
| <0.88 | 90 | BAI 04G | BES2 | e^+e^- |

 $\Gamma(\Theta(1540)K^-\bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$ Γ_{141}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|----------|
| <1.0 | 90 | BAI 04G | BES2 | e^+e^- |

 $\Gamma(\Theta(1540)K_S^0\bar{p} \rightarrow K_S^0\bar{p}K^+n)/\Gamma_{\text{total}}$ Γ_{142}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|----------|
| <0.70 | 90 | BAI 04G | BES2 | e^+e^- |

 $\Gamma(\bar{\Theta}(1540)K^+n \rightarrow K_S^0\bar{p}K^+n)/\Gamma_{\text{total}}$ Γ_{143}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|----------|
| <2.6 | 90 | BAI 04G | BES2 | e^+e^- |

 $\Gamma(\bar{\Theta}(1540)K_S^0p \rightarrow K_S^0pK^-\bar{n})/\Gamma_{\text{total}}$ Γ_{144}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|----------|
| <0.60 | 90 | BAI 04G | BES2 | e^+e^- |

| $\Gamma(K_S^0 \bar{K}_S^0)/\Gamma_{\text{total}}$ | | | Γ_{145}/Γ | | |
|---|--------------------|-------------|-----------------------|----------|--|
| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | | |
| <0.046 | ¹ BAI | 04D | BES | e^+e^- | |
| ¹ Forbidden by <i>CP</i> . | | | | | |

| $\Gamma(\Lambda_c^+ \bar{p} e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$ | | | Γ_{146}/Γ | | |
|---|------------|-------------|-----------------------|-------------|------------------------------------|
| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| <1.7 × 10⁻⁶ | 90 | 450M | ABLIKIM | 18Q | BES3 $e^+e^- \rightarrow \psi(2S)$ |


————— **RADIATIVE DECAYS** —————

| $\Gamma(\gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$ | | | Γ_{147}/Γ | | |
|---|-------------|-----------------------|-----------------------|--------------------|-------------------------------|
| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 9.79 ± 0.20 | | | | OUR FIT | |
| 9.33 ± 0.26 | | | | OUR AVERAGE | |
| 9.389 ± 0.014 ± 0.332 | 4.7M | ABLIKIM | 17U | BES3 | $e^+e^- \rightarrow \gamma X$ |
| 9.22 ± 0.11 ± 0.46 | 72k | ATHAR | 04 | CLEO | $e^+e^- \rightarrow \gamma X$ |
| 9.9 ± 0.5 ± 0.8 | | ¹ GAISER | 86 | CBAL | $e^+e^- \rightarrow \gamma X$ |
| 7.2 ± 2.3 | | ¹ BIDDICK | 77 | CNTR | $e^+e^- \rightarrow \gamma X$ |
| 7.5 ± 2.6 | | ¹ WHITAKER | 76 | MRK1 | e^+e^- |
| ¹ Angular distribution ($1+\cos^2\theta$) assumed. | | | | | |

| $\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$ | | | Γ_{148}/Γ | | |
|---|-------------|----------------------|-----------------------|--------------------|-------------------------------|
| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 9.75 ± 0.24 | | | | OUR FIT | |
| 9.54 ± 0.29 | | | | OUR AVERAGE | |
| 9.905 ± 0.011 ± 0.353 | 5.0M | ABLIKIM | 17U | BES3 | $e^+e^- \rightarrow \gamma X$ |
| 9.07 ± 0.11 ± 0.54 | 76k | ATHAR | 04 | CLEO | $e^+e^- \rightarrow \gamma X$ |
| 9.0 ± 0.5 ± 0.7 | | ¹ GAISER | 86 | CBAL | $e^+e^- \rightarrow \gamma X$ |
| 7.1 ± 1.9 | | ² BIDDICK | 77 | CNTR | $e^+e^- \rightarrow \gamma X$ |
| ¹ Angular distribution ($1-0.189 \cos^2\theta$) assumed. | | | | | |
| ² Valid for isotropic distribution of the photon. | | | | | |

| $\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$ | | | Γ_{149}/Γ | | |
|---|-------------|----------------------|-----------------------|--------------------|-------------------------------|
| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 9.52 ± 0.20 | | | | OUR FIT | |
| 9.42 ± 0.31 | | | | OUR AVERAGE | |
| Error includes scale factor of 1.3. See the ideogram below. | | | | | |
| 9.621 ± 0.013 ± 0.272 | 4.2M | ABLIKIM | 17U | BES3 | $e^+e^- \rightarrow \gamma X$ |
| 9.33 ± 0.14 ± 0.61 | 79k | ATHAR | 04 | CLEO | $e^+e^- \rightarrow \gamma X$ |
| 8.0 ± 0.5 ± 0.7 | | ¹ GAISER | 86 | CBAL | $e^+e^- \rightarrow \gamma X$ |
| 7.0 ± 2.0 | | ² BIDDICK | 77 | CNTR | $e^+e^- \rightarrow \gamma X$ |
| ¹ Angular distribution ($1-0.052 \cos^2\theta$) assumed. | | | | | |
| ² Valid for isotropic distribution of the photon. | | | | | |

WEIGHTED AVERAGE
 9.42 ± 0.31 (Error scaled by 1.3)



$$\frac{[\Gamma(\gamma\chi_{c0}(1P)) + \Gamma(\gamma\chi_{c1}(1P)) + \Gamma(\gamma\chi_{c2}(1P))]/\Gamma_{\text{total}}}{(\Gamma_{147} + \Gamma_{148} + \Gamma_{149})/\Gamma_{\text{total}}}$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$27.6 \pm 0.3 \pm 2.0$ ¹ ATHAR 04 CLEO $e^+e^- \rightarrow \gamma X$

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$$\frac{\Gamma(\gamma\chi_{c0}(1P))}{\Gamma(\gamma\chi_{c1}(1P))} \quad \Gamma_{147}/\Gamma_{148}$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.02 \pm 0.01 \pm 0.07$ ¹ ATHAR 04 CLEO $e^+e^- \rightarrow \gamma X$

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$$\frac{\Gamma(\gamma\chi_{c2}(1P))}{\Gamma(\gamma\chi_{c1}(1P))} \quad \Gamma_{149}/\Gamma_{148}$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.03 \pm 0.02 \pm 0.03$ ¹ ATHAR 04 CLEO $e^+e^- \rightarrow \gamma X$

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$$\frac{\Gamma(\gamma\chi_{c0}(1P))}{\Gamma(\gamma\chi_{c2}(1P))} \quad \Gamma_{147}/\Gamma_{149}$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.99 \pm 0.02 \pm 0.08$ ¹ ATHAR 04 CLEO $e^+e^- \rightarrow \gamma X$

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.


$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$ **Γ_{150}/Γ**

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|--------------------|---|------|------------------------------------|
| 0.34 ± 0.05 | OUR AVERAGE | Error includes scale factor of 1.3. See the ideogram below. | | |
| $0.432 \pm 0.016 \pm 0.060$ | | MITCHELL | 09 | CLEO $e^+e^- \rightarrow \gamma X$ |
| $0.32 \pm 0.04 \pm 0.06$ | 2.5k | ¹ ATHAR | 04 | CLEO $e^+e^- \rightarrow \gamma X$ |
| 0.28 ± 0.06 | | ² GAISER | 86 | CBAL $e^+e^- \rightarrow \gamma X$ |

¹ATHAR 04 used $\Gamma_{\eta_c(1S)} = 24.8 \pm 4.9$ MeV to obtain this result.

²GAISER 86 used $\Gamma_{\eta_c(1S)} = 11.5 \pm 4.5$ MeV to obtain this result.

WEIGHTED AVERAGE
 0.34 ± 0.05 (Error scaled by 1.3)



$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$ **Γ_{151}/Γ**

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|-----|-----------------------------|------|---|
| $7 \pm 2 \pm 4$ | | ¹ ABLIKIM | 12G | BES3 $\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$ |
| < 8 | 90 | ² CRONIN-HEN..10 | CLEO | $\psi(2S) \rightarrow \gamma K \bar{K} \pi$ |
| < 20 | 90 | ATHAR | 04 | CLEO $e^+e^- \rightarrow \gamma X$ |
| 20–130 | 95 | EDWARDS | 82C | CBAL $e^+e^- \rightarrow \gamma X$ |

¹ABLIKIM 12G reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] = (1.30 \pm 0.20 \pm 0.30) \times 10^{-5}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (1.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²CRONIN-HENNESSY 10 reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] < 14.5 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = 1.9 \times 10^{-2}$. This measurement assumes $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ Γ_{152}/Γ

| VALUE (units 10^{-6}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-----------------------|----------|------------------------------------|
| 1.04 ± 0.22 OUR AVERAGE Error includes scale factor of 1.4. | | | | | |
| 0.95 ± 0.16 ± 0.05 | | 423 | ABLIKIM | 17X BES3 | $\psi(2S) \rightarrow \gamma\pi^0$ |
| 1.58 ± 0.40 ± 0.13 | | 37 | ABLIKIM | 10F BES3 | $\psi(2S) \rightarrow \gamma\pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| < 5 | 90 | | PEDLAR | 09 CLE3 | $\psi(2S) \rightarrow \gamma X$ |
| < 5400 | 95 | | ¹ LIBERMAN | 75 SPEC | e^+e^- |
| < 1×10^4 | 90 | | WIIK | 75 DASP | e^+e^- |

¹ Restated by us using $B(\psi(2S) \rightarrow \mu^+\mu^-) = 0.0077$.

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ Γ_{153}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|--------------------------|----------|---|
| 1.24 ± 0.04 OUR AVERAGE | | | | | |
| 1.251 ± 0.022 ± 0.062 | | 56K | ABLIKIM | 17X BES3 | $\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta, \gamma\pi^0\pi^0\eta$ |
| 1.26 ± 0.03 ± 0.08 | | 2226 | ¹ ABLIKIM | 10F BES3 | $\psi(2S) \rightarrow 3\gamma\pi^+\pi^-, 2\gamma\pi^+\pi^-$ |
| 1.19 ± 0.08 ± 0.03 | | | PEDLAR | 09 CLE3 | $\psi(2S) \rightarrow \gamma X$ |
| 1.24 ± 0.27 ± 0.15 | | 23 | ABLIKIM | 06R BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| 1.54 ± 0.31 ± 0.20 | | ~ 43 | BAI | 98F BES | $\psi(2S) \rightarrow \pi^+\pi^-2\gamma, \pi^+\pi^-3\gamma$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| < 60 | 90 | | ² BRAUNSCH... | 77 DASP | e^+e^- |
| < 11 | 90 | | ³ BARTEL | 76 CNTR | e^+e^- |

¹ Combining the results from $\eta' \rightarrow \pi^+\pi^-\eta$ and $\eta' \rightarrow \pi^+\pi^-\gamma$ decay modes.

² Restated by us using total decay width 228 keV.

³ The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{154}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|--------------|------------------|---------|---|
| 2.73^{+0.29}_{-0.25} OUR AVERAGE Error includes scale factor of 1.8. | | | | |
| 2.84 ± 0.15 ^{+0.03} _{-0.10} | 1.9k | 1,2 DOBBS | 15 | $\psi(2S) \rightarrow \gamma\pi\pi$ |
| 2.12 ± 0.19 ± 0.32 | | 3,4 BAI | 03C BES | $\psi(2S) \rightarrow \gamma\pi\pi$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 2.08 ± 0.19 ± 0.33 | 200.6 ± 18.8 | ³ BAI | 03C BES | $\psi(2S) \rightarrow \gamma\pi^+\pi^-$ |
| 2.90 ± 1.08 ± 1.07 | 29.9 ± 11.1 | ³ BAI | 03C BES | $\psi(2S) \rightarrow \gamma\pi^0\pi^0$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (2.39 \pm 0.09 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

⁴ Combining the results from $\pi^+\pi^-$ and $\pi^0\pi^0$ decay modes.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{155}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|--------------------|------|---|
| 3.1±1.0±1.4 | 175 | ¹ DOBBS | 15 | $\psi(2S) \rightarrow \gamma K \bar{K}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$ Γ_{156}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|----------------------|------|---------------------------------------|
| 9.3±1.8±0.6 | 274 | ^{1,2} DOBBS | 15 | $\psi(2S) \rightarrow \gamma \pi \pi$ |

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_0(1500))/\Gamma_{\text{total}}] \times [B(f_0(1500) \rightarrow \pi \pi)] = (3.2 \pm 0.6 \pm 0.2) \times 10^{-5}$ which we divide by our best value $B(f_0(1500) \rightarrow \pi \pi) = (34.5 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ Γ_{157}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|----------------------|------|---|
| 3.3±0.8±0.1 | 136 | ^{1,2} DOBBS | 15 | $\psi(2S) \rightarrow \gamma K \bar{K}$ |

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = (2.9 \pm 0.6 \pm 0.3) \times 10^{-5}$ which we divide by our best value $B(f'_2(1525) \rightarrow K \bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{159}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|------------|--------------------|---------|---|
| 3.5 ±0.6 OUR AVERAGE | | | | |
| 3.6 ±0.4 ±0.5 | 290 | ¹ DOBBS | 15 | $\psi(2S) \rightarrow \gamma \pi \pi$ |
| 3.01±0.41±1.24 | 35.6 ± 4.8 | ² BAI | 03C BES | $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{160}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|------------|------|--------------------|---------|---|
| 6.6 ±0.7 OUR AVERAGE | | | | | |
| 6.7 ±0.6 ±0.6 | | 375 | ¹ DOBBS | 15 | $\psi(2S) \rightarrow \gamma K \bar{K}$ |
| 6.04±0.90±1.32 | 39.6 ± 5.9 | | ^{2,3} BAI | 03C BES | $\psi(2S) \rightarrow \gamma K^+ K^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 15.6 90 6.8 ± 3.1 ^{2,3} BAI 03C BES $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Includes unknown branching fractions to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied the $K^+ K^-$ result by a factor of 2 and the $K_S^0 K_S^0$ result by a factor of 4 to obtain the $K \bar{K}$ result.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{161}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|--------------------|------|---------------------------------------|
| 4.8±0.5±0.9 | 373 | ¹ DOBBS | 15 | $\psi(2S) \rightarrow \gamma \pi \pi$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{162}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|--------------------|------|---|
| $3.2 \pm 0.6 \pm 0.8$ | 207 | ¹ DOBBS | 15 | $\psi(2S) \rightarrow \gamma K \bar{K}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{163}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|----------------------|------|---------------------------------------|
| $< 5.8 \times 10^{-6}$ | 90 | ^{1,2} DOBBS | 15 | $\psi(2S) \rightarrow \gamma \pi \pi$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+ \pi^-$ and $\pi^0 \pi^0$ are $3.2/4.3 \times 10^{-6}$ and $2.6/4.0 \times 10^{-6}$, respectively.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{164}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|----------------------|------|---|
| $< 9.5 \times 10^{-6}$ | 90 | ^{1,2} DOBBS | 15 | $\psi(2S) \rightarrow \gamma K \bar{K}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $2.1/4.3 \times 10^{-6}$ and $3.7/5.5 \times 10^{-6}$, respectively.

$\Gamma(\gamma \eta)/\Gamma_{\text{total}}$ Γ_{166}/Γ

| VALUE (units 10^{-6}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|----------------------|----------|--|
| 0.92 ± 0.18 OUR AVERAGE | | | | | |
| $0.85 \pm 0.18 \pm 0.04$ | | 382 | ¹ ABLIKIM | 17X BES3 | $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$, $\gamma 3\pi^0$ |
| $1.38 \pm 0.48 \pm 0.09$ | | 13 | ¹ ABLIKIM | 10F BES3 | $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$, $\gamma 3\pi^0$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---------|----|--------|-----|------|--|
| < 2 | 90 | PEDLAR | 09 | CLE3 | $\psi(2S) \rightarrow \gamma X$ |
| < 90 | 90 | BAI | 98F | BES | $\psi(2S) \rightarrow \pi^+ \pi^- 3\gamma$ |
| < 200 | 90 | YAMADA | 77 | DASP | $e^+ e^- \rightarrow 3\gamma$ |

¹ Combining the results from $\eta \rightarrow \pi^+ \pi^- \pi^0$ and $\eta \rightarrow 3\pi^0$ decay modes.

$\Gamma(\gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{167}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|---|
| $8.71 \pm 1.25 \pm 1.64$ | 418 | ABLIKIM | 06R | BES2 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$ |

$\Gamma(\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$ Γ_{169}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----|-------------|------|--|
| < 0.9 | 90 | ABLIKIM | 06R | BES2 $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---------|----|----------------------|-----|------|---|
| < 1.3 | 90 | ABLIKIM | 06R | BES2 | $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ |
| < 1.2 | 90 | ¹ SCHARRE | 80 | MRK1 | $e^+ e^-$ |

¹ Includes unknown branching fraction $\eta(1405) \rightarrow K \bar{K} \pi$.

$\Gamma(\gamma \eta(1405) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{170}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|---|
| $0.36 \pm 0.25 \pm 0.05$ | 10 | ABLIKIM | 06R | BES2 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$ |

| $\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ | | | | | Γ_{171}/Γ |
|---|-----|-------------|------|---------|--|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| $<5.0 \times 10^{-7}$ | 90 | ABLIKIM | 17AJ | BES3 | $\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$ |

| $\Gamma(\gamma\eta(1475) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}$ | | | | | Γ_{173}/Γ |
|---|-----|-------------|------|---------|---|
| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT | |
| <1.4 | 90 | ABLIKIM | 06R | BES2 | $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <1.5 | 90 | ABLIKIM | 06R | BES2 | $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$ |

| $\Gamma(\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ | | | | | Γ_{174}/Γ |
|---|-----|-------------|------|---------|---|
| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT | |
| <0.88 | 90 | ABLIKIM | 06R | BES2 | $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$ |

| $\Gamma(\gamma 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ | | | | | Γ_{175}/Γ |
|--|------|-------------|------|---------|-------------------------------|
| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $39.6 \pm 2.8 \pm 5.0$ | 583 | ABLIKIM | 07D | BES2 | $e^+e^- \rightarrow \psi(2S)$ |

| $\Gamma(\gamma K^{*0} K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ | | | | | Γ_{176}/Γ |
|---|------|-------------|------|---------|-------------------------------|
| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $37.0 \pm 6.1 \pm 7.2$ | 237 | ABLIKIM | 07D | BES2 | $e^+e^- \rightarrow \psi(2S)$ |

| $\Gamma(\gamma K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$ | | | | | Γ_{177}/Γ |
|--|------|-------------|------|---------|-------------------------------|
| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $24.0 \pm 4.5 \pm 5.0$ | 41 | ABLIKIM | 07D | BES2 | $e^+e^- \rightarrow \psi(2S)$ |

| $\Gamma(\gamma K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ | | | | | Γ_{178}/Γ |
|--|------|-------------|------|---------|-------------------------------|
| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $25.6 \pm 3.6 \pm 3.6$ | 115 | ABLIKIM | 07D | BES2 | $e^+e^- \rightarrow \psi(2S)$ |

| $\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ | | | | | Γ_{179}/Γ |
|--|------|-------------|------|---------|-------------------------------|
| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $19.1 \pm 2.7 \pm 4.3$ | 132 | ABLIKIM | 07D | BES2 | $e^+e^- \rightarrow \psi(2S)$ |

| $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$ | | | | | Γ_{180}/Γ |
|---|------|------------------------|------|---------|--|
| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 3.9 ± 0.5 OUR AVERAGE | | | | | Error includes scale factor of 2.0. |
| $4.18 \pm 0.26 \pm 0.18$ | 348 | ¹ ALEXANDER | 10 | CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |
| $2.9 \pm 0.4 \pm 0.4$ | 142 | ABLIKIM | 07D | BES2 | $e^+e^- \rightarrow \psi(2S)$ |

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

$\Gamma(\gamma f_2(1950) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{181}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------------|------|--|
| $1.2 \pm 0.2 \pm 0.1$ | 111 | ¹ ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

$\Gamma(\gamma f_2(2150) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{182}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|---------------------------|------|--|
| $0.72 \pm 0.18 \pm 0.03$ | 73 | ¹ ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

$\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{183}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|--------------------------------------|
| $4.57 \pm 0.36^{+1.77}_{-4.26}$ | | ABLIKIM 12D | BES3 | $J/\psi \rightarrow \gamma p\bar{p}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|--------------|------|--|
| <1.6 | 90 | ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |
| <5.4 | 90 | ABLIKIM 07D | BES | $\psi(2S) \rightarrow \gamma p\bar{p}$ |

$\Gamma(\gamma X \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{184}/Γ

For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|--------------|------|--|
| <2 | 90 | ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |

$\Gamma(\gamma \pi^+ \pi^- p\bar{p})/\Gamma_{\text{total}}$ Γ_{185}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|--------------------------------|
| $2.8 \pm 1.2 \pm 0.7$ | 17 | ABLIKIM 07D | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\gamma 2(\pi^+ \pi^-) K^+ K^-)/\Gamma_{\text{total}}$ Γ_{186}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|--------------------------------|
| <22 | 90 | ABLIKIM 07D | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{187}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|--------------------------------|
| <17 | 90 | ABLIKIM 07D | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\gamma K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{188}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|--------------------------------|
| <4 | 90 | ABLIKIM 07D | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{189}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|------------------------------------|
| $3.1 \pm 0.6^{+0.8}_{-1.0}$ | 1.1k | ABLIKIM | 120 | BES3 $e^+e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---------------|------|----------------------|-----|---|
| 3.2 ± 0.6 | 1.1k | ¹ ABLIKIM | 17N | BES3 $\psi(2S) \rightarrow \gamma\gamma J/\psi$ |
|---------------|------|----------------------|-----|---|

¹ Uses $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$. No systematic error estimation.

$\Gamma(e^+e^-\eta')/\Gamma_{\text{total}}$ Γ_{190}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|--|
| 1.90 ± 0.26 OUR AVERAGE | | | | |
| $1.99 \pm 0.33 \pm 0.12$ | 57 | ABLIKIM | 18Z | BES3 $\psi(2S) \rightarrow \eta' e^+e^-$, $\eta' \rightarrow \gamma\pi^+\pi^-$ |
| $1.79 \pm 0.38 \pm 0.11$ | 20 | ABLIKIM | 18Z | BES3 $\psi(2S) \rightarrow \eta' e^+e^-$, $\eta' \rightarrow \eta\pi^+\pi^-$ |

$\Gamma(e^+e^-\chi_{c0}(1P))/\Gamma_{\text{total}}$ Γ_{191}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|----------------------|------|---|
| $10.6 \pm 2.4 \pm 0.4$ | 48 | ¹ ABLIKIM | 17I | BES3 $\psi(2S) \rightarrow e^+e^-\gamma J/\psi$ |

¹ ABLIKIM 17I reports $(11.7 \pm 2.5 \pm 1.0) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+e^-\chi_{c0}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.27 \pm 0.06) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.40 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(e^+e^-\chi_{c1}(1P))/\Gamma_{\text{total}}$ Γ_{192}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|------|---|
| $8.5 \pm 0.6 \pm 0.2$ | 873 | ¹ ABLIKIM | 17I | BES3 $\psi(2S) \rightarrow e^+e^-\gamma J/\psi$ |

¹ ABLIKIM 17I reports $(8.6 \pm 0.3 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+e^-\chi_{c1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (33.9 \pm 1.2) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(e^+e^-\chi_{c2}(1P))/\Gamma_{\text{total}}$ Γ_{193}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|------|---|
| $7.0 \pm 0.7 \pm 0.2$ | 227 | ¹ ABLIKIM | 17I | BES3 $\psi(2S) \rightarrow e^+e^-\gamma J/\psi$ |

¹ ABLIKIM 17I reports $(6.9 \pm 0.5 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+e^-\chi_{c2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.2 \pm 0.7) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.0 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(e^+e^-\chi_{c0}(1P))/\Gamma(\gamma\chi_{c0}(1P))$ $\Gamma_{191}/\Gamma_{147}$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|--------------------------|------|--|
| $9.4 \pm 1.9 \pm 0.6$ | 48 | ¹ ABLIKIM 17I | BES3 | $\psi(2S) \rightarrow e^+e^-\gamma J/\psi$ |
| ¹ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors. | | | | |

$\Gamma(e^+e^-\chi_{c1}(1P))/\Gamma(\gamma\chi_{c1}(1P))$ $\Gamma_{192}/\Gamma_{148}$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|--------------------------|------|--|
| $8.3 \pm 0.3 \pm 0.4$ | 873 | ¹ ABLIKIM 17I | BES3 | $\psi(2S) \rightarrow e^+e^-\gamma J/\psi$ |
| ¹ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors. | | | | |

$\Gamma(e^+e^-\chi_{c2}(1P))/\Gamma(\gamma\chi_{c2}(1P))$ $\Gamma_{193}/\Gamma_{149}$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|--------------------------|------|--|
| $6.6 \pm 0.5 \pm 0.4$ | 227 | ¹ ABLIKIM 17I | BES3 | $\psi(2S) \rightarrow e^+e^-\gamma J/\psi$ |
| ¹ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (199.6 \pm 0.8 \pm 7.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors. | | | | |

———— WEAK DECAYS ————

$\Gamma(D^0 e^+e^- + c.c.)/\Gamma_{total}$ Γ_{194}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|---------------------------|------|-------------------------------|
| $< 1.4 \times 10^{-7}$ | 90 | ¹ ABLIKIM 17AF | BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| ¹ Using D^0 decays to $K^-\pi^+$, $K^-\pi^+\pi^0$, and $K^-\pi^+\pi^+\pi^-$. | | | | |

———— OTHER DECAYS ————

$\Gamma(\text{invisible})/\Gamma(e^+e^-)$ Γ_{195}/Γ_6

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----|-------------|------|---------------------------------|
| < 2.0 | 90 | LEES 13I | BABR | $B \rightarrow K^{(*)}\psi(2S)$ |

$\psi(2S)$ CROSS-PARTICLE BRANCHING RATIOS

For measurements involving $B(\psi(2S) \rightarrow \gamma\chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$ see the corresponding entries in the $\chi_{cJ}(1P)$ sections.

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

$\psi(2S) \rightarrow \gamma\chi_{cJ}(1P)$ and $\chi_{cJ} \rightarrow \gamma J/\psi(1S)$

$a_2(\chi_{c1})/a_2(\chi_{c2})$ Magnetic quadrupole transition amplitude ratio

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|--------------------------|------|---|
| 63 ± 7 OUR AVERAGE | | | | |
| 61.7 ± 8.3 | 253k | ¹ ABLIKIM 17N | BES3 | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| 67^{+19}_{-13} | 59k | ² ARTUSO 09 | CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |

¹ Statistical and systematic errors combined.

² Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $a_2(\chi_{c1}(1P))$ and $a_2(\chi_{c2}(1P))$ from ARTUSO 09.

$b_2(\chi_{c2})/b_2(\chi_{c1})$ Magnetic quadrupole transition amplitude ratio

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| 60±31 OUR AVERAGE | | | | |
| 74±40 | 253k | ¹ ABLIKIM | 17N BES3 | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| 37 ⁺⁵³ ₋₄₇ | 59k | ² ARTUSO | 09 CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |

¹ Statistical and systematic errors combined. Derived from the reported measurement of $b_2(\chi_{c1})/b_2(\chi_{c2}) = 1.35 \pm 0.72$.

² Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $b_2(\chi_{c1}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

 $\psi(2S)$ REFERENCES

| | | | | |
|-----------|------|----------------|----------------------------|---------------------|
| ABLIKIM | 19I | PR D99 012014 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 19N | PR D99 032006 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 18Q | PR D97 091102 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 18T | PR D98 032006 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 18Z | PL B783 452 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ANASHIN | 18 | PL B781 174 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |
| LEES | 18E | PR D98 112015 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 17AF | PR D96 111101 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 17AJ | PR D96 112008 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 17AK | PR D96 112012 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 17E | PL B770 217 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 17I | PRL 118 221802 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 17L | PR D95 052003 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 17N | PR D95 072004 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 17U | PR D96 032001 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 17X | PR D96 052003 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| DOBBS | 17 | PR D96 092004 | S. Dobbs <i>et al.</i> | (NWES, WAYN) |
| LEES | 17A | PR D95 052001 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| AAIJ | 16Y | JHEP 1605 132 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 16L | PR D93 072003 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 15I | PR D91 092006 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 15V | PL B749 414 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ANASHIN | 15 | PL B749 50 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |
| DOBBS | 15 | PR D91 052006 | S. Dobbs <i>et al.</i> | (NWES) |
| LEES | 15J | PR D92 072008 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 14G | PR D89 112006 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| DOBBS | 14 | PL B739 90 | S. Dobbs <i>et al.</i> | (NWES, WAYN) |
| ABLIKIM | 13A | PRL 110 022001 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 13D | PR D87 012007 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 13F | PR D87 052007 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 13M | PR D87 092006 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 13R | PR D88 032007 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 13S | PR D88 032010 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 13W | PR D88 112007 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| LEES | 13I | PR D87 112005 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 13O | PR D87 092005 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 13Q | PR D88 032013 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 13Y | PR D88 072009 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| AAIJ | 12H | EPJ C72 1972 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 12D | PRL 108 112003 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 12G | PRL 109 042003 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 12H | PL B710 594 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 12L | PR D86 072011 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 12M | PR D86 092008 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 12O | PRL 109 172002 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 12Q | CP C36 1040 | M. Ablikim <i>et al.</i> | (BES II Collab.) |
| ANASHIN | 12 | PL B711 280 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |
| LEES | 12E | PR D85 112009 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 12F | PR D86 012008 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| METREVELI | 12 | PR D85 092007 | Z. Metreveli <i>et al.</i> | (NWES, FLOR, WAYN+) |

| | | | | |
|---------------|------|-------------------------------|-----------------------------------|------------------------|
| GE | 11 | PR D84 032008 | J.Y. Ge <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 10B | PRL 104 132002 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 10F | PRL 105 261801 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ALEXANDER | 10 | PR D82 092002 | J.P. Alexander <i>et al.</i> | (CLEO Collab.) |
| CRONIN-HEN... | 10 | PR D81 052002 | D. Cronin-Hennessey <i>et al.</i> | (CLEO Collab.) |
| ADAMS | 09 | PR D80 051106 | G.S. Adams <i>et al.</i> | (CLEO Collab.) |
| ARTUSO | 09 | PR D80 112003 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| LIBBY | 09 | PR D80 072002 | J. Libby <i>et al.</i> | (CLEO Collab.) |
| MITCHELL | 09 | PRL 102 011801 | R.E. Mitchell <i>et al.</i> | (CLEO Collab.) |
| PEDLAR | 09 | PR D79 111101 | T.K. Pedlar <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 08B | PL B659 74 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 08C | PL B659 789 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| DOBBS | 08A | PRL 101 182003 | S. Dobbs <i>et al.</i> | (CLEO Collab.) |
| MENDEZ | 08 | PR D78 011102 | H. Mendez <i>et al.</i> | (CLEO Collab.) |
| PDG | 08 | PL B667 1 | C. Amsler <i>et al.</i> | (PDG Collab.) |
| ABLIKIM | 07C | PL B648 149 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 07D | PRL 99 011802 | M. Ablikim <i>et al.</i> | (BES II Collab.) |
| ABLIKIM | 07H | PR D76 092003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ANASHIN | 07 | JETPL 85 347 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |
| | | Translated from ZETFP 85 429. | | |
| ANDREOTTI | 07 | PL B654 74 | M. Andreotti <i>et al.</i> | (Femilab E835 Collab.) |
| AUBERT | 07AK | PR D76 012008 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07AU | PR D76 092005 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| Also | | PR D77 119902E (errata.) | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07BD | PR D76 092006 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| PDG | 07 | Unofficial 2007 WWW edition | | (PDG Collab.) |
| PEDLAR | 07 | PR D75 011102 | T.K. Pedlar <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 06G | PR D73 052004 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06I | PR D74 012004 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06L | PRL 97 121801 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06R | PR D74 072001 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06W | PR D74 112003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ADAM | 06 | PRL 96 082004 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| AUBERT | 06B | PR D73 012005 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06D | PR D73 052003 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 06D | PR D74 091103 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| DOBBS | 06A | PR D74 011105 | S. Dobbs <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 05E | PR D71 072006 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05H | PR D72 012002 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05I | PL B614 37 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05J | PL B619 247 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05O | PL B630 21 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ADAM | 05 | PRL 94 012005 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| ADAM | 05A | PRL 94 232002 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| ANDREOTTI | 05 | PR D71 032006 | M. Andreotti <i>et al.</i> | (FNAL E835 Collab.) |
| AUBERT | 05D | PR D71 052001 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| BRIERE | 05 | PRL 95 062001 | R.A. Briere <i>et al.</i> | (CLEO Collab.) |
| PEDLAR | 05 | PR D72 051108 | T.K. Pedlar <i>et al.</i> | (CLEO Collab.) |
| ROSNER | 05 | PRL 95 102003 | J.L. Rosner <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 04B | PR D70 012003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 04K | PR D70 112003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 04L | PR D70 112007 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ATHAR | 04 | PR D70 112002 | S.B. Athar <i>et al.</i> | (CLEO Collab.) |
| BAI | 04B | PRL 92 052001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04C | PR D69 072001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04D | PL B589 7 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04G | PR D70 012004 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04I | PR D70 012006 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| PDG | 04 | PL B592 1 | S. Eidelman <i>et al.</i> | (PDG Collab.) |
| SETH | 04 | PR D69 097503 | K.K. Seth | |
| AULCHENKO | 03 | PL B573 63 | V.M. Aulchenko <i>et al.</i> | (KEDR Collab.) |
| BAI | 03B | PR D67 052002 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 03C | PR D67 032004 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| AUBERT | 02B | PR D65 031101 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| BAI | 02 | PR D65 052004 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 02B | PL B550 24 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 02C | PRL 88 101802 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| PDG | 02 | PR D66 010001 | K. Hagiwara <i>et al.</i> | (PDG Collab.) |
| BAI | 01 | PR D63 032002 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| AMBROGIANI | 00A | PR D62 032004 | M. Ambrogiani <i>et al.</i> | (FNAL E835 Collab.) |
| ARTAMONOV | 00 | PL B474 427 | A.S. Artamonov <i>et al.</i> | |

| | | | | |
|-------------|------|------------------------------|-------------------------------|------------------------------|
| BAI | 00 | PRL 84 594 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 99C | PRL 83 1918 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 98E | PR D57 3854 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 98F | PR D58 097101 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 98J | PRL 81 5080 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| ARMSTRONG | 97 | PR D55 1153 | T.A. Armstrong <i>et al.</i> | (E760 Collab.) |
| GRIBUSHIN | 96 | PR D53 4723 | A. Gribushin <i>et al.</i> | (E672 Collab., E706 Collab.) |
| ARMSTRONG | 93B | PR D47 772 | T.A. Armstrong <i>et al.</i> | (FNAL E760 Collab.) |
| ALEXANDER | 89 | NP B320 45 | J.P. Alexander <i>et al.</i> | (LBL, MICH, SLAC) |
| COHEN | 87 | RMP 59 1121 | E.R. Cohen, B.N. Taylor | (RISC, NBS) |
| GAISER | 86 | PR D34 711 | J. Gaiser <i>et al.</i> | (Crystal Ball Collab.) |
| KURAEV | 85 | SJNP 41 466 | E.A. Kuraev, V.S. Fadin | (NOVO) |
| | | Translated from YAF 41 733. | | |
| FRANKLIN | 83 | PRL 51 963 | M.E.B. Franklin <i>et al.</i> | (LBL, SLAC) |
| EDWARDS | 82C | PRL 48 70 | C. Edwards <i>et al.</i> | (CIT, HARV, PRIN+) |
| LEMOIGNE | 82 | PL 113B 509 | Y. Lemoigne <i>et al.</i> | (SACL, LOIC, SHMP+) |
| HIMEL | 80 | PRL 44 920 | T. Himel <i>et al.</i> | (LBL, SLAC) |
| OREGLIA | 80 | PRL 45 959 | M.J. Oreglia <i>et al.</i> | (SLAC, CIT, HARV+) |
| SCHARRE | 80 | PL 97B 329 | D.L. Scharre <i>et al.</i> | (SLAC, LBL) |
| ZHOLENTZ | 80 | PL 96B 214 | A.A. Zholents <i>et al.</i> | (NOVO) |
| | Also | SJNP 34 814 | A.A. Zholents <i>et al.</i> | (NOVO) |
| | | Translated from YAF 34 1471. | | |
| BRANDELIK | 79B | NP B160 426 | R. Brandelik <i>et al.</i> | (DASP Collab.) |
| BRANDELIK | 79C | ZPHY C1 233 | R. Brandelik <i>et al.</i> | (DASP Collab.) |
| BARTEL | 78B | PL 79B 492 | W. Bartel <i>et al.</i> | (DESY, HEIDP) |
| TANENBAUM | 78 | PR D17 1731 | W.M. Tanenbaum <i>et al.</i> | (SLAC, LBL) |
| BIDDICK | 77 | PRL 38 1324 | C.J. Biddick <i>et al.</i> | (UCSD, UMD, PAVI+) |
| BRAUNSCH... | 77 | PL 67B 249 | W. Braunschweig <i>et al.</i> | (DASP Collab.) |
| BURMESTER | 77 | PL 66B 395 | J. Burmester <i>et al.</i> | (DESY, HAMB, SIEG+) |
| FELDMAN | 77 | PRPL 33C 285 | G.J. Feldman, M.L. Perl | (LBL, SLAC) |
| YAMADA | 77 | Hamburg Conf. 69 | S. Yamada | (DASP Collab.) |
| BARTEL | 76 | PL 64B 483 | W. Bartel <i>et al.</i> | (DESY, HEIDP) |
| TANENBAUM | 76 | PRL 36 402 | W.M. Tanenbaum <i>et al.</i> | (SLAC, LBL) IG |
| WHITAKER | 76 | PRL 37 1596 | J.S. Whitaker <i>et al.</i> | (SLAC, LBL) |
| ABRAMS | 75 | Stanford Symp. 25 | G.S. Abrams | (LBL) |
| ABRAMS | 75B | PRL 34 1181 | G.S. Abrams <i>et al.</i> | (LBL, SLAC) |
| BOYARSKI | 75C | Palermo Conf. 54 | A.M. Boyarski <i>et al.</i> | (SLAC, LBL) |
| HILGER | 75 | PRL 35 625 | E. Hilger <i>et al.</i> | (STAN, PENN) |
| LIBERMAN | 75 | Stanford Symp. 55 | A.D. Liberman | (STAN) |
| LUTH | 75 | PRL 35 1124 | V. Luth <i>et al.</i> | (SLAC, LBL) JPC |
| WIIK | 75 | Stanford Symp. 69 | B.H. Wiik | (DESY) |