

$\Upsilon(2S)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

### $\Upsilon(2S)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10023.26±0.31 OUR AVERAGE</b>			
10023.5 ±0.5	<sup>1</sup> ARTAMONOV 00	MD1	$e^+e^- \rightarrow \text{hadrons}$
10023.1 ±0.4	BARBER 84	REDE	$e^+e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10023.6 ±0.5	<sup>2,3</sup> BARU	86B REDE	$e^+e^- \rightarrow \text{hadrons}$
<sup>1</sup> Reanalysis of BARU 86B using new electron mass (COHEN 87).			
<sup>2</sup> Reanalysis of ARTAMONOV 84.			
<sup>3</sup> Superseded by ARTAMONOV 00.			

### $m\Upsilon(3S) - m\Upsilon(2S)$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>331.50±0.02±0.13</b>	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$

### $\Upsilon(2S)$ WIDTH

VALUE (keV)	DOCUMENT ID	COMMENT
<b>31.98±2.63 OUR EVALUATION</b>		See the Note on "Width Determinations of the $\Upsilon$ States"

### $\Upsilon(2S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $\Upsilon(1S)\pi^+\pi^-$	(17.85±0.26) %	
$\Gamma_2$ $\Upsilon(1S)\pi^0\pi^0$	(8.6 ± 0.4) %	
$\Gamma_3$ $\tau^+\tau^-$	(2.00±0.21) %	
$\Gamma_4$ $\mu^+\mu^-$	(1.93±0.17) %	S=2.2
$\Gamma_5$ $e^+e^-$	(1.91±0.16) %	
$\Gamma_6$ $\Upsilon(1S)\pi^0$	< 4 × 10 <sup>-5</sup>	CL=90%
$\Gamma_7$ $\Upsilon(1S)\eta$	(2.9 ± 0.4) × 10 <sup>-4</sup>	S=2.0
$\Gamma_8$ $J/\psi(1S)$ anything	< 6 × 10 <sup>-3</sup>	CL=90%
$\Gamma_9$ $J/\psi(1S)\eta_c$	< 5.4 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{10}$ $J/\psi(1S)\chi_{c0}$	< 3.4 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{11}$ $J/\psi(1S)\chi_{c1}$	< 1.2 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{12}$ $J/\psi(1S)\chi_{c2}$	< 2.0 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{13}$ $J/\psi(1S)\eta_c(2S)$	< 2.5 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{14}$ $J/\psi(1S)X(3940)$	< 2.0 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{15}$ $J/\psi(1S)X(4160)$	< 2.0 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{16}$ $\chi_{c1}$ anything	(2.2 ± 0.5) × 10 <sup>-4</sup>	

$\Gamma_{17}$	$\chi_{c1}(1P)^0 X_{tetra}$	$< 3.67$	$\times 10^{-5}$	CL=90%
$\Gamma_{18}$	$\chi_{c2}$ anything	$( 2.3 \pm 0.8 )$	$\times 10^{-4}$	
$\Gamma_{19}$	$\psi(2S)\eta_c$	$< 5.1$	$\times 10^{-6}$	CL=90%
$\Gamma_{20}$	$\psi(2S)\chi_{c0}$	$< 4.7$	$\times 10^{-6}$	CL=90%
$\Gamma_{21}$	$\psi(2S)\chi_{c1}$	$< 2.5$	$\times 10^{-6}$	CL=90%
$\Gamma_{22}$	$\psi(2S)\chi_{c2}$	$< 1.9$	$\times 10^{-6}$	CL=90%
$\Gamma_{23}$	$\psi(2S)\eta_c(2S)$	$< 3.3$	$\times 10^{-6}$	CL=90%
$\Gamma_{24}$	$\psi(2S)X(3940)$	$< 3.9$	$\times 10^{-6}$	CL=90%
$\Gamma_{25}$	$\psi(2S)X(4160)$	$< 3.9$	$\times 10^{-6}$	CL=90%
$\Gamma_{26}$	$Z_c(3900)^+ Z_c(3900)^-$	$< 1.0$	$\times 10^{-6}$	CL=90%
$\Gamma_{27}$	$Z_c(4200)^+ Z_c(4200)^-$	$< 1.67$	$\times 10^{-5}$	CL=90%
$\Gamma_{28}$	$Z_c(3900)^\pm Z_c(4200)^\mp$	$< 7.3$	$\times 10^{-6}$	CL=90%
$\Gamma_{29}$	$X(4050)^+ X(4050)^-$	$< 1.35$	$\times 10^{-5}$	CL=90%
$\Gamma_{30}$	$X(4250)^+ X(4250)^-$	$< 2.67$	$\times 10^{-5}$	CL=90%
$\Gamma_{31}$	$X(4050)^\pm X(4250)^\mp$	$< 2.72$	$\times 10^{-5}$	CL=90%
$\Gamma_{32}$	$Z_c(4430)^+ Z_c(4430)^-$	$< 2.03$	$\times 10^{-5}$	CL=90%
$\Gamma_{33}$	$X(4055)^\pm X(4055)^\mp$	$< 1.11$	$\times 10^{-5}$	CL=90%
$\Gamma_{34}$	$X(4055)^\pm Z_c(4430)^\mp$	$< 2.11$	$\times 10^{-5}$	CL=90%
$\Gamma_{35}$	$\overline{2H}$ anything	$( 2.78_{-0.26}^{+0.30} )$	$\times 10^{-5}$	S=1.2
$\Gamma_{36}$	hadrons	$(94 \pm 11)$	%	
$\Gamma_{37}$	$ggg$	$(58.8 \pm 1.2)$	%	
$\Gamma_{38}$	$\gamma gg$	$( 1.87 \pm 0.28 )$	%	
$\Gamma_{39}$	$\phi K^+ K^-$	$( 1.6 \pm 0.4 )$	$\times 10^{-6}$	
$\Gamma_{40}$	$\omega \pi^+ \pi^-$	$< 2.58$	$\times 10^{-6}$	CL=90%
$\Gamma_{41}$	$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$( 2.3 \pm 0.7 )$	$\times 10^{-6}$	
$\Gamma_{42}$	$\phi f_2^l(1525)$	$< 1.33$	$\times 10^{-6}$	CL=90%
$\Gamma_{43}$	$\omega f_2(1270)$	$< 5.7$	$\times 10^{-7}$	CL=90%
$\Gamma_{44}$	$\rho(770) a_2(1320)$	$< 8.8$	$\times 10^{-7}$	CL=90%
$\Gamma_{45}$	$K^*(892)^0 \overline{K}_2^*(1430)^0 + \text{c.c.}$	$( 1.5 \pm 0.6 )$	$\times 10^{-6}$	
$\Gamma_{46}$	$K_1(1270)^\pm K^\mp$	$< 3.22$	$\times 10^{-6}$	CL=90%
$\Gamma_{47}$	$K_1(1400)^\pm K^\mp$	$< 8.3$	$\times 10^{-7}$	CL=90%
$\Gamma_{48}$	$b_1(1235)^\pm \pi^\mp$	$< 4.0$	$\times 10^{-7}$	CL=90%
$\Gamma_{49}$	$\rho \pi$	$< 1.16$	$\times 10^{-6}$	CL=90%
$\Gamma_{50}$	$\pi^+ \pi^- \pi^0$	$< 8.0$	$\times 10^{-7}$	CL=90%
$\Gamma_{51}$	$\omega \pi^0$	$< 1.63$	$\times 10^{-6}$	CL=90%
$\Gamma_{52}$	$\pi^+ \pi^- \pi^0 \pi^0$	$( 1.30 \pm 0.28 )$	$\times 10^{-5}$	
$\Gamma_{53}$	$K_S^0 K^+ \pi^- + \text{c.c.}$	$( 1.14 \pm 0.33 )$	$\times 10^{-6}$	
$\Gamma_{54}$	$K^*(892)^0 \overline{K}^0 + \text{c.c.}$	$< 4.22$	$\times 10^{-6}$	CL=90%
$\Gamma_{55}$	$K^*(892)^- K^+ + \text{c.c.}$	$< 1.45$	$\times 10^{-6}$	CL=90%
$\Gamma_{56}$	$f_1(1285)$ anything	$( 2.2 \pm 1.6 )$	$\times 10^{-3}$	
$\Gamma_{57}$	$f_1(1285) X_{tetra}$	$< 6.47$	$\times 10^{-5}$	CL=90%
$\Gamma_{58}$	Sum of 100 exclusive modes	$( 2.90 \pm 0.30 )$	$\times 10^{-3}$	

### Radiative decays

$\Gamma_{59}$	$\gamma\chi_{b1}(1P)$		$(6.9 \pm 0.4) \%$	
$\Gamma_{60}$	$\gamma\chi_{b2}(1P)$		$(7.15 \pm 0.35) \%$	
$\Gamma_{61}$	$\gamma\chi_{b0}(1P)$		$(3.8 \pm 0.4) \%$	
$\Gamma_{62}$	$\gamma f_0(1710)$	$< 5.9$	$\times 10^{-4}$	CL=90%
$\Gamma_{63}$	$\gamma f_2'(1525)$	$< 5.3$	$\times 10^{-4}$	CL=90%
$\Gamma_{64}$	$\gamma f_2(1270)$	$< 2.41$	$\times 10^{-4}$	CL=90%
$\Gamma_{65}$	$\gamma f_J(2220)$			
$\Gamma_{66}$	$\gamma\eta_c(1S)$	$< 2.7$	$\times 10^{-5}$	CL=90%
$\Gamma_{67}$	$\gamma\chi_{c0}$	$< 1.0$	$\times 10^{-4}$	CL=90%
$\Gamma_{68}$	$\gamma\chi_{c1}$	$< 3.6$	$\times 10^{-6}$	CL=90%
$\Gamma_{69}$	$\gamma\chi_{c2}$	$< 1.5$	$\times 10^{-5}$	CL=90%
$\Gamma_{70}$	$\gamma\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi$	$< 8$	$\times 10^{-7}$	CL=90%
$\Gamma_{71}$	$\gamma\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0 J/\psi$	$< 2.4$	$\times 10^{-6}$	CL=90%
$\Gamma_{72}$	$\gamma X(3915) \rightarrow \omega J/\psi$	$< 2.8$	$\times 10^{-6}$	CL=90%
$\Gamma_{73}$	$\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi$	$< 1.2$	$\times 10^{-6}$	CL=90%
$\Gamma_{74}$	$\gamma X(4350) \rightarrow \phi J/\psi$	$< 1.3$	$\times 10^{-6}$	CL=90%
$\Gamma_{75}$	$\gamma\eta_b(1S)$		$(5.5 \pm_{-0.9}^{+1.1}) \times 10^{-4}$	S=1.2
$\Gamma_{76}$	$\gamma\eta_b(1S) \rightarrow \gamma$ Sum of 26 exclusive modes	$< 3.7$	$\times 10^{-6}$	CL=90%
$\Gamma_{77}$	$\gamma X_{b\bar{b}} \rightarrow \gamma$ Sum of 26 exclusive modes	$< 4.9$	$\times 10^{-6}$	CL=90%
$\Gamma_{78}$	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] $< 1.95$	$\times 10^{-4}$	CL=95%
$\Gamma_{79}$	$\gamma A^0 \rightarrow \gamma$ hadrons	$< 8$	$\times 10^{-5}$	CL=90%
$\Gamma_{80}$	$\gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$	$< 8.3$	$\times 10^{-6}$	CL=90%

### Lepton Family number (*LF*) violating modes

$\Gamma_{81}$	$e^\pm \tau^\mp$	<i>LF</i>	$< 3.2$	$\times 10^{-6}$	CL=90%
$\Gamma_{82}$	$\mu^\pm \tau^\mp$	<i>LF</i>	$< 3.3$	$\times 10^{-6}$	CL=90%

[a]  $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

### CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 13 measurements and one constraint to determine 3 parameters. The overall fit has a  $\chi^2 = 11.8$  for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$$x_7 \begin{array}{|c} \hline 2 \\ \hline x_1 \end{array}$$

$\Upsilon(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$  $\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_4\Gamma_5/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>6.5±1.5±1.0</b>	KOBEL	92	CBAL $e^+e^- \rightarrow \mu^+\mu^-$

 $\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1\Gamma_5/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>105.4±1.0±4.2</b>	11.8K	<sup>1</sup> AUBERT	08BP BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$

<sup>1</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ . $\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{36}\Gamma_5/\Gamma$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.577±0.009 OUR AVERAGE</b>			
0.581±0.004±0.009	<sup>1</sup> ROSNER	06	CLEO $10.0 e^+e^- \rightarrow \text{hadrons}$
0.552±0.031±0.017	<sup>1</sup> BARU	96	MD1 $e^+e^- \rightarrow \text{hadrons}$
0.54 ±0.04 ±0.02	<sup>1</sup> JAKUBOWSKI	88	CBAL $e^+e^- \rightarrow \text{hadrons}$
0.58 ±0.03 ±0.04	<sup>2</sup> GILES	84B	CLEO $e^+e^- \rightarrow \text{hadrons}$
0.60 ±0.12 ±0.07	<sup>2</sup> ALBRECHT	82	DASP $e^+e^- \rightarrow \text{hadrons}$
0.54 ±0.07 <sup>+0.09</sup> <sub>-0.05</sub>	<sup>2</sup> NICZYPORUK	81C	LENA $e^+e^- \rightarrow \text{hadrons}$
0.41 ±0.18	<sup>2</sup> BOCK	80	CNTR $e^+e^- \rightarrow \text{hadrons}$

<sup>1</sup> Radiative corrections evaluated following KURAEV 85.<sup>2</sup> Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85. $\Upsilon(2S)$  PARTIAL WIDTHS $\Gamma(e^+e^-) \quad \Gamma_5$ 

VALUE (keV)	DOCUMENT ID
<b>0.612±0.011 OUR EVALUATION</b>	

 $\Upsilon(2S)$  BRANCHING RATIOS $\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$ 

Abbreviation MM in the COMMENT field below stands for missing mass.

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>17.85±0.26 OUR FIT</b>				
<b>17.92±0.26 OUR AVERAGE</b>				
16.8 ±1.1 ±1.3	906k	<sup>1</sup> LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
17.80±0.05±0.37	170k	<sup>2</sup> LEES	11L BABR	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
18.02±0.02±0.61	851k	<sup>3</sup> BHARI	09 CLEO	$e^+e^- \rightarrow \pi^+\pi^- \text{MM}$
17.22±0.17±0.75	11.8K	<sup>4</sup> AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
19.2 ±0.2 ±1.0	52.6k	<sup>5</sup> ALEXANDER	98 CLE2	$\pi^+\pi^-\ell^+\ell^-, \pi^+\pi^- \text{MM}$

18.1 ± 0.5 ± 1.0	11.6k	ALBRECHT	87	ARG	$e^+e^- \rightarrow \pi^+\pi^-$ MM
16.9 ± 4.0		GELPHMAN	85	CBAL	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
19.1 ± 1.2 ± 0.6		BESSON	84	CLEO	$\pi^+\pi^-$ MM
18.9 ± 2.6		FONSECA	84	CUSB	$e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$
21 ± 7	7	NICZYPORUK	81B	LENA	$e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$

<sup>1</sup> LEES 11C reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

<sup>3</sup> A weighted average of the inclusive and exclusive results.

<sup>4</sup> Using  $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$ ,  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$  and,  $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$  keV.

<sup>5</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$ .

### $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.6 ± 0.4 OUR AVERAGE</b>				
8.43 ± 0.16 ± 0.42	38k	<sup>1</sup> BHARI 09	CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.2 ± 0.6 ± 0.8	275	<sup>2</sup> ALEXANDER 98	CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.5 ± 1.9 ± 1.9	25	ALBRECHT 87	ARG	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
8.0 ± 1.5		GELPHMAN 85	CBAL	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
10.3 ± 2.3		FONSECA 84	CUSB	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

<sup>1</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$ .

<sup>2</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$ .

### $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ $\Gamma_2/\Gamma_1$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.462 ± 0.037 <sup>1</sup> BHARI 09 CLEO  $e^+e^- \rightarrow \Upsilon(2S)$

<sup>1</sup> Not independent of other values reported by BHARI 09.

### $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.00 ± 0.21 OUR AVERAGE</b>				
2.00 ± 0.12 ± 0.18	22k	<sup>1</sup> BESSON 07	CLEO	$e^+e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+\tau^-$
1.7 ± 1.5 ± 0.6		HAAS 84B	CLEO	$e^+e^- \rightarrow \tau^+\tau^-$


<sup>1</sup> BESSON 07 reports  $[\Gamma(\Upsilon(2S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = 1.04 \pm 0.04 \pm 0.05$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

$\Gamma_4/\Gamma$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0193±0.0017 OUR AVERAGE</b>			Error includes scale factor of 2.2. See the ideogram below.		
0.0203±0.0003±0.0008		120k	ADAMS	05	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
0.0122±0.0028±0.0019			<sup>1</sup> KOBEL	92	CBAL $e^+e^- \rightarrow \mu^+\mu^-$
0.0138±0.0025±0.0015			KAARSBERG	89	CSB2 $e^+e^- \rightarrow \mu^+\mu^-$
0.009 ±0.006 ±0.006			<sup>2</sup> ALBRECHT	85	ARG $e^+e^- \rightarrow \mu^+\mu^-$
0.018 ±0.008 ±0.005			HAAS	84B	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<0.038		90	NICZYPORUK	81C	LENA $e^+e^- \rightarrow \mu^+\mu^-$

<sup>1</sup> Taking into account interference between the resonance and continuum.  
<sup>2</sup> Re-evaluated using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 0.026$ .



$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$

$\Gamma_3/\Gamma_4$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.04±0.04±0.05</b>	22k	BESSION	07	CLEO $e^+e^- \rightarrow \Upsilon(2S)$

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$

$\Gamma_6/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

< 4	90	<sup>1</sup> TAMPONI	13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^0$
< 18	90	<sup>2</sup> HE	08A	CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
<110	90	ALEXANDER	98	CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
<800	90	LURZ	87	CBAL	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

<sup>1</sup> TAMPONI 13 reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)] < 2.3 \times 10^{-4}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = 17.85 \times 10^{-2}$ .

<sup>2</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$ .

### $\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ $\Gamma_6/\Gamma_1$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;2.3</b>	90	TAMPONI	13	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^0$

### $\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$ $\Gamma_7/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	-------------	--------------------	-------------	----------------

**2.9 ± 0.4 OUR FIT** Error includes scale factor of 2.0.

**2.9 ± 0.4 OUR AVERAGE** Error includes scale factor of 1.9. See the ideogram below.

2.39 ± 0.31 ± 0.14      112      <sup>1</sup> LEES      11L      BABR       $\Upsilon(2S) \rightarrow \ell^+\ell^-\eta$

2.1  $^{+0.7}_{-0.6}$  ± 0.3      14      <sup>2</sup> HE      08A      CLEO       $e^+e^- \rightarrow \ell^+\ell^-\eta$

• • • We use the following data for averages but not for fits. • • •

3.55 ± 0.32 ± 0.05      241      <sup>3</sup> TAMPONI      13      BELL       $e^+e^- \rightarrow \Upsilon(1S)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9      90      <sup>1,4</sup> AUBERT      08BP      BABR       $e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$

< 28      90      ALEXANDER98      CLE2       $e^+e^- \rightarrow \ell^+\ell^-\eta$

< 50      90      ALBRECHT      87      ARG       $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$  MM

< 70      90      LURZ      87      CBAL       $e^+e^- \rightarrow \ell^+\ell^-(\gamma\gamma, 3\pi^0)$

< 100      90      BESSON      84      CLEO       $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$  MM


< 20      90      FONSECA      84      CUSB       $e^+e^- \rightarrow \ell^+\ell^-(\gamma\gamma, \pi^+\pi^-\pi^0)$

<sup>1</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

<sup>2</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$ .

<sup>3</sup> TAMPONI 13 reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\eta)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (17.85 \pm 0.26) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> Using  $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$  keV.



### $\Gamma(\tau(1S)\eta)/\Gamma(\tau(1S)\pi^+\pi^-)$

$\Gamma_7/\Gamma_1$

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.64±0.25 OUR FIT</b>					Error includes scale factor of 2.0.
<b>1.99±0.14±0.11</b>		241	TAMPONI 13	BELL	$e^+e^- \rightarrow \tau(1S)\eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1.35±0.17±0.08			<sup>1</sup> LEES 11L	BABR	$\tau(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$
< 5.2	90		<sup>2</sup> AUBERT 08BP	BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

<sup>1</sup> Not independent of other values reported by LEES 11L.

<sup>2</sup> Not independent of other values reported by AUBERT 08BP.

### $\Gamma(\tau(1S)\pi^0)/\Gamma(\tau(1S)\eta)$

$\Gamma_6/\Gamma_7$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.13	90	TAMPONI 13	BELL	$e^+e^- \rightarrow \tau(1S)\pi^0$

### $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$

$\Gamma_8/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.006	90	MASCHMANN 90	CBAL	$e^+e^- \rightarrow \text{hadrons}$

### $\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$

$\Gamma_9/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.4 × 10 <sup>-6</sup>	90	YANG 14	BELL	$e^+e^- \rightarrow J/\psi X$



$\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$					$\Gamma_{10}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.4 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow J/\psi X$
$\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$					$\Gamma_{11}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.2 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow J/\psi X$
$\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$					$\Gamma_{12}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.0 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow J/\psi X$
$\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_{13}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.5 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow J/\psi X$
$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$					$\Gamma_{14}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.0 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow J/\psi X$
$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$					$\Gamma_{15}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.0 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow J/\psi X$
$\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_{16}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.24 \pm 0.44 \pm 0.20$	376	JIA	17	BELL	$\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$
$\Gamma(\chi_{c1}(1P)^0 X_{\text{tetra}})/\Gamma_{\text{total}}$					$\Gamma_{17}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<36.7 \times 10^{-6}$	90	<sup>1</sup> JIA	17A	BELL	$e^+e^- \rightarrow \text{hadrons}$
<sup>1</sup> For a tetraquark state $X_{\text{tetra}}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of $X_{\text{tetra}}$ mass and width range from $4.4 \times 10^{-6}$ to $36.7 \times 10^{-6}$ .					
$\Gamma(\chi_{c2} \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_{18}/\Gamma$
VALUE (units $10^{-4}$ )		DOCUMENT ID	TECN	COMMENT	
$2.28 \pm 0.73 \pm 0.34$		JIA	17	BELL	$\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$
$\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$					$\Gamma_{19}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.1 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow \psi(2S)X$
$\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$					$\Gamma_{20}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<4.7 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow \psi(2S)X$

$\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$					$\Gamma_{21}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.5 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow \psi(2S)X$
$\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$					$\Gamma_{22}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow \psi(2S)X$
$\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_{23}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.3 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow \psi(2S)X$
$\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$					$\Gamma_{24}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow \psi(2S)X$
$\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$					$\Gamma_{25}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow \psi(2S)X$
$\Gamma(Z_c(3900)^+ Z_c(3900)^-)/\Gamma_{\text{total}}$					$\Gamma_{26}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.0 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL	$\Upsilon(2S) \rightarrow J/\psi \pi^\pm X$
<sup>1</sup> Assuming $B(Z_c(3900)^\pm \rightarrow J/\psi \pi^\pm) = 1$ .					
$\Gamma(Z_c(4200)^+ Z_c(4200)^-)/\Gamma_{\text{total}}$					$\Gamma_{27}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<16.7 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL	$\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$
<sup>1</sup> Assuming $B(Z_c(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1$					
$\Gamma(Z_c(3900)^\pm Z_c(4200)^\mp)/\Gamma_{\text{total}}$					$\Gamma_{28}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.3 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL	$\Upsilon(2S) \rightarrow J/\psi \pi^\pm X$
<sup>1</sup> Assuming $B(Z_c(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1 = B(Z_c(3900)^\pm \rightarrow J/\psi \pi^\pm)$ .					
$\Gamma(X(4050)^+ X(4050)^-)/\Gamma_{\text{total}}$					$\Gamma_{29}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<13.5 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL	$\Upsilon(2S) \rightarrow \chi_{c1}(1P)\pi^\pm X$
<sup>1</sup> Assuming $B(X(4050)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm)$					
$\Gamma(X(4250)^+ X(4250)^-)/\Gamma_{\text{total}}$					$\Gamma_{30}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<26.7 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL	$\Upsilon(2S) \rightarrow \chi_{c1}(1P)\pi^\pm X$
<sup>1</sup> Assuming $B(X(4250)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm) = 1$					

$\Gamma(X(4050)^\pm X(4250)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<27.2 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(2S) \rightarrow \chi_{c1}(1P)\pi^\pm X$
<sup>1</sup> Assuming $B(X(4050)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm) = 1 = B(X(4250)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm)$				

$\Gamma(Z_c(4430)^+ Z_c(4430)^-)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<20.3 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(2S) \rightarrow \psi(2S)\pi^\pm X$
<sup>1</sup> Assuming $B(Z_c(4430)^\pm \rightarrow \psi(2P)\pi^\pm) = 1$				

$\Gamma(X(4055)^\pm X(4055)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<11.1 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(2S) \rightarrow \psi(2S)\pi^\pm X$
<sup>1</sup> Assuming $B(X(4055)^\pm \rightarrow \psi(2S)\pi^\pm) = 1$				

$\Gamma(X(4055)^\pm Z_c(4430)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<21.1 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(2S) \rightarrow \psi(2S)\pi^\pm X$
<sup>1</sup> Assuming $B(X(4055)^\pm \rightarrow \psi(2S)\pi^\pm) = 1 = B(Z_c(4430)^\pm \rightarrow \psi(2S)\pi^\pm)$				

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**$2.78^{+0.30}_{-0.26}$  OUR AVERAGE** Error includes scale factor of 1.2.

$2.64 \pm 0.11^{+0.26}_{-0.21}$		LEES	14G	BABR $e^+e^- \rightarrow \overline{2H} X$
$3.37 \pm 0.50 \pm 0.25$	58	ASNER	07	CLEO $e^+e^- \rightarrow \overline{2H} X$

$\Gamma(g g g)/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>58.8 \pm 1.2</math></b>	6M	<sup>1</sup> BESSON	06A	CLEO $\Upsilon(2S) \rightarrow \text{hadrons}$

<sup>1</sup> Calculated using the value  $\Gamma(\gamma g g)/\Gamma(g g g) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$  from BESSON 06A and PDG 08 values of  $B(\pi^+ \pi^- \Upsilon(1S)) = (18.1 \pm 0.4)\%$ ,  $B(\pi^0 \pi^0 \Upsilon(1S)) = (8.6 \pm 0.4)\%$ ,  $B(\mu^+ \mu^-) = (1.93 \pm 0.17)\%$ , and  $R_{\text{hadrons}} = 3.51$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(\gamma g g)/\Gamma_{\text{total}}$  measurement of BESSON 06A.

$\Gamma(\gamma g g)/\Gamma(g g g)$   $\Gamma_{38}/\Gamma_{37}$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.18 \pm 0.04 \pm 0.47</math></b>	6M	BESSON	06A	CLEO $\Upsilon(2S) \rightarrow (\gamma +) \text{hadrons}$

$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.58 \pm 0.33 \pm 0.18</math></b>	58	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(K^+ K^-)$

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{40}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;2.58</b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$
$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{41}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b><math>2.32 \pm 0.40 \pm 0.54</math></b>	135	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$
$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$					$\Gamma_{42}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;1.33</b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(K^+ K^-)$
$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$					$\Gamma_{43}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;0.57</b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$
$\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$					$\Gamma_{44}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;0.88</b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$
$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{45}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b><math>1.53 \pm 0.52 \pm 0.19</math></b>	32	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$
$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$					$\Gamma_{46}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;3.22</b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$
$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$					$\Gamma_{47}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;0.83</b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$
$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$					$\Gamma_{48}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;0.40</b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$
$\Gamma(\rho\pi)/\Gamma_{\text{total}}$					$\Gamma_{49}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;1.16</b>	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0$
$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$					$\Gamma_{50}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;0.80</b>	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0$

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{51}/\Gamma$
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;1.63</b>	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+\pi^-\pi^0\pi^0$

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{52}/\Gamma$
VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>13.0 \pm 1.9 \pm 2.1</math></b>	$261 \pm 37$	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+\pi^-\pi^0\pi^0$

$\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{53}/\Gamma$	
VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>1.14 \pm 0.30 \pm 0.13</math></b>		$40 \pm 10$	SHEN	13	BELL	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.2	90	<sup>1</sup> DOBBS	12A		$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$
------	----	--------------------	-----	--	--

<sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{54}/\Gamma$
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;4.22</b>	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$

$\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{55}/\Gamma$
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;1.45</b>	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$

$\Gamma(f_1(1285) \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_{56}/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>2.20 \pm 1.50 \pm 0.63</math></b>	2.9k	JIA	17A	BELL	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(f_1(1285) X_{tetra})/\Gamma_{\text{total}}$					$\Gamma_{57}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;<math>64.7 \times 10^{-6}</math></b>	90	<sup>1</sup> JIA	17A	BELL	$e^+e^- \rightarrow \text{hadrons}$

<sup>1</sup> For a tetraquark state  $X_{tetra}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of  $X_{tetra}$  mass and width range from  $7.8 \times 10^{-6}$  to  $64.7 \times 10^{-6}$ .

$\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$					$\Gamma_{58}/\Gamma$
VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT		
<b><math>0.29 \pm 0.03</math></b>	1,2	DOBBS	12A		$\Upsilon(2S) \rightarrow \text{hadrons}$

<sup>1</sup> DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

<sup>2</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$						$\Gamma_{59}/\Gamma$
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>0.069 ± 0.004</b>	<b>OUR AVERAGE</b>					
0.0693 ± 0.0012 ± 0.0041	407k	ARTUSO	05	CLEO	$e^+e^- \rightarrow \gamma X$	
0.069 ± 0.005 ± 0.009		EDWARDS	99	CLE2	$\Upsilon(2S) \rightarrow \gamma\chi(1P)$	
0.091 ± 0.018 ± 0.022		ALBRECHT	85E	ARG	$e^+e^- \rightarrow \gamma\text{conv. } X$	
0.065 ± 0.007 ± 0.012		NERNST	85	CBAL	$e^+e^- \rightarrow \gamma X$	
0.080 ± 0.017 ± 0.016		HAAS	84	CLEO	$e^+e^- \rightarrow \gamma\text{conv. } X$	
0.059 ± 0.014		KLOPFEN...	83	CUSB	$e^+e^- \rightarrow \gamma X$	

$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$						$\Gamma_{60}/\Gamma$
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>0.0715 ± 0.0035</b>	<b>OUR AVERAGE</b>					
0.0724 ± 0.0011 ± 0.0040	410k	ARTUSO	05	CLEO	$e^+e^- \rightarrow \gamma X$	
0.074 ± 0.005 ± 0.008		EDWARDS	99	CLE2	$\Upsilon(2S) \rightarrow \gamma\chi(1P)$	
0.098 ± 0.021 ± 0.024		ALBRECHT	85E	ARG	$e^+e^- \rightarrow \gamma\text{conv. } X$	
0.058 ± 0.007 ± 0.010		NERNST	85	CBAL	$e^+e^- \rightarrow \gamma X$	
0.102 ± 0.018 ± 0.021		HAAS	84	CLEO	$e^+e^- \rightarrow \gamma\text{conv. } X$	
0.061 ± 0.014		KLOPFEN...	83	CUSB	$e^+e^- \rightarrow \gamma X$	

$\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$						$\Gamma_{61}/\Gamma$
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>0.038 ± 0.004</b>	<b>OUR AVERAGE</b>					
0.0375 ± 0.0012 ± 0.0047	198k	ARTUSO	05	CLEO	$e^+e^- \rightarrow \gamma X$	
0.034 ± 0.005 ± 0.006		EDWARDS	99	CLE2	$\Upsilon(2S) \rightarrow \gamma\chi(1P)$	
0.064 ± 0.014 ± 0.016		ALBRECHT	85E	ARG	$e^+e^- \rightarrow \gamma\text{conv. } X$	
0.036 ± 0.008 ± 0.009		NERNST	85	CBAL	$e^+e^- \rightarrow \gamma X$	
0.044 ± 0.023 ± 0.009		HAAS	84	CLEO	$e^+e^- \rightarrow \gamma\text{conv. } X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.035 ± 0.014		KLOPFEN...	83	CUSB	$e^+e^- \rightarrow \gamma X$	

$\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$						$\Gamma_{62}/\Gamma$
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>&lt;59</b>	90	<sup>1</sup> ALBRECHT	89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 5.9	90	<sup>2</sup> ALBRECHT	89	ARG	$\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-$	
<sup>1</sup> Re-evaluated assuming $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$ .						
<sup>2</sup> Includes unknown branching ratio of $f_0(1710) \rightarrow \pi^+\pi^-$ .						

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$						$\Gamma_{63}/\Gamma$
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>&lt;53</b>	90	<sup>1</sup> ALBRECHT	89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$	
<sup>1</sup> Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$ .						

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$						$\Gamma_{64}/\Gamma$
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>&lt;24.1</b>	90	<sup>1</sup> ALBRECHT	89	ARG	$\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-$	
<sup>1</sup> Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$ .						

$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.8	90	<sup>1</sup> ALBRECHT	89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$
------	----	-----------------------	----	-----	---

<sup>1</sup> Includes unknown branching ratio of  $f_J(2220) \rightarrow K^+ K^-$ .

$\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<2.7 $\times 10^{-5}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

$\Gamma(\gamma \chi_{c0})/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<1.0 $\times 10^{-4}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

$\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$   $\Gamma_{68}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<3.6 $\times 10^{-6}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

$\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$   $\Gamma_{69}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<1.5 $\times 10^{-5}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

$\Gamma(\gamma \chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{70}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.8 $\times 10^{-6}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

$\Gamma(\gamma \chi_{c1}(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{71}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<2.4 $\times 10^{-6}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

$\Gamma(\gamma X(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{72}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<2.8 $\times 10^{-6}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

$\Gamma(\gamma \chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{73}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<1.2 $\times 10^{-6}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

$\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{74}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<1.3 $\times 10^{-6}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

$\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$   $\Gamma_{75}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**5.5<sup>+1.1</sup><sub>-0.9</sub> OUR AVERAGE** Error includes scale factor of 1.2.

6.1 <sup>+0.6+0.9</sup> <sub>-0.7-0.6</sub>		29k	FULSOM	18 BELL	$\Upsilon(2S) \rightarrow \gamma X$
---	--	-----	--------	---------	-------------------------------------

3.9 $\pm$ 1.1 <sup>+1.1</sup> <sub>-0.9</sub>		13 $\pm$ 5k	<sup>1</sup> AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
---	--	-------------	---------------------	-----------	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90		LEES	11J BABR	$\Upsilon(2S) \rightarrow X\gamma$
-----	----	--	------	----------	------------------------------------

< 8.4	90		<sup>1</sup> BONVICINI	10 CLEO	$\Upsilon(2S) \rightarrow \gamma X$
-------	----	--	------------------------	---------	-------------------------------------

< 5.1	90		<sup>2</sup> ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$
-------	----	--	---------------------	---------	-------------------------------

<sup>1</sup> Assuming  $\Gamma_{\eta_b(1S)} = 10$  MeV.<sup>2</sup> Superseded by BONVICINI 10. $\Gamma(\gamma\eta_b(1S) \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$   $\Gamma_{76}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<3.7 $\times 10^{-6}$	90	SANDILYA	13 BELL	$\Upsilon(2S) \rightarrow \gamma$ hadrons
-----------------------	----	----------	---------	---

 $\Gamma(\gamma X_{b\bar{b}} \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$   $\Gamma_{77}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

< 4.9	90		SANDILYA	13 BELL	$\Upsilon(2S) \rightarrow \gamma$ hadrons
-------	----	--	----------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

46.2 <sup>+29.7</sup> <sub>-14.2</sub> $\pm$ 10.6		10	<sup>1</sup> DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons
---	--	----	--------------------	----	---

<sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration. $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$   $\Gamma_{78}/\Gamma$   
(1.5 GeV <  $m_X$  < 5.0 GeV)

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.95	95	ROSNER	07A CLEO	$e^+e^- \rightarrow \gamma X$
-------	----	--------	----------	-------------------------------

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$   $\Gamma_{79}/\Gamma$   
(0.3 GeV <  $m_{A^0}$  < 7 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<8 $\times 10^{-5}$	90	<sup>1</sup> LEES	11H BABR	$\Upsilon(2S) \rightarrow \gamma$ hadrons
---------------------	----	-------------------	----------	---

<sup>1</sup> For a narrow scalar or pseudoscalar  $A^0$ , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of  $m_{A^0}$  range from  $1 \times 10^{-6}$  to  $8 \times 10^{-5}$ . $\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{80}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<8.3	90	<sup>1</sup> AUBERT	09Z BABR	$e^+e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$
------	----	---------------------	----------	--

<sup>1</sup> For a narrow scalar or pseudoscalar  $a_1^0$  with mass in the range 212–9300 MeV, excluding  $J/\psi$  and  $\psi(2S)$ . Measured 90% CL limits as a function of  $m_{a_1^0}$  range from 0.26–8.3  $\times 10^{-6}$ .



## LEPTON FAMILY NUMBER (*LF*) VIOLATING MODES

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$					$\Gamma_{81}/\Gamma$
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;3.2</b>	90	LEES	10B	BABR	$e^+ e^- \rightarrow e^\pm \tau^\mp$

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$					$\Gamma_{82}/\Gamma$
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt; 3.3</b>	90	LEES	10B	BABR	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<14.4	95	LOVE	08A	CLEO	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

### $\Upsilon(2S)$ Cross-Particle Branching Ratios

$$B(\Upsilon(2S) \rightarrow \pi^+ \pi^-) \times B(\Upsilon(3S) \rightarrow \Upsilon(2S) X)$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.78±0.02±0.11</b>	906k	LEES	11C	BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$

### $\Upsilon(2S)$ REFERENCES

FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)
SANDILYA	13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
TAMPONI	13	PR D87 011104	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)

ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE $e^+e^-$ Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
GELPHMAN	85	PR D32 2893	D. Gelpman <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>	(DESY, ARGUS Collab.+)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)